Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

Averaging techniques yield reliable a
posteriori finite element error control for
obstacle problems

by

C. Carstensen and S. Bartels

Preprint no.: 2

2001

AVERAGING TECHNIQUES YIELD RELIABLE A POSTERIORI FINITE ELEMENT ERROR CONTROL FOR OBSTACLE PROBLEMS

S. BARTELS AND C. CARSTENSEN

ABSTRACT. The reliability of frequently applied averaging techniques for a posteriori error control has recently been established for a series of finite element methods in the context of second-order partial differential equations. This paper establishes related reliable and efficient a posteriori error estimates for the energy-norm error of an obstacle problem on unstructured grids as a model example for variational inequalities. The surprising main result asserts that the distance of the piecewise constant discrete gradient to *any* continuous piecewise affine approximation is a reliable upper error bound up to known higher order terms, consistency terms, and a multiplicative constant.

1. INTRODUCTION

While a posteriori error control and adaptive mesh design is well established for (elliptic) partial differential equations [AO, BSt, EEHJ, V], their exploitation for variational inequalities started very recently [BSu, CN, LLT, V1, V2]. Amongst the a posteriori error estimation techniques are averaging schemes firstly justified by super-convergence properties on structured grids with symmetry properties. Their recent justification on unstructured grids in [BC, CA, CB, CF1, CF2, CF3, CF4, CF5] raises the question: How can averaging techniques be possibly reliable (i.e., be guaranteed upper bounds) for variational inequalities?

Our mathematical investigations recast this question into the design of a weak approximation operator that is compatible with the obstacle conditions and still enjoys local orthogonality properties to generate higher order terms. Utilising the operator J from [Ca] and its dual J^* this paper provides an affirmative answer for a simple obstacle problem with affine obstacle and studies the nonconforming case.

Given a bounded Lipschitz domain Ω in $\mathbb{R}^d$, $d = 2, 3$, $f \in H^1(\Omega)$, $g \in H^1(\Gamma_N)$, $u_D \in H^1(\Gamma_D)$, and $\chi \in H^1(\Omega)$ such that the closed and convex subset

$$K := \{v \in H^1(\Omega) : v = u_D \text{ on } \Gamma_D, \chi \leq v \text{ almost everywhere in } \Omega\}$$

of $H^1(\Omega)$ is non-void, the obstacle problem under question reads: *Seek $u \in K$ such that*

$$(1.1) \quad (\nabla u; \nabla(u - v)) \leq (f; u - v) + \int_{\Gamma_N} g(u - v) ds \quad \text{for all } v \in K.$$

Here, $(\cdot; \cdot)$ denotes the L^2 -product and Γ_D is a closed subset of $\Gamma := \partial\Omega$ with positive surface measure; $\Gamma_N := \Gamma \setminus \Gamma_D$. It is known [R, GLT, K] that (1.1) has a unique solution. The finite

Date: January 4, 2001.

1991 Mathematics Subject Classification. 65N30, 65R20, 73C50.

Key words and phrases. Adaptive finite element methods, a posteriori error estimates, elliptic variational inequalities, obstacle problems.

element approximation employs a (closed and convex) discrete set K_h (i.e., a subset of a finite-dimensional subspace of $H^1(\Omega)$) and reads: *Seek $u_h \in K_h$ such that*

$$(1.2) \quad (\nabla u_h; \nabla(u_h - v_h)) \leq (f; u_h - v_h) + \int_{\Gamma_N} g(u_h - v_h) ds \quad \text{for all } v_h \in K_h.$$

There exists a unique discrete solution u_h whose error $e := u - u_h$ is in some sense quasi-optimally small; we refer to [F, N] for a priori error estimates and focus on a posteriori estimates in this paper. The choice

$$(1.3) \quad K_h := \{v_h \in \mathcal{S}^1(\mathcal{T}) : v_h = u_{D,h} \text{ on } \Gamma_D, \chi_h \leq v_h \text{ almost everywhere in } \Omega\}$$

can model a conforming (i.e., $K_h \subseteq K$) or non-conforming (i.e., $K_h \not\subseteq K$) discretisation. Here, $\mathcal{S}^1(\mathcal{T})$ is the P_1 -finite element space defined through a regular triangulation $\mathcal{T}$ of Ω into triangles and tetrahedra if $d = 2$ and $d = 3$, respectively, [BSc, Ci]; $\chi_h \in \mathcal{S}^1(\mathcal{T})$ is an approximation to χ , $u_{D,h} \in \mathcal{S}^1(\mathcal{T})|_{\Gamma_D}$ is an approximation to u_D and we assume $K_h \neq \emptyset$.

Our first result employs [BC, CB, Ca, CV] and standard estimates for the proof of

$$(1.4) \quad \|\nabla(u - u_h)\| \lesssim \eta_Z + (\varrho_h; \chi - u_h - w) + \|\nabla w\| + \|h_{\mathcal{T}}^2 \nabla f\| + \|h_{\mathcal{E}}^{3/2} \partial_{\mathcal{E}} g / \partial s\|_{L^2(\Gamma_N)}.$$

Here, $\|\cdot\|$ denotes the $L^2(\Omega)$ -norm, “ $\lesssim$ ” substitutes “ $\leq$ up to a multiplicative mesh-size-independent constant”, w is arbitrary in $H^1(\Omega)$ with $u_h + w \in K$, $w|_{\Gamma_D} = u_D - u_{D,h}$ vanishes at nodes on Γ_D , ϱ_h is a known discrete residual, $h_{\mathcal{T}}$ and $h_{\mathcal{E}}$ are local mesh sizes, and

$$(1.5) \quad \eta_Z := \min\{\|p_h - \nabla u_h\| : p_h \in \mathcal{S}^1(\mathcal{T})^d, p_h \cdot n = g \text{ on } \mathcal{N} \cap \bar{\Gamma}_N\};$$

where n denotes the outer unit normal on Γ_N and $\mathcal{N}$ is the set of nodes in $\mathcal{T}$ ($p_h \cdot n$ interpolates g at all nodes on $\bar{\Gamma}_N$). Consistency is included in the arbitrary choice of w to assess the error in $K_h \neq K$ and $u_{D,h} \neq u_D$; in the absence of contact near the boundary, with $(\cdot)_+ := \max\{\cdot, 0\}$,

$$(1.6) \quad \|\nabla w\| \lesssim \|h_{\mathcal{E}}^{3/2} \partial_{\mathcal{E}}^2 u_D / \partial s^2\|_{L^2(\Gamma_D)} + \|\nabla(\chi - u_h)_+\|.$$

The estimate (1.4) can be recommended for practical error control since $(\varrho_h; \chi - u_h - w)$ can be evaluated. Closer investigations reveal that this term can indeed be replaced by consistency, averaging, and higher order terms. Our main result is a refined version of

$$(1.7) \quad \begin{aligned} \|\nabla(u - u_h)\| &\lesssim \eta_Z + \min_{q_h \in \mathcal{S}^1(\mathcal{T})^d} \|q_h - \nabla(\chi_h - u_h)\| + \|\nabla w\| + \|h_{\mathcal{T}}^2 \nabla f\| + \|h_{\mathcal{T}} f\|_{L^2(\cup \mathcal{T}_D)} \\ &\quad + \|h_{\mathcal{E}}^{3/2} \partial_{\mathcal{E}} g / \partial s\|_{L^2(\Gamma_N)} + \|h_{\mathcal{T}}^{-1}(\chi - \chi_h - w)_-\| + (\|f\| \|(\chi - \chi_h - w)_-\|)^{1/2} \end{aligned}$$

where $(\cdot)_- := \min\{\cdot, 0\}$. The term $\|h_{\mathcal{T}} f\|_{L^2(\cup \mathcal{T}_D)}$ is the L^2 -norm over the shrinking domain $\cup \mathcal{T}_D$, a union of a few layers of elements near Γ_D ; e.g., if $f \in L^\infty(\Omega)$ we have $\|h_{\mathcal{T}} f\|_{L^2(\cup \mathcal{T}_D)} \lesssim \|f\|_{L^\infty(\Omega)} \|h_{\mathcal{E}}^2\|_{L^2(\Gamma_D)}$ and see that this term is of higher order. In case $\chi = \chi_h$ and no contact near the boundary, the estimate reduces to

$$(1.8) \quad \|\nabla(u - u_h)\| \lesssim \eta_Z + \min_{q_h \in \mathcal{S}^1(\mathcal{T})^d} \|q_h - \nabla(\chi_h - u_h)\| + \text{h.o.t.}$$

(h.o.t. denotes higher order terms). The finer estimate of Theorem 3.2 refines (1.7)-(1.8) in the substitution of $\|q_h - \nabla(\chi_h - u_h)\|_{L^2(\Omega)}$ by the refined norm $\|q_h - \nabla(\chi_h - u_h)\|_{L^2(\Omega_s)}$ on a smaller computable region Ω_s around the free boundary of the contact zone. Numerical examples convinced us that this refinement is necessary for efficient approximation and error control.

If the obstacle $\chi = \chi_h$ is globally affine, $\nabla \chi_h = A$ is constant and $q_h = p_h + A$ in (1.8) provides

$$(1.9) \quad \|\nabla(u - u_h)\| \lesssim \eta_Z + \text{h.o.t.}$$

Hence, the averaging estimator η_Z (from the variational equality) is indeed reliable for the obstacle problem up to a multiplicative constant and up to known higher order terms. It is stressed that the averaging estimator η_Z is efficient; the proof is provided by a triangle inequality

$$(1.10) \quad \eta_Z \leq \|\nabla(u_h - u)\| + \min\{\|\nabla u - p_h\| : p_h \in \mathcal{S}^1(\mathcal{T})^d, p_h \cdot n = g \text{ on } \mathcal{N} \cap \bar{\Gamma}_N\};$$

in case u is sufficiently smooth, the minimum in the right-hand side is of higher order.

It appears to us that the reliability of averaging techniques is always related to smooth data (u_D , g , and f) and hence rough obstacles might be excluded from the assumptions; this is seen in our analysis by consistency terms which are not always of higher order and may dominate the error estimate. Consequently, this paper does not focus on coarse approximation of rough data.

The rest of this paper is organised as follows. Preliminaries and notation is introduced in Section 2 where we recall a few results and state some basic estimates. Section 3 is devoted to the a posteriori error estimates and their proofs. Section 4 outlines the numerical realisation with a penalty scheme we employed. Section 5 reports on four examples where the estimate of the error in the energy norm is extremely accurate.

2. PRELIMINARIES

Throughout this paper, $u \in K$ solves (1.1) and $u_h \in K_h$ solves (1.2). The aim is to prove reliability and efficiency of the aforementioned estimators. We let $H_D^1(\Omega) := \{v \in H^1(\Omega) : v = 0 \text{ on } \Gamma_D\}$ and define $\mathcal{S}_D^1(\mathcal{T}) := \mathcal{S}^1(\mathcal{T}) \cap H_D^1(\Omega)$.

Let $(\varphi_z : z \in \mathcal{N})$ be the nodal basis of $\mathcal{S}^1(\mathcal{T})$. Note that $(\varphi_z : z \in \mathcal{N})$ is a partition of unity and the open patches

$$(2.1) \quad \omega_z := \{x \in \Omega : 0 < \varphi_z(x)\}$$

form an open cover $(\omega_z : z \in \mathcal{N})$ of Ω with finite overlap.

Let $\mathcal{K} := \mathcal{N} \setminus \Gamma_D$ denote the set of free nodes and let $\mathcal{E}$ denote the set of all edges ($d = 2$) or faces ($d = 3$) appearing for some T in $\mathcal{T}$. In order to define a weak interpolation operator $J : H_D^1(\Omega) \rightarrow \mathcal{S}_D^1(\mathcal{T})$ we modify $(\varphi_z : z \in \mathcal{K})$ to a partition of unity $(\psi_z : z \in \mathcal{K})$. For each fixed node $z \in \mathcal{N} \setminus \mathcal{K}$, we choose a neighboring node $\zeta(z) \in \mathcal{K}$ and let $\zeta(z) := z$ if $z \in \mathcal{K}$. In this way, we define a partition of $\mathcal{N}$ into $\text{card}(\mathcal{K})$ classes $I(z) := \{\tilde{z} \in \mathcal{N} : \zeta(\tilde{z}) = z\}$, $z \in \mathcal{K}$. For each $z \in \mathcal{K}$ set

$$(2.2) \quad \psi_z := \sum_{\zeta \in I(z)} \varphi_\zeta \quad \text{and} \quad \Omega_z := \{x \in \Omega : 0 < \psi_z(x)\}$$

and notice that $(\psi_z : z \in \mathcal{K})$ is a partition of unity. It is required that Ω_z is connected and that $\psi_z \neq \varphi_z$ implies that $\Gamma_D \cap \partial\Omega_z$ has a positive surface measure.

For $g \in L^1(\Omega)$ define

$$(2.3) \quad Jg := \sum_{z \in \mathcal{K}} g_z \varphi_z \in \mathcal{S}_D^1(\mathcal{T}) \quad \text{where} \quad g_z := (g; \psi_z)/(1; \varphi_z) \in \mathbb{R}.$$

The local mesh-sizes are denoted by $h_{\mathcal{T}}$ and $h_{\mathcal{E}}$ where $h_{\mathcal{T}} \in \mathcal{L}^0(\mathcal{T})$ denotes the element size, $h_{\mathcal{T}}|_T := h_T := \text{diam}(T)$ for $T \in \mathcal{T}$, and the edge size $h_{\mathcal{E}} \in L^\infty(\cup \mathcal{E})$ is defined on the union or skeleton $\cup \mathcal{E}$ of all edges E in $\mathcal{E}$ by $h_{\mathcal{E}}|_E := h_E := \text{diam}(E)$. The patch size $h_z := \text{diam}(\Omega_z)$ is defined for each node $z \in \mathcal{K}$ separately. For $z \in \mathcal{N} \setminus \mathcal{K}$ set $h_z := \text{diam}(\omega_z)$ and for $T \in \mathcal{T}$ let $\omega_T := \cup_{z \in T \cap \mathcal{N}} \Omega_{\zeta(z)}$. Note that the sets of patches $(\omega_T : T \in \mathcal{T})$ and $(\Omega_z : z \in \mathcal{K})$ have a finite overlap.

In the following we write $\|\cdot\|_{p,A}$ instead of $\|\cdot\|_{L^p(A)}$ and $\|\cdot\|$ abbreviates $\|\cdot\|_{2,\Omega}$. Similarly, we denote by $|\cdot|_{1,2,A} := \|\nabla \cdot\|_{2,A}$ the semi-norm in $H^1(A)$ and $|\cdot|_{1,2}$ abbreviates $|\cdot|_{1,2,\Omega}$.

Theorem 2.1 ([Ca, Cl, CV, CB]). *The operator J is H^1 -stable and first-order convergent, i.e.,*

$$(2.4) \quad \|h_{\mathcal{T}}^{-1}(g - Jg)\| + \|h_{\mathcal{E}}^{-1/2}(g - Jg)\|_{2,\Gamma_N} + |g - Jg|_{1,2} \lesssim |g|_{1,2}$$

for $g \in H_D^1(\Omega)$. Moreover, for $f \in L^2(\Omega)$, there holds

$$(2.5) \quad (f; g - Jg) \lesssim |g|_{1,2} \left(\sum_{z \in \mathcal{K}} h_z^2 \min_{f_z \in \mathbb{R}} \|f - f_z\|_{2,\Omega_z}^2 \right)^{1/2}. \quad \square$$

Lemma 2.1. *We have, for all $v \in H_D^1(\Omega)$,*

$$\begin{aligned} (f; v - Jv) + \int_{\Gamma_N} g(v - Jv) dx - (\nabla u_h; \nabla(v - Jv)) \\ \lesssim |v|_{1,2} \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N} + \|h_{\mathcal{T}}^2 \nabla f\|). \end{aligned}$$

Proof. The lemma is, at least implicitly, included in [CB] (and also in [BC, CF1, CF2, CF3, CF4]) and so we merely sketch its proof. From (2.5) we have by Poincaré's inequality

$$(f; v - Jv) \lesssim |v|_{1,2} \|h_{\mathcal{T}}^2 \nabla f\|.$$

An integration by parts of $-(p_h; \nabla(v - Jv))$ and utilising that $\text{div}_{\mathcal{T}} \nabla u_h = 0$ reveals that the last two terms in the left-hand side of the asserted inequality equal

$$(2.6) \quad \begin{aligned} \int_{\Gamma_N} (g - p_h \cdot n)(v - Jv) ds + (p_h - \nabla u_h; \nabla(v - Jv)) + (\text{div}_{\mathcal{T}}(p_h - \nabla u_h); v - Jv) \\ \lesssim |v|_{1,2} \left(\|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N} + \|p_h - \nabla u_h\| + \|h_{\mathcal{T}} \text{div}_{\mathcal{T}}(p_h - \nabla u_h)\| \right) \end{aligned}$$

by (2.4) and Cauchy inequalities. This and a $\mathcal{T}$ -elementwise inverse estimate of the form $h_T \|\text{div}_{\mathcal{T}}(p_h - \nabla u_h)\|_{2,T} \lesssim \|p_h - \nabla u_h\|_{2,T}$ conclude the proof. $\square$

Lemma 2.2 ([BCD]). *Assume $u_D \in H^1(\Gamma_D) \cap C(\Gamma_D)$, $u_D|_E \in H^2(E)$ for all $E \in \mathcal{E}$ such that $E \subseteq \Gamma_D$, and let $\partial_{\xi}^2 u_D / \partial s^2$ denote the edgewise second derivative of u_D along Γ_D . Suppose $u_{D,h}$ is the nodal interpolant of u_D , i.e., $u_{D,h}(z) = u_D(z)$ for all $z \in \mathcal{N} \cap \Gamma_D$. Then there exists $w_D \in H^1(\Omega)$ such that $w_D|_{\Gamma_D} = u_D - u_{D,h}$, $\text{supp } w_D \subseteq \bigcup_{T \in \mathcal{T}, T \cap \Gamma_D \neq \emptyset} T$,*

$$(2.7) \quad \|w_D\|_{\infty} = \|u_D - u_{D,h}\|_{\infty, \Gamma_D}, \quad \text{and} \quad |w_D|_{1,2} \lesssim \|h_{\mathcal{E}}^{3/2} \partial_{\xi}^2 u_D / \partial s^2\|_{2, \Gamma_D}. \quad \square$$

Definition 2.1. Define

$$\mathcal{T}_D := \{T \in \mathcal{T} : T \cap \Gamma_D \neq \emptyset\} \quad \text{and} \quad \mathcal{T}_c := \{T \in \mathcal{T} \setminus \mathcal{T}_D : (\chi_h - u_h)|_T = 0\}.$$

The following lemma shows (1.6) and estimates the terms which include w in (1.7).

Lemma 2.3. *Suppose that u_D satisfies the conditions of Lemma 2.2, that $\chi|_{\Gamma_D} \leq u_{D,h}$, and that $(\chi - u_h)_- \leq w_D$ in $\bigcup \mathcal{T}_D$ with w_D from Lemma 2.2. Then we have*

$$\begin{aligned} \min_{\substack{w \in H^1(\Omega) \\ u_h + w \in K}} |w|_{1,2} &\lesssim \|h_\varepsilon^{3/2} \partial_\varepsilon^2 u_D / \partial s^2\|_{2,\Gamma_D} + |(\chi - u_h)_+|_{1,2} \quad \text{and} \\ \min_{\substack{w \in H^1(\Omega) \\ u_h + w \in K}} &\left(|w|_{1,2}^2 + \sum_{T \in \mathcal{T} \setminus \mathcal{T}_D} \|h_T^{-1} (\chi - \chi_h - w)_-\|_{2,T}^2 + \sum_{T \in \mathcal{T}_c} \|f\|_{2,\omega_T} \|(\chi - \chi_h - w)_-\|_{2,T} \right) \\ &\lesssim \|h_\varepsilon^{3/2} \partial_\varepsilon^2 u_D / \partial s^2\|_{2,\Gamma_D}^2 + |(\chi - u_h)_+|_{1,2}^2 + \sum_{T \in \mathcal{T} \setminus \mathcal{T}_D} \|h_T^{-1} (\chi - \chi_h)_-\|_{2,T}^2 \\ &\quad + \sum_{T \in \mathcal{T}_c} \|f\|_{2,\omega_T} \|(\chi - \chi_h)_-\|_{2,T}. \end{aligned}$$

Proof. Set $w := (\chi - u_h)_+ + w_D$ and notice $u_h + w \in K$. Then $|w|_{1,2} \leq |(\chi - u_h)_+|_{1,2} + |w_D|_{1,2}$. Utilising $w_D|_T = 0$ and $\chi_h \leq u_h$ on each $T \in \mathcal{T} \setminus \mathcal{T}_D$ we have on each $T \in \mathcal{T} \setminus \mathcal{T}_D$

$$\|(\chi - \chi_h - w)_-\|_{2,T} = \|(\chi - \chi_h - (\chi - u_h)_+)_-\|_{2,T} \leq \|(\chi - \chi_h)_-\|_{2,T}.$$

Then, Lemma 2.2 proves the assertions. $\square$

Remark 2.1. Since $\|w_D\|_\infty = \|u_D - u_{D,h}\|_{\infty,\Gamma_D}$ by Lemma 2.2, the assumption $(\chi - u_h)_- \leq -\|u_D - u_{D,h}\|_{\infty,\Gamma_D}$ in $\bigcup \mathcal{T}_D$ implies $(\chi - u_h)_- \leq w_D$ in $\bigcup \mathcal{T}_D$.

Lemma 2.4 ([BC, CB]). *Let $g|_E \in H^1(E) \cap C(E)$ for all $E \in \mathcal{E}$ such that $E \subseteq \bar{\Gamma}_N$ and, for each node $z \in \mathcal{N} \cap \bar{\Gamma}_N$ where the outer unit normal n on Γ_N is continuous (hence constant in a neighbourhood of z as Γ_N is a polygon), let g be continuous. Assume that the set*

$$\mathcal{S}_N^1(\mathcal{T}, g) := \{p_h \in \mathcal{S}^1(\mathcal{T})^d : \forall E \in \mathcal{E}, z \in E \subseteq \bar{\Gamma}_N, p_h(z) \cdot n_E = g(z)\}$$

is non-void. Then $(\partial_\varepsilon g / \partial s)$ denotes the edgewise surface gradient of g on Γ_N)

$$\begin{aligned} \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} &\left(\|\nabla u_h - p_h\| + \|h_\varepsilon^{1/2} (g - p_h \cdot n)\|_{2,\Gamma_N} \right) \\ &\lesssim \min_{q_h \in \mathcal{S}_N^1(\mathcal{T}, g)} \|\nabla u_h - q_h\| + \|h_\varepsilon^{3/2} \partial_\varepsilon g / \partial s\|_{2,\Gamma_N}. \quad \square \end{aligned}$$

Remark 2.2. For $d = 2$ the conditions on g in Lemma 2.4 suffice for $\mathcal{S}_N^1(\mathcal{T}, g) \neq \emptyset$ [CB].

Definition 2.2. Define $\varrho \in (H_D^1(\Omega))^*$ and $\varrho_h \in \mathcal{S}^1(\mathcal{T})$, for $v \in H_D^1(\Omega)$, by

$$(2.8) \quad \varrho(v) := (f; v) + \int_{\Gamma_N} g v \, ds - (\nabla u; \nabla v),$$

$$(2.9) \quad \varrho_h := \sum_{z \in \mathcal{K}} \left((f; \varphi_z) + \int_{\Gamma_N} g \varphi_z \, ds - (\nabla u_h; \nabla \varphi_z) \right) \psi_z / (1; \varphi_z).$$

Remark 2.3. Note that $0 \leq \varrho(e - w)$ for $w \in H^1(\Omega)$ satisfying $w|_{\Gamma_D} = u_D - u_{D,h}$ and $\chi - u_h \leq w$ (since $u_h + w \in K$). If $u_h \in K$ we may choose $w = 0$. If not, let, e.g., $P_K u_h$ be the projection of u_h onto K with respect to $|\cdot|_{1,2}$ and $w := P_K u_h - u_h$ minimises $|w|_{1,2}$ among all w with $u_h + w \in K$.

Lemma 2.5. *We have, for all $w \in H^1(\Omega)$ satisfying $w|_{\Gamma_D} = u_D - u_{D,h}$,*

$$(2.10) \quad \begin{aligned} \frac{1}{2}|e - w|_{1,2}^2 + \frac{1}{2}|e|_{1,2}^2 &= (f; e - w - J(e - w)) - (\nabla u_h; \nabla(e - w - J(e - w))) \\ &+ \int_{\Gamma_N} g(e - w - J(e - w)) ds + \frac{1}{2}|w|_{1,2}^2 + (\varrho_h; e - w) - \varrho(e - w). \end{aligned}$$

Proof. Note that $e - w \in H_D^1(\Omega)$. The definition of $J(e - w)$ yields, e.g.,

$$\sum_{z \in \mathcal{K}} (\nabla u_h; \nabla \varphi_z)(\psi_z; e - w) / (1; \varphi_z) = (\nabla u_h; \nabla J(e - w))$$

and eventually leads to

$$(\varrho_h; e - w) = (f; J(e - w)) - (\nabla u_h; \nabla J(e - w)) + \int_{\Gamma_N} gJ(e - w) ds.$$

This and some elementary calculations show

$$\begin{aligned} \varrho(e - w) - (\varrho_h; e - w) &= (f; e - w - J(e - w)) + \int_{\Gamma_N} g(e - w - J(e - w)) ds \\ &- (\nabla u_h; \nabla(e - w)) + (\nabla u_h; \nabla J(e - w)) = (f; e - w - J(e - w)) \\ &+ \int_{\Gamma_N} g(e - w - J(e - w)) ds - (\nabla e; \nabla(e - w)) - (\nabla u_h; \nabla(e - w - J(e - w))). \end{aligned}$$

Since $2(\nabla e; \nabla(e - w)) = |e - w|_{1,2}^2 + |e|_{1,2}^2 - |w|_{1,2}^2$ we deduce (2.10). $\square$

Our motivation for the definition of ϱ_h is that its nodal values reflect Kuhn-Tucker conditions.

Lemma 2.6. *We have $\varrho_h \leq 0 \leq u_h - \chi_h$ almost everywhere in Ω and, for $z \in \mathcal{K}$,*

$$\varrho_h(z)(\chi_h(z) - u_h(z)) = 0.$$

Proof. Given $z \in \mathcal{K}$ and a real number w define $v_h \in \mathcal{S}^1(\mathcal{T})$ by $v_h(z) := w$ and $v_h(\zeta) = u_h(\zeta)$ for $\zeta \in \mathcal{N} \setminus \{z\}$. If $\chi_h(z) \leq w$ we have $v_h \in K_h$ and calculate with (1.2)

$$\begin{aligned} (u_h(z) - w)(\nabla u_h; \nabla \varphi_z) &= (\nabla u_h; \nabla(u_h - v_h)) \leq (f; u_h - v_h) + \int_{\Gamma_N} g(u_h - v_h) ds \\ &= (u_h(z) - w)((f; \varphi_z) + \int_{\Gamma_N} g\varphi_z ds). \end{aligned}$$

According to (2.9) this gives (after a division by $(1; \varphi_z) > 0$)

$$0 \leq (u_h(z) - w)\varrho_h(z).$$

A discussion of $w \in \mathbb{R}$ under the restriction $\chi_h(z) \leq w$ yields the assertions. $\square$

3. A POSTERIORI ESTIMATES

The combination of the next result with Lemma 2.4 provides a proof of (1.4).

Theorem 3.1 (A posteriori estimate I). *If $w \in H^1(\Omega)$ is such that $u_h + w \in K$, i.e., $w|_{\Gamma_D} = u_D - u_{D,h}$ and $\chi - u_h \leq w$, then*

$$|e - w|_{1,2} + |e|_{1,2} \lesssim \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N}) \\ + \|h_{\mathcal{T}}^2 \nabla f\| + |w|_{1,2} + (\varrho_h; \chi - u_h - w).$$

Proof. Since $u_h + w \in K$ we have $\varrho(e - w) \geq 0$, cf. Remark 2.3. Moreover, $\varrho_h \leq 0 \leq u - \chi$ almost everywhere in Ω by Lemma 2.6 so that $(\varrho_h; u - \chi) \leq 0$ and hence

$$(\varrho_h; e - w) - \varrho(e - w) \leq (\varrho_h; e - w) = (\varrho_h; u - \chi) + (\varrho_h; \chi - u_h - w) \leq (\varrho_h; \chi - u_h - w).$$

Utilising this estimate and Lemma 2.1 in (2.10) we deduce the assertion. $\square$

The following lemmas are needed to obtain other bounds for $(\varrho_h; e - w)$.

Lemma 3.1. *Let $z \in \mathcal{N}$ be either an interior point of Ω or suppose that each open half-space with boundary point z has a non-void intersection with Ω . Suppose $T \in \mathcal{T}$ is such that $z \in \overline{\omega}_T$ and set $\tilde{\Omega}_z := \Omega_z \cup \omega_T$. Let $w_h \in \mathcal{S}^1(\mathcal{T})$ satisfy $w_h(z) = 0$ and $0 \leq w_h$ on $\tilde{\Omega}_z$. Then,*

$$(3.1) \quad \|w_h\|_{2,\tilde{\Omega}_z} \lesssim h_z \min_{q_z \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_z})^d} \|\nabla w_h - q_z\|_{2,\tilde{\Omega}_z}.$$

Proof. The left- and right-hand side of (3.1) define semi-norms $\|\cdot\|_l$ and $\|\cdot\|_r$, respectively, on $\mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_z})$. We claim that $\|w_h\|_r = 0$ implies $\|w_h\|_l = 0$ for all $w_h \in \mathcal{S}^1(\mathcal{T})$ with $w_h(z) = 0 \leq w_h|_{\tilde{\Omega}_z}$. Indeed, if ∇w_h equals some $q_z \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_z})^d$ it is $(\mathcal{T}|_{\tilde{\Omega}_z})$ -piecewise constant and continuous, whence w_h is affine on $\tilde{\Omega}_z$. Since $w_h(z) = 0$ we obtain that $w_h(x)$ equals $\alpha n \cdot (x - z)$ for all $x \in \tilde{\Omega}_z$ and some $n \in \mathbb{R}^d$, $|n| = 1$, and some $\alpha \in \mathbb{R}$. Let $H := \{y \in \mathbb{R}^d : m \cdot (y - z) < 0\}$ intersect with $\tilde{\Omega}_z$ ($H \cap \tilde{\Omega}_z \neq \emptyset$ is obvious for $z \in \Omega$ and assumed for $z \in \mathcal{N} \setminus \Omega$). For $x \in H \cap \tilde{\Omega}_z$,

$$(3.2) \quad 0 \leq w_h(x) = \alpha n \cdot (x - z) \text{ and } m \cdot (x - z) < 0.$$

For $m = +n$, (3.2) implies $\alpha \leq 0$ and for $m = -n$, (3.2) yields $0 \leq \alpha$. Together, $\alpha = 0$, i.e., $w_h = 0$ and so $\|w_h\|_l = 0$. Since $\|\cdot\|_l$ and $\|\cdot\|_r$ are norms on the finite-dimensional affine space $\{w_h \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_z}) : w_h(z) = 0, 0 \leq w_h|_{\tilde{\Omega}_z}\}$, they are equivalent. The constant $C > 0$ in $\|\cdot\|_l \leq C \|\cdot\|_r$ depends on $\mathcal{T}|_{\tilde{\Omega}_z}$. A scaling argument concludes the proof. $\square$

Remark 3.1. If $z \in \mathcal{N}$ is a boundary point of Ω and $\{x \in \Omega : |x - z| < \varepsilon\}$ is convex for some $\varepsilon > 0$ then z does not satisfy the condition of Lemma 3.1. Convex corners may yield unexpected difficulties for higher approximation [NW].

The next result shows that ϱ_h can be controlled by averaging terms.

Lemma 3.2. *We have, for $T \in \mathcal{T}$,*

$$h_T \|\varrho_h\|_{2,T} \lesssim h_T \|f\|_{2,\omega_T} + \min_{q_T \in \mathcal{S}^1(\mathcal{T}|_{\omega_T})^d} (\|\nabla u_h - q_T\|_{2,\omega_T} + h_T^{1/2} \|(g - q_T \cdot n)\|_{2,\Gamma_N \cap \partial\omega_T}), \\ h_T^2 |\varrho_h|_{1,2,T} \lesssim h_T^2 \|f\|_{1,2,\omega_T} + \min_{q_T \in \mathcal{S}^1(\mathcal{T}|_{\omega_T})^d} (\|\nabla u_h - q_T\|_{2,\omega_T} + h_T^{1/2} \|(g - q_T \cdot n)\|_{2,\Gamma_N \cap \partial\omega_T}).$$

Proof. Set $J^*f := \sum_{z \in \mathcal{K}} (f; \varphi_z) / (1; \varphi_z) \psi_z$ and note that J^*f is the first summand in the definition of ϱ_h in (2.9). We have $\|J^*f\|_{2,T} \lesssim \|f\|_{2,\omega_T}$ and $|J^*f|_{1,2,T} \lesssim |f|_{1,2,\omega_T}$ for $T \in \mathcal{T}$, cf. [Ci, CV, CB]. Note also that $h_T^d \lesssim (1, \varphi_z) \lesssim h_T^d$ and $|\psi_z|_{1,2,T} \lesssim h_T^{d/2-1}$, $\|\psi_z\|_{2,T} \lesssim h_T^{d/2}$ for all $T \in \mathcal{T}$ with $T \subseteq \overline{\Omega}_z$. Using this in (2.9) we deduce

$$(3.3) \quad \begin{aligned} \|\varrho_h\|_{2,T} &\lesssim \|f\|_{2,\omega_T} + \sum_{z \in \mathcal{K}, T \subseteq \overline{\Omega}_z} h_T^{-d/2} |(\nabla u_h; \nabla \varphi_z) - \int_{\Gamma_N} g \varphi_z ds|, \\ |\varrho_h|_{1,2,T} &\lesssim |f|_{1,2,\omega_T} + \sum_{z \in \mathcal{K}, T \subseteq \overline{\Omega}_z} h_T^{-d/2-1} |(\nabla u_h; \nabla \varphi_z) - \int_{\Gamma_N} g \varphi_z ds|. \end{aligned}$$

Let q_T be an element of $\mathcal{S}^1(\mathcal{T}|_{\omega_T})^d$. An elementwise inverse estimate shows

$$h_z \|\operatorname{div}_{\mathcal{T}}(q_T - \nabla u_h)\|_{2,\omega_T} \lesssim \|\nabla u_h - q_T\|_{2,\omega_T}.$$

This, an integration by parts, $\operatorname{div}_{\mathcal{T}} \nabla u_h = 0$, $|\varphi_z|_{1,2} \lesssim h_z^{d/2-1}$, $\|\varphi_z\| \lesssim h_z^{d/2}$, and noting that for $z \in T \cap \mathcal{K}$ there holds $\omega_z \subseteq \omega_T$ lead to

$$(\nabla u_h; \nabla \varphi_z) - \int_{\Gamma_N} \varphi_z q_T \cdot n ds = (\nabla u_h - q_T; \nabla \varphi_z) - (\operatorname{div}_{\mathcal{T}} q_T; \varphi_z) \lesssim h_T^{d/2-1} \|\nabla u_h - q_T\|_{2,\omega_T}.$$

This, the fact that each element T belongs to a finite number of patches $\overline{\Omega}_z$ only, (3.3), and $\int_{\partial\omega_T \cap \Gamma_N} \varphi_z (g - q_T \cdot n) ds \lesssim h_z^{d/2-1} \|h_z^{1/2} (g - q_T \cdot n)\|_{2,\Gamma_N \cap \partial\omega_T}$ conclude the proof of the lemma. $\square$

Definition 3.1. With each $T \in \mathcal{T}_i$,

$$\mathcal{T}_i := \{T \in \mathcal{T} : T \cap \Gamma_D = \emptyset, \exists x, y \in \mathcal{K} \cap \overline{\omega}_T, \chi_h(x) = u_h(x), \chi_h(y) < u_h(y)\},$$

we associate some $z_T \in \mathcal{K} \cap \overline{\omega}_T$ such that $\chi_h(z_T) = u_h(z_T)$ and set $\tilde{\Omega}_{z_T} := \Omega_{z_T} \cup \omega_T$,

$$\mathcal{T}_s := \{T \in \mathcal{T} : \exists K \in \mathcal{T}_i, T \subseteq \tilde{\Omega}_{z_K}\}, \quad \text{and} \quad \Omega_s := \bigcup_{T \in \mathcal{T}_s} T.$$

Remark 3.2. For each $T \in \mathcal{T}_i$ we preferably choose $z_T \in \Omega$ (i.e., $z_T \notin \partial\Omega$ if possible). This allows us to impose the condition of Lemma 3.1 in as few as possible nodes on the boundary.

Remark 3.3. The region Ω_s may be regarded as a layer between the discrete contact zone and the discrete non-contact zone.

Lemma 3.3. Assume that for each $T \in \mathcal{T}_i$ for which $z_T \in \mathcal{K} \cap \Gamma_N$ the intersection of Ω with any open half-space with boundary point z_T is non-void. For all $w \in H^1(\Omega)$ satisfying $w|_{\Gamma_D} = u_D - u_{D,h}$ and $\chi - u_h \leq w$, we have

$$\begin{aligned} (\varrho_h; e - w) &\lesssim \sum_{T \in \mathcal{T}_D} h_T \|\varrho_h\|_{2,T} |e - w|_{1,2,\omega_T} + \sum_{T \in \mathcal{T}_i} \min_{q_z \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_{z_T}})^d} \|\nabla(\chi_h - u_h) - q_z\|_{2,\tilde{\Omega}_{z_T}}^2 \\ &\quad + \sum_{T \in \mathcal{T} \setminus \mathcal{T}_D} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T}^2 + \sum_{T \in \mathcal{T}_e} h_T \|\varrho_h\|_{2,T} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T} + \|h_{\mathcal{T}}^2 \nabla \varrho_h\|^2. \end{aligned}$$

Proof. Since $(\varrho_h; e - w) = \sum_{T \in \mathcal{T}} \int_T \varrho_h(e - w) dx$ we may estimate the contribution from each $T \in \mathcal{T}$ separately. In the first case we assume $T \in \mathcal{T}_D$. Since $e - w = 0$ on Γ_D , Friedrichs' inequality implies

$$(3.4) \quad \int_T \varrho_h(e - w) dx \lesssim h_T \|\varrho_h\|_{2,T} |e - w|_{1,2,\omega_T}.$$

In the second case we assume $T \in \mathcal{T}$, $T \cap \Gamma_D = \emptyset$, and $\chi_h|_{\omega_T} < u_h|_{\omega_T}$. Lemma 2.6 guarantees $\varrho_h|_T = 0$ and so

$$(3.5) \quad \int_T \varrho_h(e - w) dx = 0.$$

In the third case we assume $T \in \mathcal{T}_i$ and so there exist $y, z_T \in \mathcal{K} \cap \overline{\omega}_T$ such that $\chi_h(z_T) = u_h(z_T)$ and $\chi_h(y) < u_h(y)$. The conditions of Lemma 3.1 are satisfied for z_T by assumption. Since $\chi_h \leq u_h$, $\chi \leq u$,

$$(\chi - \chi_h - w)_- + \chi_h - u_h \leq (\chi - u_h - w)_- \leq e - w.$$

Because of $\varrho_h \leq 0$, this leads to

$$(3.6) \quad \int_T \varrho_h(e - w) dx \leq \int_T \varrho_h(\chi - \chi_h - w)_- dx + \int_T \varrho_h(\chi_h - u_h) dx.$$

Owing to Lemma 2.6 we have $\varrho_h(y) = 0$ and so $\|\varrho_h\|_{2,T} \lesssim h_T |\varrho_h|_{1,2,\omega_T}$ by a discrete Friedrichs' inequality. This, (3.6), and Lemma 3.1 show

$$(3.7) \quad \begin{aligned} \int_T \varrho_h(e - w) dx &\lesssim h_T^2 |\varrho_h|_{1,2,\omega_T} \\ &\times \left(\|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T} + \min_{q_z \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_{z_T}})^d} \|\nabla(\chi_h - u_h) - q_z\|_{2,\tilde{\Omega}_{z_T}} \right). \end{aligned}$$

In the remaining fourth case we assume $T \in \mathcal{T}_c$ and obtain with (3.6)

$$(3.8) \quad \int_T \varrho_h(e - w) dx \lesssim h_T \|\varrho_h\|_{2,T} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T}.$$

The summation of (3.4), (3.5), (3.7), and (3.8) verifies the assertion. $\square$

The combination of the next result with Lemma 2.4 provides the proof of (1.7) and so specifies to the reliability of all averaging techniques for affine obstacles.

Theorem 3.2 (A posteriori estimate II). *Assume that for each $T \in \mathcal{T}_i$ such that $z_T \in \mathcal{K} \cap \Gamma_N$ the intersection of Ω with any open half-space with boundary point z_T is non-void. For all $w \in H^1(\Omega)$ satisfying $w|_{\Gamma_D} = u_D - u_{D,h}$ and $\chi - u_h \leq w$, we have*

$$\begin{aligned} |e - w|_{1,2} + |e|_{1,2} &\lesssim \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N}) + \min_{q_h \in \mathcal{S}^1(\mathcal{T}_s)^d} \|\nabla(\chi_h - u_h) - q_h\|_{2,\Omega_s} \\ &+ \left(\sum_{T \in \mathcal{T}_c} \|f\|_{2,\omega_T} \|(\chi - \chi_h - w)_-\|_{2,T} \right)^{1/2} + |w|_{1,2} + \|h_{\mathcal{T}}^2 \nabla f\| \\ &+ \left(\sum_{T \in \mathcal{T} \setminus \mathcal{T}_D} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T}^2 \right)^{1/2} + \left(\sum_{T \in \mathcal{T}_D} h_T^2 \|f\|_{2,\omega_T}^2 \right)^{1/2}. \end{aligned}$$

Proof. As in the proof of Theorem 3.1 we have

$$|e - w|_{1,2} + |e|_{1,2} \lesssim \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N}) + \|h_{\mathcal{T}}^2 \nabla f\| + |w|_{1,2} + (\varrho_h; e - w).$$

Employing Lemma 3.3 and absorbing $|e - w|_{1,2}$ we have

$$(3.9) \quad |e - w|_{1,2}^2 + |e|_{1,2}^2 \lesssim \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N})^2 + \|h_{\mathcal{T}}^2 \nabla f\|^2 + |w|_{1,2}^2 + \|h_{\mathcal{T}}^2 \nabla \varrho_h\|^2 + \sum_{T \in \mathcal{T}_D} h_T^2 \|\varrho_h\|_{2,T}^2 + \sum_{T \in \mathcal{T} \setminus \mathcal{T}_D} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T}^2 + \sum_{T \in \mathcal{T}_i} \min_{q_z \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_{z_T}})^d} \|\nabla(\chi_h - u_h) - q_z\|_{2,\tilde{\Omega}_{z_T}}^2 + \sum_{T \in \mathcal{T}_c} h_T \|\varrho_h\|_{2,T} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T}.$$

Lemma 3.2 shows

$$(3.10) \quad \|h_{\mathcal{T}}^2 \nabla \varrho_h\|^2 + \sum_{T \in \mathcal{T}_D} h_T^2 \|\varrho_h\|_{2,T}^2 + \sum_{T \in \mathcal{T}_c} h_T \|\varrho_h\|_{2,T} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T} \lesssim \|h_{\mathcal{T}}^2 \nabla f\|^2 + \sum_{T \in \mathcal{T}} \min_{q_T \in \mathcal{S}^1(\mathcal{T}|_{\omega_T})^d} (\|\nabla u_h - q_T\|_{2,\omega_T} + h_T^{1/2} \|(g - q_T \cdot n)\|_{2,\Gamma_N \cap \partial\omega_T})^2 + \sum_{T \in \mathcal{T} \setminus \mathcal{T}_D} \|h_T^{-1}(\chi - \chi_h - w)_-\|_{2,T}^2 + \sum_{T \in \mathcal{T}_c} \|h_T f\|_{2,\omega_T} \|h_T^{-1}(\chi - \chi_h - w)_-\| + \sum_{T \in \mathcal{T}_D} h_T^2 \|f\|_{2,\omega_T}^2.$$

Let $\tilde{p}_h \in \mathcal{S}^1(\mathcal{T})^d$ denote the minimiser of

$$\min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N}).$$

Since $\tilde{p}_h|_{\omega_T} \in \mathcal{S}^1(\mathcal{T}|_{\omega_T})^d$ for all $T \in \mathcal{T}$ and since the patches ω_T have a finite overlap we have

$$(3.11) \quad \sum_{T \in \mathcal{T}} \min_{q_T \in \mathcal{S}^1(\mathcal{T}|_{\omega_T})^d} (\|\nabla u_h - q_T\|_{2,\omega_T} + h_T^{1/2} \|(g - q_T \cdot n)\|_{2,\Gamma_N \cap \partial\omega_T})^2 \lesssim \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N})^2.$$

A similar argument and the definition of $\mathcal{T}_s$ show

$$(3.12) \quad \sum_{T \in \mathcal{T}_i} \min_{q_z \in \mathcal{S}^1(\mathcal{T}|_{\tilde{\Omega}_{z_T}})^d} \|\nabla(\chi_h - u_h) - q_z\|_{2,\tilde{\Omega}_{z_T}}^2 \lesssim \min_{q_h \in \mathcal{S}^1(\mathcal{T}_s)^d} \|\nabla(\chi_h - u_h) - q_h\|_{2,\Omega_s}^2.$$

The combination of (3.9)-(3.12) proves the theorem. $\square$

Remark 3.4. Two applications of the triangle inequality indicate efficiency of the error estimate of Theorem 3.2 in case $\chi_h = \chi$,

$$\min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u_h - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N}) + \min_{q_h \in \mathcal{S}^1(\mathcal{T}_s)^d} \|\nabla(u_h - \chi_h) - q_h\|_{2,\Omega_s} \leq |e|_{1,2} + |e|_{1,2,\Omega_s} + |\chi - \chi_h|_{1,2,\Omega_s} + 2 \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} (\|\nabla u - p_h\| + \|h_{\mathcal{E}}^{1/2}(g - p_h \cdot n)\|_{2,\Gamma_N}) + \min_{q_h \in \mathcal{S}^1(\mathcal{T}_s)^d} \|\nabla \chi - q_h\|_{2,\Omega_s}. \quad \square$$

4. NUMERICAL REALISATION VIA PENALISATION

In the examples presented in the subsequent section we have $\chi_h = \chi$, $\Gamma_N = \emptyset$, $\chi|_{\Gamma_D} \leq u_{D,h}$, and $f \in H^2(\Omega)$. Then the error estimate of Theorem 3.2 reduces to

$$(4.1) \quad |e|_{1,2} \lesssim \min_{p_h \in \mathcal{S}^1(\mathcal{T})^d} \|\nabla u_h - p_h\| + \min_{q_h \in \mathcal{S}^1(\mathcal{T}_s)^d} \|\nabla(u_h - \chi_h) - q_h\|_{2,\Omega_s} + \text{h.o.t.},$$

where h.o.t. denotes higher order contributions. In the numerical experiments we do not compute the minima in (4.1) but calculate an upper bound for the first two terms by applying the operator $\mathcal{A} : L^2(\Omega)^2 \rightarrow \mathcal{S}_N^1(\mathcal{T}, g)$ and $\mathcal{B} : L^2(\Omega)^2 \rightarrow \mathcal{S}^1(\mathcal{T})^2$ of [CB], defined for $p \in L^2(\Omega)^2$ and $\Gamma_N = \emptyset$ by

$$(4.2) \quad \mathcal{A}p = \mathcal{B}p := \sum_{z \in \mathcal{N}} p_z \varphi_z$$

with $p_z := \frac{1}{|\omega_z|} \int_{\omega_z} p \, dx \in \mathbb{R}^2$ for $z \in \mathcal{N}$; i.e., we calculate

$$\|\nabla u_h - \mathcal{A}\nabla u_h\| + \|\nabla(u_h - \chi_h) - \mathcal{B}\nabla(u_h - \chi_h)\|_{2,\Omega_s}.$$

We refer to [CB] for a definition of $\mathcal{A}$ in case $\Gamma_N \neq \emptyset$ where $g - (\mathcal{A}\nabla u_h) \cdot n$ vanishes at all nodes $z \in \mathcal{N} \cap \bar{\Gamma}_N$.

Our numerical approximations are obtained utilising a penalisation method which reads: *Given $\varepsilon > 0$, seek $u_\varepsilon \in \{v \in H^1(\Omega) : v = u_D \text{ on } \Gamma_D\}$ such that*

$$(4.3) \quad (\nabla u_\varepsilon; \nabla v) = (f; v) + \int_{\Gamma_N} g v \, ds - \frac{1}{\varepsilon}((u_\varepsilon - \chi)_-; v) \quad \text{for all } v \in H_D^1(\Omega).$$

It is not difficult to show that (4.3) has a unique solution $u_\varepsilon \in H^1(\Omega)$ with $u_\varepsilon = u_D$ on Γ_D . Moreover, if the solution u of problem (1.1) satisfies $u \in H^2(\Omega)$ we have

$$|u - u_\varepsilon|_{1,2} \leq \varepsilon^{1/2} \|f + \Delta u\|.$$

The discrete version of (4.3) reads: *Given $\varepsilon > 0$, seek $u_{\varepsilon,h} \in \{v_h \in \mathcal{S}^1(\mathcal{T}) : v_h = u_{D,h} \text{ on } \Gamma_D\}$ such that*

$$(4.4) \quad (\nabla u_{\varepsilon,h}; \nabla v_h) = (f; v_h) + \int_{\Gamma_N} g v_h \, ds - \frac{1}{\varepsilon}((u_{\varepsilon,h} - \chi_h)_-; v_h) \quad \text{for all } v_h \in \mathcal{S}_D^1(\mathcal{T}).$$

We solved the nonlinear equation (4.4) with a Newton-Raphston scheme (without damping). The implementation was performed in Matlab in the spirit of [ACF] with a direct solution of linear systems of equations.

We stress that we do not solve (1.2) but rather an approximation to it and we use the penalisation (4.3) even if we know $u \notin H^2(\Omega)$.

The following adaptive algorithm generates all the sequences of meshes $\mathcal{T}_0, \mathcal{T}_1, \mathcal{T}_2, \dots$ in this paper which are uniform for $\Theta = 0$ or adapted for $\Theta = 1/2$ in (4.5). For details on the red-blue-green-refinements in the algorithm we refer to [V].

- Algorithm* (A_Θ). (a) Start with a coarse mesh $\mathcal{T}_0$, $k = 0$.
 (b) Set $\varepsilon := 1/N$ where N is the number of degrees of freedom of the triangulation $\mathcal{T}_k$ and compute the discrete solution $u_{\varepsilon,h}$ of (4.4) on the actual mesh $\mathcal{T}_k$.
 (c) Define

$$\begin{aligned}\mathcal{M} &:= \{z \in \mathcal{K} : u_{\varepsilon,h} \leq \chi_h(z), \exists T \in \mathcal{T}_k \exists y \in \mathcal{N} \cap T, \chi_h(y) < u_{\varepsilon,h}(y)\}, \\ \mathcal{T}_{s,\varepsilon} &:= \{T \in \mathcal{T}_k : \exists z \in \mathcal{M}, T \subseteq \overline{\omega}_z\}.\end{aligned}$$

For $T \in \mathcal{T}_k$ compute the refinement indicator

$$\eta_{Z,T} := \begin{cases} \|\nabla u_{\varepsilon,h} - \mathcal{A}\nabla u_{\varepsilon,h}\|_{2,T} & \text{if } T \notin \mathcal{T}_{s,\varepsilon}, \\ \frac{1}{2}(\|\nabla u_{\varepsilon,h} - \mathcal{A}\nabla u_{\varepsilon,h}\|_{2,T} + \|\nabla(u_{\varepsilon,h} - \chi_h) - \mathcal{B}\nabla(u_{\varepsilon,h} - \chi_h)\|_{2,T}) & \text{if } T \in \mathcal{T}_{s,\varepsilon}, \end{cases}$$

for the energy error $e_N := \|\nabla(u - u_{\varepsilon,h})\|_{2,\Omega}$ and compute its estimator $\eta_N := (\sum_{T \in \mathcal{T}} \eta_{Z,T}^2)^{1/2}$.

- (d) Mark the element T for *red*-refinement provided

$$(4.5) \quad \eta_{Z,T} \geq \Theta \max_{T' \in \mathcal{T}_k} \eta_{Z,T'}.$$

- (e) Mark further elements (*red-blue-green*-refinement) to avoid hanging nodes and generate a new triangulation $\mathcal{T}_{k+1}$. Update k and go to (b).

Remarks 4.1. (i) Note that the discrete contact zone is $\{x \in \Omega : u_{\varepsilon,h}(x) \leq \chi_h(x)\}$ and $u_{\varepsilon,h}(x) < \chi_h(x)$ can occur for some $x \in \Omega$.

(ii) For simplicity, we only computed an approximation $\mathcal{T}_{s,\varepsilon}$ of $\mathcal{T}_s$.

(iii) Since we only consider lowest order methods with optimal convergence rate $\mathcal{O}(h)$ the choice $\varepsilon = 1/N$ is motivated by $1/N \propto h^2$ in two dimensions.

(iv) The choice of the factors in the definition of $\eta_{Z,T}$ is motivated by the efficiency estimate of Remark 3.4.

5. NUMERICAL EXPERIMENTS

The theoretical results of this paper are supported by numerical experiments. In this section, we report on four examples of problem (1.1) on uniform and adapted meshes.

5.1. Example with smooth rotational symmetric solution [LLT]. Let $f := -2$ on $\Omega := (-3/2, 3/2)^2$ and $u_D(x, y) := r^2/2 - \ln(r) - 1/2$ where $r := (x^2 + y^2)^{1/2}$ on the Dirichlet boundary $\Gamma_D := \partial\Omega$. For $\chi_h = \chi := 0$ the exact solution of problem (1.1) then reads

$$u(x, y) = \begin{cases} r^2/2 - \ln(r) - 1/2 & \text{if } r \geq 1, \\ 0 & \text{otherwise.} \end{cases}$$

Note that $u \in H^2(\Omega)$. In our numerical experiments the coarse triangulation $\mathcal{T}_0$ of Fig. 1 consists of 16 squares halved along a diagonal.

The left plot in Fig. 1 shows a sequence of triangulations generated by Algorithm ($A_{1/2}$). The algorithm refines the mesh in the complement of the contact zone $\{(x, y) \in \Omega : x^2 + y^2 \leq 1\}$ in which the solution vanishes. The approximate discrete contact zone $\{T \in \mathcal{T}_k : u_{\varepsilon,h}(x_T) \leq \chi_h(x_T)\}$, where x_T denotes the center of a triangle T , is plotted in white while its complement is shaded (we chose this color since in most of the examples the complement of the contact zone is refined and appears darker). The right plot of Fig. 1 displays the solution $u_{\varepsilon,h}$

on the adaptively generated mesh $\mathcal{T}_8$ with 865 degrees of freedom. In Fig. 2 we plotted the error and its estimator versus the degrees of freedom for uniform and adaptive mesh refinement. A logarithmic scaling used for both axes allows a slope $-\alpha$ to be interpreted as an experimental convergence rate 2α (owing to $h \propto N^{-2}$ in two dimensions). We obtain experimental convergence rates 1 for both refinement strategies. The error on the adaptively refined meshes is however smaller than the error on uniform meshes at comparable numbers of degrees of freedoms. The plot shows that η_N serves as a very accurate approximate of the error e_N : The entries (N, e_N) and (N, η_N) almost coincide.

FIGURE 1. Adaptively refined meshes $\mathcal{T}_0$ (left upper) to $\mathcal{T}_8$ (right lower) (left) with approximate discrete contact zone shown in white and solution $u_{\epsilon,h}$ on $\mathcal{T}_8$ with 865 free nodes (right) in Example 5.1.

5.2. Example with corner singularity. Using polar coordinates (r, φ) on the L-shaped domain $\Omega := (-2, 2)^2 \setminus [0, 2] \times [-2, 0]$, $u_D := 0$ on $\Gamma_D := \partial\Omega$, $\chi_h = \chi := 0$, let $f(r, \varphi) := -r^{2/3} \sin(2\varphi/3)(\gamma_1'(r)/r + \gamma_1''(r)) - \frac{4}{3}r^{-1/3}\gamma_1'(r) \sin(2\varphi/3) - \gamma_2(r)$ where,

$$\gamma_1(r) := \begin{cases} 1 & \text{if } \bar{r} < 0, \\ -6\bar{r}^5 + 15\bar{r}^4 - 10\bar{r}^3 + 1 & \text{if } 0 \leq \bar{r} < 1, \\ 0 & \text{if } 1 \leq \bar{r}, \end{cases} \quad \text{for } \bar{r} := 2(r - 1/4),$$

and $\gamma_2(r) := 0$ if $r \leq 5/4$ and $\gamma_2(r) := 1$ otherwise. The exact solution of (1.1) is then given by $u(r, \varphi) := r^{2/3} \gamma_1(r) \sin(2\varphi/3)$ and has a typical corner singularity at the origin. The coarsest triangulation $\mathcal{T}_0$ of Fig. 3 consists of 48 halved squares.

The sequence of triangulations generated by Algorithm $(A_{1/2})$ in Example 5.2 and displayed in the left plot of Fig. 3 shows a refinement towards the origin where the solution has a singularity in the gradient and a refinement in the region $\{(x, y) \in \Omega : 1/4 \leq (x^2 + y^2)^{1/2} \leq 3/4\}$ where the solution has big gradients. This behavior can also be seen in the right plot of Fig. 3 where we plotted the numerical solution $u_{\epsilon,h}$ on triangulation $\mathcal{T}_8$ with 726 degrees of freedom. Fig. 4 shows that the adaptive Algorithm $(A_{1/2})$ improves the experimental

FIGURE 2. Error and error estimator for uniform and adaptive mesh-refinement in Example 5.1.

convergence rate about 0.9 for uniform mesh-refinement to the optimal value 1. Note that for uniform mesh-refinement we expect an asymptotic convergence rate $2/3$ due to the corner singularity. The error in the region where u has a large gradient seems to dominate in this preasymptotic range with $N \leq 10^5$. Again, the entries for η_N and e_N almost coincide and this behavior improves for increasing numbers of degrees of freedom.

5.3. First problem from elastoplastic torsion [R]. Let $f := 1$ on $\Omega := (-1, 1)^2$, $u_D := 0$ on $\Gamma_D := \partial\Omega$, $\Gamma_N := \emptyset$, and $\chi(x, y) := \text{dist}((x, y), \partial\Omega)$. In this example the exact solution of (1.1) is not known and cannot be expected to be smooth since $\chi \notin H^2(\Omega)$. The coarsest triangulation $\mathcal{T}_0$ of Fig. 5 consists of 64 elements with $\chi_h = \chi$ on $\mathcal{T}_0$.

This example is different from Examples 5.1 and 5.2 in the sense that the solution and the obstacle are non-smooth along the lines $\{(x, y) \in \Omega : x = y \text{ or } x = 1 - y\}$ of the contact zone. Algorithm $(A_{1/2})$ refines the mesh towards these lines as can be seen in the left and right plot of Fig. 5. Moreover, the approximate discrete contact zone reduces to these lines. Using Algorithm $(A_{1/2})$ the experimental convergence rate 0.5 for Algorithm (A_0) improves to the optimal value 1 for the error estimator at least in Fig 6.

5.4. Second problem from elastoplastic torsion [LLT]. Let $f := -3$ on $\Omega := (-1, 1)^2$, $u_D := 0$ on $\Gamma_D := \partial\Omega$, $\Gamma_N := \emptyset$, and $\chi_h(x, y) = \chi(x, y) := -\text{dist}((x, y), \partial\Omega)$. As in the previous example the solution is not known. The coarsest triangulation $\mathcal{T}_0$ is the same as in Example 5.3.

This example underlines the efficiency of our estimator and its robustness with respect to non-smoothness of the obstacle. The obstacle is non-smooth along the lines $\{(x, y) \in \Omega :$

FIGURE 3. Adaptively refined meshes $\mathcal{T}_0$ (left upper) to $\mathcal{T}_8$ (right lower) (left) with approximate discrete contact zone shown in white and solution $u_{\varepsilon,h}$ on $\mathcal{T}_8$ with 726 free nodes (right) in Example 5.2.

FIGURE 4. Error and error estimator for uniform and adaptive mesh-refinement in Example 5.2.

$x = y$ or $x = 1 - y$ but these lines do not belong to the contact zone and hence there should be no refinement towards them. The sequence of meshes in the left plot of Fig. 7 shows that the complement of the contact zone is indeed refined but the lines of non-smoothness of the

FIGURE 5. Adaptively refined meshes $\mathcal{T}_0$ (left upper) to $\mathcal{T}_8$ (right lower) (left) with approximate discrete contact zone shown in white and solution $u_{\varepsilon,h}$ on $\mathcal{T}_8$ with 2541 free nodes (right) in Example 5.3.

FIGURE 6. Error estimator for uniform and adaptive mesh-refinement in Example 5.3.

obstacle do not seem to play a special role in the refinement. The experimental convergence rate of the error estimator for uniform mesh-refinement equals one and this can be seen as

an indication that $u \in H^2(\Omega)$. The adaptive Algorithm $(A_{1/2})$ leads to smaller errors than Algorithm (A_0) , cf. Fig. 8, and shows also the optimal experimental convergence rate 1.

To illustrate that the term $\min_{q_h \in \mathcal{S}^1(\mathcal{T}_s)^d} \|\nabla(u_h - \chi_h) - q_h\|_{2,\Omega_s}$ should not be coarsened to $\min_{q_h \in \mathcal{S}^1(\mathcal{T})^d} \|\nabla(u_h - \chi_h) - q_h\|$ of (1.7) we compared η_N with the error estimator $\eta_{N,c}$,

$$2\eta_{N,c} := \|\nabla u_{\varepsilon,h} - \mathcal{A}\nabla u_{\varepsilon,h}\| + \|\nabla(u_{\varepsilon,h} - \chi_h) - \mathcal{B}\nabla(u_{\varepsilon,h} - \chi_h)\|.$$

As can be seen in Fig. 8 this error estimator leads to worse experimental convergence rates for uniform and adaptive mesh-refinement.

FIGURE 7. Adaptively refined meshes $\mathcal{T}_0$ (left upper) to $\mathcal{T}_8$ (right lower) (left) with approximate discrete contact zone plotted in white and solution $u_{\varepsilon,h}$ on $\mathcal{T}_8$ with 1429 free nodes (right) in Example 5.4.

5.5. Remarks. (i) The numerical results for Examples 5.1-5.4 show that the adaptive Algorithm $(A_{1/2})$ yields significant error reduction.

(ii) The error estimate performed extremely accurate although we only computed an approximation $u_{\varepsilon,h}$ to u_h .

(iii) As an initial function for the Newton scheme we used χ_h on $\mathcal{T}_0$ for the first mesh and successively the prolongation to $\mathcal{T}_{k+1}$ of the solution $u_{\varepsilon,h}$ on $\mathcal{T}_k$ for subsequent refinement levels (nested iteration). We stopped the iteration process when the Euclidean norm of the coefficient vector of the residual r_h of (4.4) satisfied $|r_h| \leq 10^{-12}$. In the above examples, the scheme converged after at most ten iteration steps.

(iv) The meshes generated by Algorithm (A_Θ) show local symmetries. A similar error estimator as η_N designed for second order partial differential equations performed well also on randomly perturbed meshes without any symmetry [CB, BC].

(v) The error estimator is reliable and efficient in Examples 5.1, 5.2, and 5.4. It is reliable (but possibly not efficient) in Example 5.3 owing to non-smoothness of the obstacle.

FIGURE 8. Error estimator η_N and coarsened estimator $\eta_{N,c}$ for uniform and adaptive mesh-refinement in Example 5.4.

Acknowledgments. The first author (S.B.) thankfully acknowledges partial support by the German Research Foundation (DFG) within the Graduiertenkolleg 'Effiziente Algorithmen und Mehrskalenmethoden'.

REFERENCES

- [AO] M. AINSWORTH, J.T. ODEN: *A Posteriori Error Estimation in Finite Element Analysis*. John Wiley & Sons 2000.
- [ACF] J. ALBERTY, C. CARSTENSEN, S.A. FUNKEN: Remarks around 50 lines of Matlab: short finite element implementation. *Num. Alg.* 20 (1999), 117–137.
- [BSt] I. BABUŠKA, T. STROUBOULIS: *The Finite Element Method and its Reliability*. Oxford University Press, publication date: 20 January 2001.
- [BC] S. BARTELS, C. CARSTENSEN: Each Averaging Technique Yields Reliable A Posteriori Error Control in FEM on Unstructured Grids. Part II: Higher Order FEM. *Berichtsreihe des Mathematischen Seminars Kiel*, Technical report **00-5**, Christian-Albrechts-Universität zu Kiel, Kiel (2000). (<http://www.numerik.uni-kiel.de/reports/2000/00-5.html>)
- [BCD] S. BARTELS, C. CARSTENSEN, G. DOLZMANN: Inhomogeneous Dirichlet conditions in a priori and a posteriori finite element error analysis. In preparation.
- [BSu] H. BLUM, F.-T. SUTTMEIER: Weighted error estimates for finite element solutions of variational inequalities. Preprint.
- [BSc] S.C. BRENNER, L.R. SCOTT: *The Mathematical Theory of Finite Element Methods*. Texts Appl. Math. 15, Springer, New-York (1994).
- [Ca] C. CARSTENSEN: Quasi interpolation and a posteriori error analysis in finite element method. *M²AN* 33 (1999) 1187-1202.
- [CA] C. CARSTENSEN, J. ALBERTY: Averaging techniques for reliable a posteriori FE-error control in elastoplasticity with hardening? *Berichtsreihe des Mathematischen Seminars Kiel*, Technical

- report **00-23**, Christian-Albrechts-Universität zu Kiel, Kiel (2000), (<http://www.numerik.uni-kiel.de/reports/2000/00-23.html>)
- [CB] C. CARSTENSEN, S. BARTELS: Each Averaging Technique Yields Reliable A Posteriori Error Control in FEM on Unstructured Grids Part I: Low Order Conforming, Nonconforming, and Mixed FEM. *Math. Comp.* (to appear). *Berichtsreihe des Mathematischen Seminars Kiel*, Technical report **99-11**, Christian-Albrechts-Universität zu Kiel, Kiel (1999). (<http://www.numerik.uni-kiel.de/reports/1999/99-11.html>)
- [CF1] C. CARSTENSEN, S.A. FUNKEN: Constants in Clément-interpolation error and residual based a posteriori error estimates in Finite Element Methods. *East-West Journal of Numerical Analysis* 8 (3) (2000) 153-256.
- [CF2] C. CARSTENSEN, S.A. FUNKEN: Averaging technique for FE a posteriori error control in elasticity. Part I: conforming FEM. *Berichtsreihe des Mathematischen Seminars Kiel*, Technical report **99-14**, Christian-Albrechts-Universität zu Kiel, Kiel (1999). *Comp. Meth. Appl. Mech. Engrg.* (accepted). (<http://www.numerik.uni-kiel.de/reports/1999/00-14.html>)
- [CF3] C. CARSTENSEN, S.A. FUNKEN: Averaging technique for FE a posteriori error control in elasticity. Part II: λ -independent estimates. *Berichtsreihe des Mathematischen Seminars Kiel*, Technical report **00-6**, Christian-Albrechts-Universität zu Kiel, Kiel (2000). *Comp. Meth. Appl. Mech. Engrg.* (accepted). (<http://www.numerik.uni-kiel.de/reports/2000/00-6.html>)
- [CF4] C. CARSTENSEN, S.A. FUNKEN: Averaging technique for FE a posteriori error control in elasticity. Part III: Locking-free nonconforming FEM. *Comp. Meth. Appl. Mech. Engrg.* (accepted). *Berichtsreihe des Mathematischen Seminars Kiel*, Technical report **00-11**, Christian-Albrechts-Universität zu Kiel, Kiel (2000). (<http://www.numerik.uni-kiel.de/reports/2000/00-11.html>)
- [CF5] C. CARSTENSEN, S.A. FUNKEN: A posteriori error control in low-order finite element discretisations of incompressible stationary flow problems. *Math. Comp.* (accepted). *Berichtsreihe des Mathematischen Seminars Kiel*, Technical report **99-5**, Christian-Albrechts-Universität zu Kiel, Kiel (1999). (<http://www.numerik.uni-kiel.de/reports/1999/99-5.html>)
- [CV] C. CARSTENSEN, R. VERFÜRTH: Edge residuals dominate a posteriori error estimates for low order finite element methods. *SIAM J. Numer. Anal.* **36** No. 5 (1999) 1571-1587.
- [CN] Z. CHEN, R.H. NOCHETTO: Residual type a posteriori error estimates for elliptic obstacle problems. *Numer. Math.* **84**, 4, 527-548 (2000).
- [Ci] P.G. CIARLET: *The finite element method for elliptic problems*. North-Holland, Amsterdam, 1978.
- [Cl] P. CLÉMENT: Approximation by finite element functions using local regularization. *RAIRO Sér. Rouge Anal. Numér.* **R-2** (1975) 77-84.
- [EEHJ] K. ERIKSSON, D. ESTEP, P. HANSBO, C. JOHNSON: Introduction to adaptive methods for differential equations. *Acta Numerica* (1995) 105-158.
- [EG] L.C. EVANS, R.F. GARIEPY: *Measure Theory and fine properties of functions*. Studies in Advanced Mathematics. CRC Press (1992).
- [F] R.S. FALK: Error estimates for the approximation of a class of variational inequalities. *Math. Comp.* **28**, 963-971 (1974).
- [GLT] R. GLOWINSKI, J.L. LIONS, R. TREMOLIERES: *Numerical analysis of variational inequalities*. Studies in Mathematics and its Applications, Vol. 8. Amsterdam, New York, Oxford: North-Holland Publishing Company (1981).
- [K] R. KORNUBER: *Adaptive monotone multigrid methods for nonlinear variational problems*. B. G. Teubner (1997).
- [LLT] R. LI, W. LIU, AND TAO TANG: Moving Mesh Finite Element Approximations for Variational Inequality I: Static Obstacle Problems. Preprint.
- [N] F. NATTERER: Optimale L^2 -Konvergenz finiter Elemente bei Variationsungleichungen. *Bonner Math. Schr.*, **89**, 1-12, (1976).
- [NW] R.H. NOCHETTO, L.B. WAHLBIN: Positivity preserving finite element interpolation. In preparation.
- [R] J.-F. RODRIGUES: *Obstacle problems in mathematical physics*. North Holland Studies in Mathematics, Vol 134, North-Holland (1987).
- [V1] A. VEESER: On a posteriori error estimation for constant obstacle problems. Freiburg, Preprint 29 (1999).

- [V2] A. VEESER: Efficient and reliable a posteriori error estimators for elliptic obstacle problems. Preprint (2000).
- [V] R. VERFÜRTH: *A review of a posteriori error estimation and adaptive mesh-refinement techniques*. Wiley-Teubner (1996).

MATHEMATISCHES SEMINAR, CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL, LUDEWIG-MEYN-STR. 4, D-24098 KIEL, GERMANY.

E-mail address: `sba@numerik.uni-kiel.de`

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTR. 22-26, D-04103 LEIPZIG, GERMANY (ON LEAVE FROM MATHEMATISCHES SEMINAR, CHRISTIAN-ALBRECHTS-UNIVERSITÄT ZU KIEL, LUDEWIG-MEYN-STR. 4, D-24098 KIEL, GERMANY).

E-mail address: `cc@numerik.uni-kiel.de`