

ADDENDUM 1 : CHERN-WEIL CHARACTERISTIC CLASSES

HERE WE WILL SIMPLY RECORD THE PERTINENT DEFINITIONS AND RESULTS AND, AS AN ILLUSTRATION, SKETCH ONE SIMPLE CALCULATION.

A SMOOTH PRINCIPAL BUNDLE CONSISTS OF A SMOOTH (C^∞) MANIFOLD P (BUNDLE SPACE), A SMOOTH MANIFOLD M (BASE SPACE), A SMOOTH MAP π (PROJECTION) OF P ONTO M , A LIE GROUP G (STRUCTURE GROUP) AND A SMOOTH RIGHT ACTION

$$\sigma : P \times G \rightarrow P$$

$$\sigma(p, g) = p \cdot g = \sigma_p(g) = \sigma_g(p)$$

OF G ON P SUCH THAT THE FOLLOWING ARE SATISFIED :

1. σ PRESERVES FIBERS OF π : $\pi(p \cdot g) = \pi(p)$
 $\forall p \in P \forall g \in G$

2. (LOCAL TRIVIALITY) FOR EACH $x \in M$ \exists OPEN NEIGHBORHOOD U OF x IN M (TRIVIALIZING NBD) AND A DIFFEOMORPHISM $\Psi : \pi^{-1}(U) \rightarrow U \times G$ (TRIVIALIZATION) OF THE FORM $\Psi(p) = (\pi(p), \psi(p))$, WHERE $\psi : \pi^{-1}(U) \rightarrow G$ SATISFIES

$$\psi(p \cdot g) = \psi(p) \cdot g$$

$\forall p \in \pi^{-1}(U) \forall g \in G$ (ψ IS EQUIVARIANT WITH RESPECT TO σ AND THE NATURAL ACTION OF G ON $U \times G$).

$$G \hookrightarrow P \xrightarrow{\pi} M$$

(LOCAL) SECTION/GAUGE : $\Delta : U \rightarrow \pi^{-1}(U)$, $\pi \circ \Delta = id_U$

SECTIONS \longleftrightarrow TRIVIALIZATIONS

$$\Delta(x) \cdot g \longleftrightarrow (x, g)$$

TRANSITION FUNCTIONS : $\Delta_i : U_i \rightarrow \pi^{-1}(U_i)$

$$\Delta_j : U_j \rightarrow \pi^{-1}(U_j)$$


$$\Delta_j(x) = \Delta_i(x) \cdot g_{ij}(x)$$

ASSOCIATED FIBER BUNDLES : F SMOOTH MANIFOLD WITH LEFT ACTION OF G

$$(g, \xi) \rightarrow g \cdot \xi$$

RIGHT ACTION OF G ON $P \times F$: $(p, \xi) \cdot g = (p \cdot g, g^{-1} \cdot \xi)$

ORBIT SPACE : $P \times_G F$



IF $F = V$ IS A VECTOR SPACE AND THE ACTION OF G COMES FROM A REPRESENTATION $\rho : G \rightarrow GL(V)$, THEN THE ASSOCIATED BUNDLE $P \times_{\rho} V$ IS A VECTOR BUNDLE, E.G., ADJOINT BUNDLE

$$ad P = P \times_{ad} \mathfrak{g}$$

THREE (EQUIVALENT) DEFINITIONS OF A CONNECTION ON $G \hookrightarrow P \xrightarrow{\pi} M$:

ASSUME G IS A MATRIX LIE GROUP WITH LIE ALGEBRA \mathfrak{g} AND $\dim M = n$

1. n -DIMENSIONAL DISTRIBUTION $p \in P \rightarrow \text{HOR}_p(P) \subseteq T_p(P)$ S.T.

$$(a) \quad T_p(P) = \text{HOR}_p(P) \oplus \text{VERT}_p(P), \text{ WHERE}$$

$$\text{VERT}_p(P) = T_p(\pi^{-1}(\pi(p)))$$

$$(b) \quad \text{HOR}_{p \cdot g}(P) = (\sigma_g)_{*p}(\text{HOR}_p(P))$$

2. \mathfrak{g} -VALUED 1-FORM ω ON P S.T.

$$(a) \quad \omega_{p \cdot g}((\sigma_g)_{*p}(v_p)) = g^{-1} \omega_p(v_p) g$$

$$(b) \quad \omega_p(\xi^*(p)) = \xi \quad \forall \xi \in \mathfrak{g}, \text{ WHERE}$$

$$\xi^*(p) = \left. \frac{d}{dt} (p \cdot \exp(t\xi)) \right|_{t=0}$$

3. A TRIVIALIZING COVER $\{(U_j, \psi_j)\}$ OF M FOR $G \hookrightarrow P \xrightarrow{\pi} M$

AND, FOR EACH j , A \mathfrak{g} -VALUED 1-FORM a_j ON U_j S.T.

WHENEVER $U_j \cap U_i \neq \emptyset$

$$a_j = g_{ij}^{-1} a_i g_{ij} + g_{ij}^{-1} dg_{ij}$$

RELATIONS : $\text{HOR}_p(P) = \text{KER } \omega_p$ AND $a_j = \psi_j^* \omega$

CURVATURE OF A CONNECTION 1-FORM ω IS THE \mathfrak{g} -VALUED 2-FORM

Ω ON P DEFINED BY

$$\Omega_p(\nu_p, \omega_p) = d\omega_p(\nu_p^{\text{HOR}}, \omega_p^{\text{HOR}})$$

WHERE ν_p^{HOR} AND ω_p^{HOR} ARE THE PROJECTIONS OF ν_p AND ω_p ONTO $\text{HOR}_p(P)$ IN $T_p(P) = \text{HOR}_p(P) \oplus \text{VERT}_p(P)$.

CARTAN STRUCTURE EQUATION: $\Omega = d\omega + \omega \wedge \omega$

LOCAL FIELD STRENGTHS: $\mathcal{F}_j = \nu_j^* \Omega = d\mathcal{A}_j + \mathcal{A}_j \wedge \mathcal{A}_j$

$$\mathcal{F}_j = g_{ij}^{-1} \mathcal{F}_i g_{ij}$$

\Rightarrow THE \mathcal{F}_j PIECE TOGETHER INTO A 2-FORM

$$F_\omega \in \Omega^2(M, \text{ad}P)$$

ON M WITH VALUES IN THE ADJOINT BUNDLE $\text{ad}P$. THIS IS ALSO CALLED THE CURVATURE OF ω .

CHERN-WEIL PROCEDURE:

ASSUME NOW THAT G IS A COMPACT MATRIX LIE GROUP WITH LIE ALGEBRA

\mathfrak{g} AND CONSIDER THE ALGEBRA

$$\mathbb{C}[\mathfrak{g}]^G$$

OF $\text{ad}G$ -INVARIANT, COMPLEX-VALUED POLYNOMIAL FUNCTIONS ON \mathfrak{g} .

MORE DETAIL: LET $\{\xi_1, \dots, \xi_n\}$ BE A BASIS FOR \mathfrak{g}
 AND $\{x^1, \dots, x^n\}$ THE DUAL BASIS FOR \mathfrak{g}^* ($x^a(\xi_b) = \delta_b^a$).
 THE SYMMETRIC ALGEBRA $S(\mathfrak{g}^*) = S[x^1, \dots, x^n]$ IS
 THE ALGEBRA OF POLYNOMIALS WITH REAL COEFFICIENTS
 IN x^1, \dots, x^n (REAL-VALUED POLYNOMIAL FUNCTIONS
 ON \mathfrak{g}). THERE IS A LEFT ACTION OF G ON $S(\mathfrak{g}^*)$:
 $\rho \in S(\mathfrak{g}^*)$ AND $g \in G$ GIVE $g \cdot \rho \in S(\mathfrak{g}^*)$ DEFINED BY

$$(g \cdot \rho)(\xi) = \rho(\text{ad}_{g^{-1}} \xi) = \rho(g^{-1} \xi g).$$

$$S(\mathfrak{g}^*)^G = \{\rho \in S(\mathfrak{g}^*) : g \cdot \rho = \rho \ \forall g \in G\}, \text{ I.E.,}$$

$$\rho(g^{-1} \xi g) = \rho(\xi).$$

$$\mathbb{C}[\mathfrak{g}] = S(\mathfrak{g}^*) \otimes \mathbb{C} \text{ AND } \mathbb{C}[\mathfrak{g}]^G = S(\mathfrak{g}^*)^G \otimes \mathbb{C}.$$

EVERY $\rho \in \mathbb{C}^{\wedge}[\mathfrak{g}]^G$ GIVES RISE, VIA POLARIZATION, TO
 A G -INVARIANT \mathbb{R} -MULTILINEAR FUNCTION, ALSO DENOTED

$$\rho : \mathfrak{g} \times \dots \times \mathfrak{g} \rightarrow \mathbb{C}$$

CHERN-WEIL IS BASICALLY A MAP

$$\mathbb{C}[\mathfrak{g}]^G \rightarrow \Omega^*(P)_{\text{BASIC}}$$

FROM G -INVARIANT POLYNOMIALS ON \mathfrak{g} TO DIFFERENTIAL FORMS
 ON P THAT ARE BASIC, I.E., G -INVARIANT ($\sigma_g^* \omega = \omega \ \forall g \in G$)
 AND HORIZONTAL ($i_V \omega = 0 \ \forall$ VERTICAL VECTOR FIELD V ON P)

NOTE: BASIC FORMS ON THE PRINCIPAL BUNDLE SPACE P ARE PRECISELY THOSE $\varphi \in \Omega^*(P)$ WHICH DESCEND TO THE BASE MANIFOLD M , I.E., FOR WHICH $\exists \bar{\varphi} \in \Omega^*(M)$ WITH $\pi^* \bar{\varphi} = \varphi$.

GIVEN $\varrho \in [\mathfrak{g}]^G$ (THOUGHT OF AS A k -MULTILINEAR FUNCTION) CHOOSE A CONNECTION ω ON $G \hookrightarrow P \xrightarrow{\pi} M$ AND LET Ω BE ITS CURVATURE. WRITE $\omega = \omega^a \xi_a$ AND $\Omega = \Omega^a \xi_a$. DEFINE $\varrho(\Omega) \in \Omega^{2k}(P)$ BY

$$\begin{aligned} \varrho(\Omega) &= \varrho(\Omega^{a_1} \xi_{a_1}, \dots, \Omega^{a_k} \xi_{a_k}) \\ &= \varrho(\xi_{a_1}, \dots, \xi_{a_k}) \Omega^{a_1} \wedge \dots \wedge \Omega^{a_k} \end{aligned}$$

THEN $\varrho(\Omega)$ IS A CLOSED, BASIC $2k$ -FORM ON P SO IT DESCENDS TO A CLOSED $2k$ -FORM $\bar{\varrho}(\Omega)$ ON M :

$$\pi^*(\bar{\varrho}(\Omega)) = \varrho(\Omega)$$

ESSENTIALLY, $\bar{\varrho}(\Omega)$ IS JUST THE RESTRICTION OF $\varrho(\Omega)$ TO THE ω -HORIZONTAL SPACES IN TP .

THE COHOMOLOGY CLASS $[\bar{\varrho}(\Omega)] \in H^{2k}(M; \mathbb{R})$ DOES NOT DEPEND ON THE CHOICE OF ω OR THE BASIS $\{\xi_1, \dots, \xi_n\}$ FOR \mathfrak{g} AND IS CALLED A CHARACTERISTIC CLASS OF $G \hookrightarrow P \xrightarrow{\pi} M$.

AS AN EXAMPLE WE WILL CONSIDER THE 2ND CHERN CLASS OF A PRINCIPAL $SU(2)$ -BUNDLE

$$SU(2) \hookrightarrow P \xrightarrow{\pi} M$$

OVER A COMPACT, CONNECTED, ORIENTED 4-MANIFOLD M

NOTE : ANOTHER EXAMPLE (THE EULER CLASS) WILL BE DISCUSSED IN THE NEXT SECTION ON "EQUIVARIANT COHOMOLOGY AND THE WITTEN LAGRANGIAN".

NEED TO CHOOSE

1. A BASIS $\{\xi_1, \xi_2, \xi_3\}$ FOR $\mathfrak{su}(2)$ (2×2 COMPLEX MATRICES THAT ARE SKEW-HERMITIAN AND TRACEFREE), E.G.,

$$\xi_1 = -\frac{i}{2} \sigma_1 = -\frac{i}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 0 & -i \\ -i & 0 \end{pmatrix}$$

$$\xi_2 = -\frac{i}{2} \sigma_2 = -\frac{i}{2} \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

$$\xi_3 = -\frac{i}{2} \sigma_3 = -\frac{i}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = \frac{1}{2} \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$$

2. A CONNECTION ω WITH CURVATURE Ω WHICH WE WRITE AS

$$\omega = \omega^a \xi_a = \frac{1}{2} \begin{pmatrix} -\omega^3 i & -\omega^2 - i\omega^1 \\ \omega^2 - i\omega^1 & \omega^3 i \end{pmatrix}$$

$$\Omega = \Omega^a \xi_a = \frac{1}{2} \begin{pmatrix} -\Omega^3 i & -\Omega^2 - i\Omega^1 \\ \Omega^2 - i\Omega^1 & \Omega^3 i \end{pmatrix}$$

3. AN INVARIANT POLYNOMIAL ρ ON $SU(2)$. FOR THIS WE TAKE

$$\rho : SU(2) \rightarrow \mathbb{C}$$

$$\rho(A) = \frac{1}{8\pi^2} \text{Tr}(A^2)$$

THE CORRESPONDING BILINEAR FORM ON $SU(2)$ IS CLEARLY

$$\rho : SU(2) \times SU(2) \rightarrow \mathbb{C}$$

$$\rho(A, B) = \frac{1}{8\pi^2} \text{Tr}(AB)$$

THUS,

$$\begin{aligned} \rho(\Omega) &= \rho(\Omega^{a_1} \xi_{a_1}, \Omega^{a_2} \xi_{a_2}) \\ &= \rho(\xi_{a_1}, \xi_{a_2}) \Omega^{a_1} \wedge \Omega^{a_2} \\ &= \frac{1}{8\pi^2} \text{Tr}(\xi_{a_1}, \xi_{a_2}) \Omega^{a_1} \wedge \Omega^{a_2} \end{aligned}$$

FOR THE BASIS CHOSEN ABOVE

$$\text{Tr}(\xi_{a_1}, \xi_{a_2}) = \begin{cases} 0 & , a_1 \neq a_2 \\ -\frac{1}{2} & , a_1 = a_2 \end{cases}$$

SO

$$\begin{aligned} \rho(\Omega) &= \frac{1}{8\pi^2} \left(-\frac{1}{2}\right) (\Omega^1 \wedge \Omega^1 + \Omega^2 \wedge \Omega^2 + \Omega^3 \wedge \Omega^3) \\ &= \frac{1}{8\pi^2} \text{Tr}(\Omega \wedge \Omega) \end{aligned}$$

WHERE $\Omega \wedge \Omega$ IS THE MATRIX PRODUCT WITH ENTRIES MULTIPLIED BY THE ORDINARY WEDGE PRODUCT.

WE KNOW THAT

$$\frac{1}{8\pi^2} \text{Tr} (\Omega \wedge \Omega)$$

IS A CLOSED, BASIC 4-FORM ON P AND SO DESCENDS TO A CLOSED 4-FORM ON M (I.E., IS π^* OF SOME CLOSED 4-FORM ON M).

THIS CLOSED 4-FORM ON M CAN BE DESCRIBED LOCALLY IN TERMS OF LOCAL FIELD STRENGTHS

$$\mathcal{F}_i = \mathcal{A}_i^* \Omega$$

$$\text{SINCE } \pi^* \mathcal{F}_i = \pi^* (\mathcal{A}_i^* \Omega) = (\mathcal{A}_i \circ \pi)^* \Omega = \text{id}^* \Omega = \Omega,$$

$$\pi^* \left(\frac{1}{8\pi^2} \text{Tr} (\mathcal{F}_i \wedge \mathcal{F}_i) \right) = \frac{1}{8\pi^2} \text{Tr} (\Omega \wedge \Omega)$$

ON ITS DOMAIN. SINCE THE LOCAL FIELD STRENGTHS TRANSFORM UNDER THE ADJOINT REPRESENTATION ($\mathcal{F}_i = g_{ij}^{-1} \mathcal{F}_i g_{ij}$)

THE SAME IS TRUE OF $\mathcal{F}_i \wedge \mathcal{F}_i$ AND Tr IS AD-INVARIANT SO THE

$$\frac{1}{8\pi^2} \text{Tr} (\mathcal{F}_i \wedge \mathcal{F}_i)$$

PIECE TOGETHER INTO THE GLOBALLY DEFINED 4-FORM ON M TO WHICH $\frac{1}{8\pi^2} \text{Tr} (\Omega \wedge \Omega)$ DESCENDS. WE WRITE THIS AS

$$\frac{1}{8\pi^2} \text{Tr} (F_\omega \wedge F_\omega)$$

AND CALL ITS COHOMOLOGY CLASS

$$c_2(P) \in H^4(M; \mathbb{R})$$

THE 2ND CHERN CLASS OF $SU(2) \hookrightarrow P \xrightarrow{\pi} M$, ITS INTEGRAL OVER M

$$c_2(P)[M] = \int_M c_2(P) = \frac{1}{8\pi^2} \int_M \text{Tr}(F_\omega \wedge F_\omega),$$

WHICH IS ACTUALLY AN INTEGER, IS CALLED THE 2ND CHERN NUMBER OF $SU(2) \hookrightarrow P \xrightarrow{\pi} M$

NOTE : FOR THE QUATERNIONIC HOPF BUNDLE

$$SU(2) \hookrightarrow S^7 \xrightarrow{\pi} S^4$$

ONE CAN USE ANY OF THE BPST INSTANTON CONNECTIONS TO COMPUTE REPRESENTATIVES OF THE CHERN CLASS. MOREOVER, THE BUNDLE IS TRIVIAL OVER $S^4 - \{\text{POINT}\}$ WHICH STEREOGRAPHICALLY PROJECTS TO \mathbb{R}^4 SO THE CHERN NUMBER CAN BE COMPUTED AS AN INTEGRAL OVER \mathbb{R}^4 . THE CALCULATION WAS SKETCHED IN THE LECTURE. THE RESULT IS 1.

FOR THE COMPACT, SIMPLY CONNECTED, ORIENTED, SMOOTH 4-MANIFOLDS M OF INTEREST TO US HERE, PRINCIPAL $SU(2)$ -BUNDLES OVER M ARE CLASSIFIED UP TO EQUIVALENCE BY THEIR 2ND CHERN CLASS / NUMBER.