

THE THRESHOLDING SCHEME FOR MEAN CURVATURE FLOW AND DE GIORGI'S IDEAS FOR MINIMIZING MOVEMENTS

TIM LAUX AND FELIX OTTO

ABSTRACT. We consider the thresholding scheme and explore its connection to De Giorgi's ideas on gradient flows in metric spaces; here applied to mean curvature flow as the steepest descent of the interfacial area. The basis of our analysis is the observation by Esedoğlu and the second author that thresholding can be interpreted as a minimizing movements scheme for an energy that approximates the interfacial area. De Giorgi's framework provides an optimal energy dissipation relation for the scheme in which we pass to the limit to derive a dissipation-based weak formulation of mean curvature flow. Although applicable in the general setting of arbitrary networks, here we restrict ourselves to the case of a single interface, which allows for a compact, self-contained presentation.

1. INTRODUCTION AND CONTEXT

The purpose of these notes is to draw a connection between De Giorgi's tools for minimizing movements, that is, gradient flows in metric spaces on the one hand, and the very popular thresholding scheme for flow of a hyper-surface by its mean curvature on the other hand. While we have developed this connection in the case of multiple phases with surface energies and mobilities depending on the pair of phases, as is relevant for grain growth in polycrystals, and when the notion of viscosity solution is not available, we present our results here in the simplest setting of two phases. Our presentation is essentially self-contained.

What makes the evolution of the boundary $\partial\Omega$ of a set Ω by its mean curvature H valuable for modeling in materials science is that it is driven by the reduction of the (total) interfacial area of $\partial\Omega$, which relies on the mean curvature H , the sum of the principal curvatures, being the first variation of the interfacial area. There is a more intimate connection between mean curvature flow (MCF) and the functional E of interfacial area of a configuration: MCF can formally be understood as a gradient flow of E . We stress that a dynamical system that can be written as a gradient flow, that is, a steepest descent in an energy landscape, does not just rely on the height function E , but also on a notion of distance on configuration space, which is typically described by a metric tensor g in the sense of Riemannian geometry. In case of

MCF, the tangent space in some configuration Ω should be thought of as consisting of all normal velocities V , i.e., functions on $\partial\Omega$, while the configuration-dependent metric tensor g_Ω is given by the L^2 -inner product on $\partial\Omega$.

Still formally, any gradient flow allows for a natural discretization in time. Every step of the discretization comes in form of a variational problem, just involving the functional E and the induced distance d , cf. (9), but not the metric tensor g and the differential of E – it thus relies rather on the “metric”, but not the differential structure. Following De Giorgi, we call such a scheme a minimizing movements scheme. We recall that, as in elementary differential geometry, the induced distance d on a Riemannian manifold (\mathcal{M}, g) is defined via $d^2(\chi_0, \chi_1) := \inf\{\int_0^1 g_{\chi_s}(\frac{d\chi}{ds}, \frac{d\chi}{ds})ds\}$, where the infimum is taken over all curves $[0, 1] \ni s \mapsto \chi_s$ connecting χ_0 to χ_1 (we use the letter χ because we think of a characteristic function describing the configuration). We note that in the Euclidean case, the Euler-Lagrange equation of (9) turns into the implicit Euler scheme for $\frac{dx}{dt} = \text{grad } E|_\chi$.

However, this infinite-dimensional Riemannian structure making MCF a gradient flow leads to a degenerate induced metric (i.e. $d \equiv 0$): It can be seen that the infimum of $\int_0^1 \int_{\partial\Omega_s} V_s^2 ds$ over all curves of configurations $[0, 1] \ni s \mapsto \Omega_s$ with normal velocity $[0, 1] \ni s \mapsto V_s$, connecting some given configurations Ω_0 and Ω_1 , vanishes [9].

Nonetheless, a minimizing movements scheme (before the latter) for MCF has been formulated by Almgren et. al. [1], with $E(\Omega)$ being the surface area of $\partial\Omega$ and with $d^2(\Omega_1, \Omega_0) = 4 \int_{\Omega_1 \Delta \Omega_0} \text{dist}(\cdot, \partial\Omega_0)$. Luckhaus et. al. [7] have established a (long-time) convergence result for this scheme. This convergence result is conditional in the sense that a condition like in (5) has to be imposed.

Thresholding, cf. (1), is a very well performing and widely used numerical scheme for MCF, introduced by Osher et. al. [8]. Also the convolution step, which after spatial discretization can be carried out by the Fast Fourier Transform, is of low complexity. Right from the beginning, thresholding has attracted the attention of analysts; since it obviously conserves the comparison principle for MCF, it has been shown to converge to MCF in the sense of viscosity solution in the two-phase case [4].

Esedoglu and the second author [3] realized that thresholding also respects the gradient-flow structure of MCF, in the sense that it can be interpreted as a minimizing movements scheme, cf. Lemma 2. This was used in the multi-phase case to extend thresholding to surface tensions and mobilities [10] that depend on the pair of grains, while keeping its low complexity. It was also used by the present authors to provide several types of convergence results; presently, all of them are conditional in the sense of assumption (5), in the tradition of [7].

The first result [5] provided the same limiting notion of solution for MCF as in [7]. However, this weak notion of solution does not imply the dissipation inequality natural to a gradient flow. It is Brakke's weak notion of solution for MCF that is based on a localization of the dissipation inequality; in [6], we establish a (still conditional) convergence result towards this inequality-based notion of solution.

For any gradient flow in a Riemannian context (\mathcal{M}, g, E) , there is yet another notion of weak solution based on a single inequality, namely $E(\chi(T)) + \int_0^T \frac{1}{2} g_\chi(\frac{d\chi}{dt}, \frac{d\chi}{dt}) + \frac{1}{2} |\text{grad} E|_\chi|^2 dt \leq E(\chi(0))$. This elementary observation is credited to De Giorgi; its appeal lies in the fact that it is potentially more stable in limiting procedures because only lower semi-continuity is needed (as provided by Propositions 1 and 2). The main result of this paper, Theorem 1, precisely establishes this inequality in the case of MCF, cf. (8).

One advantage of a minimizing movements scheme, cf. (9), lies in the fact that it automatically comes with the a priori estimate $E(\chi^N) + \sum_{n=1}^N \frac{1}{2h} d^2(\chi^n, \chi^{n-1}) \leq E(\chi^0)$, which is obtained by using χ^{n-1} as a competitor in (9). In the limit $h \downarrow 0$, this inequality formally turns into $E(\chi(T)) + \int_0^T \frac{1}{2} g_\chi(\frac{d\chi}{dt}, \frac{d\chi}{dt}) dt \leq E(\chi(0))$, which misses the formally correct identity by a factor of 2. On the level of the metric structure, De Giorgi provides tools to capture the missing term $\int_0^T \frac{1}{2} |\text{grad} E|_\chi|^2 dt$, see Lemma 1. We take the proof from the monograph [2].

As a consequence of these notions and tools of De Giorgi, our (conditional) convergence proof for the thresholding scheme in fact is rather "soft", softer than [5] which relied on the notion of tilt excess and the fine structure of Caccioppoli sets, and certainly softer than [7] which relied on regularity theory for minimal surfaces. We believe that these tools have a wider potential for geometric evolutions or non-linear PDE of gradient-flow type. For the broader context and more references, we refer to [5].

2. MAIN RESULT AND STRUCTURE OF PROOF

Given an initial configuration, as described by its (Lebesgue-measurable) characteristic function $\chi^0: \mathbb{R}^d \rightarrow \{0, 1\}$, and a time step size $h > 0$, the thresholding scheme iteratively produces configurations at time steps $n = 1, 2, \dots$, encoded by their characteristic functions χ^n , via convolution and "thresholding":

$$(1) \quad \chi^n := \left\{ \begin{array}{ll} 1 & \text{where } G_h * \chi^{n-1} > \frac{1}{2} \\ 0 & \text{else} \end{array} \right\},$$

where G_h denotes the heat kernel at time $\frac{h}{2}$, that is,

$$(2) \quad G_h(z) := \frac{1}{\sqrt{2\pi h}^d} \exp\left(-\frac{|z|^2}{2h}\right)$$

(like in stochastic analysis, we take $\frac{h}{2}$ so that G_1 is the standard Gaussian). We interpolate piecewise constant in time:

$$(3) \quad \chi_h(t) = \chi^n \quad \text{for } t \in [nh, (n+1)h), \quad \chi_h(t) = \chi^0 \quad \text{for } t \leq 0.$$

For simplicity, we pass from the whole space \mathbb{R}^d to a torus as the spatial domain; by rescaling, we may w. l. o. g. take the unit torus $[0, 1]^d$. We also restrict to the finite time horizon $T < \infty$ and $h \leq 1$.

Our main result is the following convergence result, which is only a conditional one since assumption (5) on the energies E_h defined below in (15) presumably cannot be verified. It is the opposite direction, $c_0 \int_{(0,T) \times [0,1]^d} |\nabla \chi| dt \leq \liminf_{h \downarrow 0} \int_0^T E_h(\chi_h(t)) dt$, that follows from (4), see (19) in Lemma 3. Here and in the sequel, $|\nabla \chi| dt$ denotes the total variation of the distribution $\nabla \chi$ in $(0, T) \times [0, 1]^d$, provided the latter is a bounded measure. This notation is justified since in the present case, $|\nabla \chi| dt$ is (Lebesgue) equi-integrable in t and thus admits a density $|\nabla \chi|$. In the sequel, $\nu \in L^1(|\nabla \chi| dt)$ denotes the measure-theoretic normal (characterized through the polar factorization $\nabla \chi = \nu |\nabla \chi| dt$ and $|\nu| = 1$ $|\nabla \chi| dt$ -almost everywhere).

Theorem 1. *Given χ^0 as above and such that $\nabla \chi^0$ is a bounded measure, and a sequence $h \downarrow 0$; let χ_h be defined by (1) and (3). Suppose that there exists a $\chi: (0, T) \times [0, 1]^d \rightarrow [0, 1]$ such that*

$$(4) \quad \chi_h \rightharpoonup \chi \quad \text{in } L^1((0, T) \times [0, 1]^d).$$

Then we have $\chi \in \{0, 1\}$ (Lebesgue)-a. e. and $\nabla \chi$ is a bounded measure which is equi-integrable in t . If we assume in addition

$$(5) \quad \limsup_{h \downarrow 0} \int_0^T E_h(\chi_h(t)) dt \leq c_0 \int_{(0,T) \times [0,1]^d} |\nabla \chi| dt,$$

where $c_0 := \frac{1}{\sqrt{2\pi}}$, then there exists $H \in L^2(|\nabla \chi| dt)$ with

$$(6) \quad \int_{(0,T) \times [0,1]^d} (\nabla \cdot \xi - \nu \cdot \nabla \xi \nu) |\nabla \chi| dt = - \int_{(0,T) \times [0,1]^d} H \nu \cdot \xi |\nabla \chi| dt$$

for all $\xi \in C_0^\infty((0, T) \times [0, 1]^d)$, and $V \in L^2(|\nabla \chi| dt)$ with

$$(7) \quad \int_{[0,1]^d} \zeta(t=0) \chi^0 dx + \int_{(0,T) \times [0,1]^d} \partial_t \zeta \chi dx dt + \int_{(0,T) \times [0,1]^d} \zeta V |\nabla \chi| dt = 0$$

for all $\zeta \in C_0^\infty([0, T) \times [0, 1]^d)$, such that

$$(8) \quad \limsup_{\tau \downarrow 0} \frac{1}{\tau} \int_{(T-\tau, T) \times [0,1]^d} |\nabla \chi| dt + \int_{(0,T) \times [0,1]^d} (V^2 + (\frac{H}{2})^2) |\nabla \chi| dt \leq \int_{[0,1]^d} |\nabla \chi^0|.$$

We note that in case of $\{\chi = 1\}$ being smooth in time-space $[0, T] \times [0, 1]^d$, $|\nabla\chi|$ coincides with the surface measure, ν with the (inner) normal, and H and V coincide with the mean curvature (with the convention that convex sets have positively curved boundary) and normal velocity (with the convention that growing sets have positive velocity), respectively. In addition, (7) yields that $\chi(t = 0) = \chi^0$. Moreover, expanding the square $V^2 + (\frac{H}{2})^2 = (V + \frac{H}{2})^2 - VH$, and appealing to the classical formula $\frac{d}{dt} \int_{[0,1]^d} |\nabla\chi| = \int_{[0,1]^d} VH|\nabla\chi|$ (which relies on the fact that mean curvature describes the first variation of the surface area), we see that (8) turns into $\int_{(0,T) \times [0,1]^d} (V + \frac{H}{2})^2 |\nabla\chi| dt \leq 0$, and thus MCF in form of $V = -\frac{H}{2}$ (the factor $\frac{1}{2}$ stems from the normalization in (2)). Therefore, the inequality (8) may be considered a weak notion of MCF.

In the sequel, we omit writing the time-space domain $(0, T) \times [0, 1]^d$ when integrating the Lebesgue measure $dxdt$ or the limiting surface measure $|\nabla\chi|dt$. However, the convolution $*$, for which we reserve the z -variable, is always w. r. t. \mathbb{R}^d .

The next elementary lemma provides the necessary notions and results on abstract minimizing movements schemes.

Lemma 1. *Let (\mathcal{M}, d) be a compact metric space and $E: \mathcal{M} \rightarrow \mathbb{R}$ be continuous. Given $\chi^0 \in \mathcal{M}$ and $h > 0$ consider a sequence $\{\chi^n\}_{n \in \mathbb{N}}$ satisfying*

$$(9) \quad \chi^n \text{ minimizes } \frac{1}{2h} d^2(u, \chi^{n-1}) + E(u) \text{ among all } u \in \mathcal{M}.$$

Then we have for all $t \in \mathbb{N}h$

$$(10) \quad \begin{aligned} & E(\chi(t)) \\ & + \frac{1}{2} \int_0^t \left(\frac{1}{h^2} d^2(\chi(s), \chi(s-h)) + |\partial E(u(s))|^2 \right) ds \leq E(\chi^0). \end{aligned}$$

Here $\chi(t)$ is the piecewise constant interpolation, cf. (3), $u(t)$ is another interpolation (the “variational interpolation”) satisfying

$$(11) \quad \int_0^\infty \frac{1}{2h^2} d^2(u(t), \chi(t)) dt \leq E(\chi^0),$$

$$(12) \quad E(u(t)) \leq E(\chi(t)) \text{ for all } t \geq 0,$$

and $|\partial E(u)|$ is the “metric slope” defined through

$$(13) \quad |\partial E(u)| := \limsup_{v: d(v,u) \rightarrow 0} \frac{(E(u) - E(v))_+}{d(v, u)} \in [0, \infty].$$

The next elementary but crucial lemma establishes that the thresholding scheme is a minimizing movements scheme.

Lemma 2. *Expression (1) satisfies (9) provided we define*

$$(14) \quad \mathcal{M} := \{u: [0, 1]^d \rightarrow [0, 1] \text{ measurable}\},$$

$$(15) \quad E_h(u) := \frac{1}{\sqrt{h}} \int (1 - u) G_h * u dx,$$

$$(16) \quad d_h(u, u') := (2\sqrt{h} \int |G_{\frac{h}{2}} * (u - u')|^2 dx)^{\frac{1}{2}}.$$

Furthermore, (\mathcal{M}, d_h) is a compact metric space and E_h continuous.

We will mostly use (16) in form of

$$(17) \quad \frac{1}{2h} d_h^2(u, u') = \frac{1}{\sqrt{h}} \int |G_{\frac{h}{2}} * (u - u')|^2 dx.$$

The first part of the next lemma provides compactness. The second part contains the (only) way we use the convergence assumption (5); loosely speaking, it ensures convergence of the (oriented) normal down to (spatial) scales of $O(\sqrt{h})$. In particular, it rules out ghost interfaces. Since it will also be used for the variational interpolation, cf. Lemma 1, it is formulated for a $[0, 1]$ -valued sequence $\{u_h\}_{h \downarrow 0}$.

Lemma 3. *i) Consider a sequence $\{\chi_h\}_{h \downarrow 0}$ of $\{0, 1\}$ -valued functions on $(0, T) \times [0, 1]^d$ that satisfies*

$$(18) \quad \text{ess sup}_{t \in (0, T)} E_h(\chi_h(t)) + \int_0^T \frac{1}{2h^2} d_h^2(\chi_h(t), \chi_h(t - h)) dt$$

stays bounded as $h \downarrow 0$,

and that is piecewise constant in the sense of (3). Such a sequence is compact in $L^1((0, T) \times [0, 1]^d)$; any (weak) limit χ is such that $\nabla \chi$ is a bounded measure, equi-integrable in t , with

$$(19) \quad c_0 \int |\nabla \chi| dt \leq \liminf_{h \downarrow 0} \int_0^T E_h(\chi_h(t)) dt.$$

ii) Consider a sequence $\{u_h\}_{h \downarrow 0}$ of $[0, 1]$ -valued functions on $(0, T) \times [0, 1]^d$ and a $\{0, 1\}$ -valued function χ on $(0, T) \times [0, 1]^d$ that satisfies (4) and (5) (with χ_h replaced by u_h) and

$$(20) \quad \text{ess sup}_{t \in (0, T)} E_h(u_h(t)) \quad \text{stays bounded as } h \downarrow 0.$$

Then, as measures on (z, t, x) -space, we have the weak convergences

$$(21) \quad G_1(z) \frac{1}{\sqrt{h}} u_h(t, x) (1 - u_h)(t, x - \sqrt{h}z) dx dt dz$$

$\rightarrow G_1(z) (\nu \cdot z)_+ |\nabla \chi| dt dz,$

$$(22) \quad G_1(z) \frac{1}{\sqrt{h}} (1 - u_h)(t, x) u_h(t, x - \sqrt{h}z) dx dt dz$$

$\rightarrow G_1(z) (\nu \cdot z)_- |\nabla \chi| dt dz.$

The test functions may even have polynomial growth in z .

The next two propositions are at the core and provide the link between (10) and (8). Proposition 1 ensures that the metric d_h , cf. (16), is strong enough to control the right notion of energy of curves in configuration space. Proposition 2 makes sure that it is not too strong so that the metric slope $|\partial E_h|$, cf. (13), controls the gradient of the limiting functional.

Proposition 1. *Suppose that (4) and the conclusion of Lemma 3 hold (with u_h replaced by χ_h). Provided the l. h. s. of (24) is finite, there exists $V \in L^2(|\nabla\chi|dt)$ that is the normal velocity in the sense of*

$$(23) \quad \partial_t \chi = V |\nabla \chi| \quad \text{distributionally,}$$

and that is dominated in the sense of

$$(24) \quad \liminf_{h \downarrow 0} \int_0^T \frac{1}{2h^2} d_h^2(\chi_h(t), \chi_h(t-h)) dt \geq c_0 \int V^2 |\nabla \chi| dt.$$

Proposition 2. *Suppose that the conclusions of Lemma 3 ii) hold. Then there exists $H \in L^2(|\nabla\chi|dt)$ that is the mean curvature in the sense of (6) and that is dominated in the sense of*

$$(25) \quad \liminf_{h \downarrow 0} \int_0^T \frac{1}{2} |\partial E_h(u_h(t))|^2 dt \geq c_0 \int \left(\frac{H}{2}\right)^2 |\nabla \chi| dt.$$

3. PROOFS

We will repeatedly use the (parabolic) scaling of G_h , cf. (2),

$$(26) \quad G_h(z) = \frac{1}{\sqrt{h}^d} G_1\left(\frac{z}{\sqrt{h}}\right)$$

and its semi-group property in form of

$$(27) \quad G_h * G_{h'} = G_{h+h'} \quad \text{in particular} \quad G_{\frac{h}{2}} * G_{\frac{h}{2}} = G_h.$$

The constant $c_0 = \frac{1}{\sqrt{2\pi}}$ appears because of the identity

$$(28) \quad \int G_1(z)(z_1)_+ dz = G_1^{d=1}(0) = c_0,$$

where $G_1^{d=1}(z_1) := \frac{1}{\sqrt{2\pi}} \exp(-\frac{z_1^2}{2})$ denotes the standard Gaussian in a single variable. Indeed, by the factorization of the d -dimensional standard Gaussian into $G_1^{d=1}$ and the $(d-1)$ -dimensional one, and by the normalization of the latter, the integral in (28) reduces to $\int_0^\infty G_1^{d=1} z_1 dz_1$. The formula then follows from writing $z_1 G_1^{d=1} = -\frac{d}{dz_1} G_1^{d=1}$.

PROOF OF THEOREM 1.

Note that Lemma 2 allows to make use of Lemma 1, so that we have (10) with (E, d, χ, u) replaced by (E_h, d_h, χ_h, u_h) . We start with the

l. h. s. of (10), for which we plainly have

$$(29) \quad E_h(\chi^0) \leq c_0 \int |\nabla \chi^0|.$$

Indeed, dropping the index 0, this follows by making the l. h. s. explicit $\frac{1}{\sqrt{h}} \int G_h(z)(1 - \chi)\chi(\cdot - z)dx dz$, which by (26) and $\chi \in \{0, 1\}$ coincides with $\int G_1(z)\frac{1}{\sqrt{h}}(\chi - \chi(\cdot - \sqrt{h}z))_ - dx dz$. It remains to appeal to the mean-value inequality $\int \frac{1}{\sqrt{h}}(\chi - \chi(\cdot - \sqrt{h}z))_ - dx \leq \int (z \cdot \nu)_ - |\nabla \chi|$ and to (28).

Note that because of (10) and (29), (18) is satisfied. Hence we may apply Lemma 3 i), which yields $\chi \in \{0, 1\}$ a. e. and that $\nabla \chi$ is a bounded measure which is equi-integrable in t . By Lemma 3 ii), in view of the theorem's assumption (5), we obtain (21) & (22) with u_h replaced by χ_h , so that we may apply Proposition 1. We now argue that (11) & (12) imply that (4) & (5) hold with χ_h replaced by u_h , so that we may use Proposition 2 also for u_h . Indeed, (5) for u_h follows immediately from (5) for χ_h and (12). We now turn to (4); because it is $[0, 1]$ -valued, the sequence $\{u_h\}_{h \downarrow 0}$ always admits a subsequence that has a weak limit u , so that it remains to argue that $u = \chi$, w. l. o. g. assuming that the entire sequence converges. We momentarily fix $h_0 > 0$ and note that by (27) together with Jensen's inequality we have for all $h \leq 2h_0$ that

$$\begin{aligned} \int |G_{h_0} * (u_h - \chi_h)|^2 dx dt &\leq \int |G_{\frac{h}{2}} * (u_h - \chi_h)|^2 dx dt \\ &\stackrel{(17)}{=} \frac{1}{2\sqrt{h}} \int_0^T d_h^2(u_h(t), \chi_h(t)) dt \stackrel{(11)}{\leq} h\sqrt{h} E_h(\chi^0) \stackrel{(29)}{\leq} c_0 h\sqrt{h} \int |\nabla \chi^0|, \end{aligned}$$

so that by lower-semi continuity of the l. h. s. under weak convergence we obtain $\int |G_{h_0} * (u - \chi)|^2 dx dt = 0$. From letting h_0 tend to zero we obtain the desired $u = \chi$.

Momentarily setting $\rho(t) := E_h(\chi_h(t)) + \frac{1}{2} \int_0^t (\frac{1}{h^2} d_h^2(\chi_h(s), \chi_h(s-h)) + |\partial E_h(u_h(s))|^2) ds$, we note that by definition $\rho(t) = \rho(nh) + \delta(t)$, for $t \in [nh, (n+1)h)$, where $\delta(t) := \frac{1}{2} \int_{nh}^t (\frac{1}{h^2} d_h^2(\chi_h(s), \chi_h(s-h)) + |\partial E_h(u_h(s))|^2) ds$. By (10), $\int_0^T \delta(t) dt \leq h E_h(\chi^0)$. Hence if we multiply (10) in form of $\rho(nh) \leq E_h(\chi^0)$ with $\eta(nh) - \eta((n+1)h)$ for some non-increasing $\eta \in C_0^\infty([0, T])$, we obtain $\int_0^\infty (-\frac{d\eta}{dt}) \rho dt \leq (\eta(0) + h \sup |\frac{d\eta}{dt}|) E_h(\chi^0)$. By an integration by parts and with the choice $\eta(t) = \max\{\min\{\frac{T-t}{\tau}, 1\}, 0\}$, this turns into

$$\begin{aligned} &\frac{1}{\tau} \int_{T-\tau}^T E_h(\chi_h(t)) dt \\ &+ \frac{1}{2} \int_0^{T-\tau} (\frac{1}{h^2} d_h^2(\chi_h(t), \chi_h(t-h)) + |\partial E_h(u_h(t))|^2) dt \leq (1 + \frac{h}{\tau}) E_h(\chi^0). \end{aligned}$$

Passing to the limit $h \downarrow 0$, for the first l. h. s. term, we appeal to (19) in Lemma 3 with $(0, T)$ replaced by $(T - \tau, T)$. For the second l. h. s. term, we apply (24) (note that we may extend the integral down to 0 because of the second item in (3)) and (25), both with $(0, T)$ replaced by $(0, T - \tau)$. For the r. h. s. term, we use (29). Summing up, we obtain

$$\begin{aligned} & \frac{c_0}{\tau} \int_{(T-\tau, T) \times [0, 1]^d} |\nabla \chi| dt \\ & + c_0 \int_{(0, T-\tau) \times [0, 1]^d} \left(V^2 + \left(\frac{H}{2} \right)^2 \right) |\nabla \chi| dt \leq c_0 \int |\nabla \chi^0|. \end{aligned}$$

Dividing by c_0 and letting $\tau \downarrow 0$ yields (8).

Finally, we argue why (23) is sufficient to infer (7). Indeed, by the trivial extension of χ_h to $t \leq 0$, cf. (3), the assumptions (4) & (5) extend to $(-T, T)$, where for (5) we appeal to (29). Likewise, the l. h. s. integral in (24) extends to $(-T, T)$. Hence (23) holds distributionally on $(-T, T) \times [0, 1]^d$, which turns into (7) because of $\chi = \chi^0$ for $t < 0$.

PROOF OF LEMMA 1.

We reproduce the proof of [2, Theorem 3.1.4 & Lemma 3.1.3]. We start with the definition of the variational interpolation u . Since by assumption, (\mathcal{M}, d) is compact and E continuous, for any $n \in \mathbb{N}$ and any $t \in ((n-1)h, nh]$, there exists $u(t)$ that minimizes

$$(30) \quad \frac{d^2(u, \chi^{n-1})}{2(t - (n-1)h)} + E(u) \quad \text{among } u \in \mathcal{M}.$$

W. l. o. g. we may assume that $u(nh) = \chi^n$, cf. (9), so that u is indeed an interpolation of $\{\chi^n\}_{n \in \mathbb{N}}$. Since by comparison with $u = \chi^{n-1}$ we have $E(u(t)) \leq E(\chi^{n-1})$, (12) follows immediately from the way we defined the piecewise linear interpolation, cf. (3).

Fixing $n \in \mathbb{N}$ and introducing

$$(31) \quad e(t) := \min_{u \in \mathcal{M}} \left(\frac{d^2(u, \chi^{n-1})}{2(t - (n-1)h)} + E(u) \right), \quad (n-1)h < t \leq nh,$$

we now establish the two crucial inequalities

$$(32) \quad \begin{aligned} & \frac{d^2(u(s), \chi^{n-1})}{2(s - (n-1)h)(t - (n-1)h)} \leq \frac{e(s) - e(t)}{t - s} \\ & \leq \frac{d^2(u(t), \chi^{n-1})}{2(s - (n-1)h)(t - (n-1)h)}, \quad (n-1)h < s < t \leq nh. \end{aligned}$$

For notational simplicity we consider the case $n = 1$; for any $s, t > 0$ we have by definitions (30) and (31)

$$\begin{aligned} e(s) &= \frac{1}{2s} d^2(u(s), \chi^0) + E(u(s)) \\ &\leq \frac{1}{2s} d^2(u(t), \chi^0) + E(u(t)) = \left(\frac{1}{2s} - \frac{1}{2t}\right) d^2(u(t), \chi^0) + e(t). \end{aligned}$$

Writing $\frac{1}{2s} - \frac{1}{2t} = \frac{t-s}{2st}$ this gives the upper bound in (32) after division by $t - s > 0$. Exchanging the roles of s and t , we likewise get the lower one.

We now argue that

$$(33) \quad |\partial E(u(t))| \leq \frac{d(u(t), \chi^{n-1})}{t - (n-1)h} \quad \text{for } t \in ((n-1)h, nh].$$

Again, for notational simplicity we consider $n = 1$ and give ourselves a $v \in \mathcal{M}$. By the characterizing property (30) of $u(t)$ we have $\frac{1}{2t} d^2(u(t), \chi^0) + E(u(t)) \leq \frac{1}{2t} d^2(v, \chi^0) + E(v)$, so that $E(u(t)) - E(v) \leq \frac{1}{2t} (d(v, \chi^0) - d(u(t), \chi^0))(d(v, \chi^0) + d(u(t), \chi^0))$. By the triangle inequality, this implies

$$E(u(t)) - E(v) \leq d(v, u(t)) \frac{1}{t} (d(u(t), \chi^0) + \frac{1}{2} d(v, u(t))),$$

so that (33) follows from definition (13) of the metric slope.

We now may conclude on (10). By telescoping and according to the piecewise constant interpolation, it is sufficient to establish

$$E(\chi^n) + \frac{1}{2h} d^2(\chi^n, \chi^{n-1}) + \int_{(n-1)h}^{nh} \frac{1}{2} |\partial E(u(s))|^2 ds \leq E(\chi^{n-1}),$$

which according to (33) follows from

$$E(\chi^n) + \frac{1}{2h} d^2(\chi^n, \chi^{n-1}) + \int_{(n-1)h}^{nh} \frac{d^2(u(s), \chi^{n-1})}{2(s - (n-1)h)^2} ds \leq E(\chi^{n-1}),$$

and with help of (31) may be rewritten as

$$(34) \quad e(nh) + \int_{(n-1)h}^{nh} \frac{d^2(u(s), \chi^{n-1})}{2(s - (n-1)h)^2} ds \leq E(\chi^{n-1}).$$

Here comes the argument for (34): We first learn from (32) that

$$(35) \quad ((n-1)h, nh] \ni s \mapsto d^2(u(s), \chi^{n-1}) \quad \text{is monotone increasing}$$

and thus continuous outside of a countable set of s 's. We then learn that e is locally Lipschitz continuous on $((n-1)h, nh]$ and differentiable where (35) is continuous. In particular, we have in those (Lebesgue) almost every time points s , $\frac{de}{ds}(s) = -\frac{1}{2s^2} d^2(u(s), \chi^{n-1})$. Integrating this relationship from some $t \in ((n-1)h, nh]$ to nh we obtain $e(nh) + \int_t^{nh} \frac{1}{2s^2} d^2(u(s), \chi^{n-1}) ds \leq e(t)$. Using the obvious $e(t) \leq E(\chi^{n-1})$,

cf. (31), and letting $t \downarrow (n-1)h$ we obtain (34) by monotone convergence.

We finally turn to (11). According to (35) we have $d^2(u(t), \chi^{n-1}) \leq d^2(\chi^n, \chi^{n-1})$ for $t \in ((n-1)h, nh]$ and thus $\int_0^\infty d^2(u(t), \chi(t)) dt \leq \int_0^\infty d^2(\chi(t), \chi(t-h)) dt$, so that (11) follows from (10).

PROOF OF LEMMA 2.

By the definitions (15) and (17), the latter in conjunction with (27), we have

$$\frac{1}{2h} d_h^2(u, \chi^{n-1}) + E_h(u) = \langle u - \chi^{n-1}, u - \chi^{n-1} \rangle + \langle 1 - u, u \rangle,$$

where we momentarily introduced the bilinear form $\langle u, u' \rangle := \frac{1}{\sqrt{h}} \int u G_h * u' dx$. Since this form is symmetric, we may rewrite the r. h. s. as $\langle u, 1 - 2\chi^{n-1} \rangle + \langle \chi^{n-1}, \chi^{n-1} \rangle$, so that

$$\frac{1}{2h} d_h^2(u, \chi^{n-1}) + E_h(u) = \frac{1}{\sqrt{h}} \int u(1 - 2G_h * \chi^{n-1}) + C,$$

where $C := \langle \chi^{n-1}, \chi^{n-1} \rangle$ does not depend on u . It is now obvious that (1) minimizes $\frac{1}{2h} d_h^2(u, \chi^{n-1}) + E_h(u)$ among all $u \in \mathcal{M}$, cf. (14).

It remains to argue that the metric space (\mathcal{M}, d_h) is compact and E_h continuous. Both follows from the fact that d_h metrizes weak convergence on $\mathcal{M} \subset L^2([0, 1]^d)$. The latter can be seen as follows: In terms of Fourier series, we have $\frac{2}{\sqrt{h}} d_h^2(u, u') = \sum_{k \in 2\pi\mathbb{Z}^d} \exp(-\frac{|hk|^2}{2}) |\mathcal{F}(u - u')|^2(k)$, and note $|\mathcal{F}(u - u')|^2(k) \leq \int (u - u')^2 dx \leq 1$. Hence by dominated convergence, $d_h(u_n, u) \rightarrow 0$ is equivalent to $\mathcal{F}(u_n - u)(k) \rightarrow 0$ for all $k \in 2\pi\mathbb{Z}^d$, which by the L^2 -boundedness of $\{u_n\}_{n \uparrow \infty} \subset \mathcal{M}$ is equivalent to weak convergence.

PROOF OF LEMMA 3.

STEP 1. Some useful inequalities on E_h and d_h . We claim for any $[0, 1]$ -valued function u of space:

$$(36) \quad \int |u - G_h * u| dx \leq 2\sqrt{h} E_h(u),$$

$$(37) \quad E_{h_0}(u) \leq E_h(u) \quad \text{for } h_0 \in \mathbb{N}^2 h,$$

which we claim combine to

$$(38) \quad \int (u - G_{h_0} * u)^2 dx \leq 4\sqrt{h_0} E_h(u) \quad \text{for all } h_0 \geq h.$$

We also claim for any pair of $\{0, 1\}$ -valued functions χ, χ' of space

$$(39) \quad \int |\chi - \chi'| dx \leq \frac{1}{2\sqrt{h}} d_h^2(\chi, \chi') + \sqrt{h} (E_h(\chi) + E_h(\chi')).$$

We first tackle (36); by Jensen's inequality in form of $|u - G_h * u|(x) \leq \int G_h(z) |u(x) - u(x-z)| dz$, and by the definition (15) of E_h which

together with the symmetry of G_h* yields $2\sqrt{h}E_h(u) = \int((1-u)G_h*u + uG_h*(1-u))dx$, (36) follows from the elementary inequality

$$(40) \quad |u - u'| \leq (1-u)u' + u(1-u') \quad \text{for } u, u' \in [0, 1].$$

We now turn to (37) which we (iteratively) establish in the more general form of

$$(41) \quad \sqrt{h_0}E_{h_0} \leq \sqrt{h}E_h + \sqrt{h'}E_{h'} \quad \text{provided } \sqrt{h_0} = \sqrt{h} + \sqrt{h'}.$$

Indeed, by definition (15) and the scaling (26) we have $\sqrt{h}E_h(u) = \int G_1(z)(1-u)(x)u(x - \sqrt{h}z)dx dz$ and by a change of variables in x , $\sqrt{h'}E_{h'}(u) = \int G_1(z)(1-u)(x - \sqrt{h}z)u(x - (\sqrt{h} + \sqrt{h'})z)dx dz$. Hence (41) follows from the elementary inequality

$$(1-u)u'' \leq (1-u)u' + (1-u')u'' \quad \text{for } u, u', u'' \in [0, 1].$$

This is equivalent to $u'(u + u'' - 1) \leq uu''$, which because of $u' \in [0, 1]$ and $uu'' \geq 0$ follows from $u + u'' - 1 \leq uu''$. The latter is equivalent to $u''(1-u) \leq 1-u$, which holds because of $u, u'' \in [0, 1]$.

We now turn to the upgrade (38). We first observe that the l. h. s. is monotone increasing in h_0 , as can be seen by the Fourier representation $\sum_k (1 - \exp(-\frac{h_0 k^2}{2})) |\mathcal{F}u(k)|^2$, where $\mathcal{F}u(k)$, $k \in 2\pi\mathbb{Z}^d$, denotes the Fourier series of u and $\exp(-\frac{h_0 k^2}{2})$ is the Fourier transform of G_{h_0} . Now given $h_0 \geq h$, we write $\sqrt{h_0} = N\sqrt{h} - s$ with $N \in \mathbb{N}$ and $s \in [0, \sqrt{h}]$. By the above monotonicity we have $\int (u - G_{h_0} * u)^2 \leq \int (u - G_{N^2 h} * u)^2 \leq \int |u - G_{N^2 h} * u|$, to which we first apply (36) with h replaced by $N^2 h$ and second apply (37) with $N^2 h$ playing the role of h_0 . Hence we end up with $\int (u - G_{h_0} * u)^2 \leq 2N\sqrt{h}E_h(u)$, which because of $N\sqrt{h} \leq \sqrt{h_0} + \sqrt{h}$ turns into (38).

We finally address (39), which according to the definitions (15)&(16) of E_h and d_h follows from integrating the following inequality in x , appealing to the symmetry of G_h* ,

$$|\chi - \chi'| \leq (\chi - \chi')G_h * (\chi - \chi') + \chi G_h * (1 - \chi) + \chi' G_h * (1 - \chi').$$

Writing $|\chi - \chi'| = (\chi - \chi')^2 = (\chi - \chi')G_h * (\chi - \chi') + (\chi - \chi')(\chi - G_h * \chi) + (\chi - \chi')(\chi' - G_h * \chi')$, we see that the inequality relies on $(\chi - \chi')(\chi - G_h * \chi) \leq \chi(\chi - G_h * \chi) = \chi(1 - G_h * \chi) = \chi G_h * (1 - \chi)$ and on the same inequality with the roles of χ and χ' exchanged.

STEP 2. Modulus of continuity in time: We claim that for every $\{0, 1\}$ -valued function χ of time-space that is piecewise constant in time,

cf. (3), and any time shift $s \in [0, 1]$ we have

$$(42) \quad I(s) \leq C_0 \left\{ \begin{array}{ll} \frac{s}{\sqrt{h}} & \text{for } s \leq h \\ 2\sqrt{h} & \text{for } h \leq s \leq \sqrt{h} \\ 4s & \text{for } \sqrt{h} \leq s \end{array} \right\} \leq 4C_0\sqrt{s},$$

where in this step of the proof, we use the abbreviation

$$(43) \quad \begin{aligned} I(s) &:= \int_{(s,T) \times [0,1]^d} |\chi(t,x) - \chi(t-s,x)| dx dt, \\ C_0 &:= \int_h^T \frac{1}{2h^2} d_h^2(\chi(t), \chi(t-h)) dt + 2 \int_0^T E_h(\chi(t)) dt. \end{aligned}$$

Indeed, for $s \geq h$, we use (39) with $(\chi, \chi') = (\chi(t), \chi(t-s))$ and integrate in $t \in (s, T)$ to obtain

$$(44) \quad I(s) \leq \int_s^T \frac{1}{2\sqrt{h}} d_h^2(\chi, \chi(\cdot - s)) dt + 2\sqrt{h} \int_0^T E_h(\chi) dt,$$

where we write $\chi(\cdot - s)$ for the time-shifted function $(t, x) \mapsto \chi(t-s, x)$. We first use this to treat (42) in case of $s \leq h$: By the piecewise constant interpolation, cf. (3), we have $I(s) \leq \frac{s}{h} I(h)$, into which we insert (44) for $s = h$ in form of $I(h) \leq C_0\sqrt{h}$. We now treat (42) in case of $h \leq s \leq \sqrt{h}$, and first restrict ourselves to $s = Nh$ with $N \in \mathbb{N}$ in order to use the triangle inequality for d_h in form of $d_h^2(\chi, \chi(\cdot - s)) \leq N \sum_{n=1}^N d_h^2(\chi(\cdot - (n-1)h), \chi(\cdot - nh))$ so that we obtain from (44)

$$(45) \quad \begin{aligned} I(s) &\leq \left(\frac{s}{h}\right)^2 \int_h^T \frac{1}{2\sqrt{h}} d_h^2(\chi, \chi(\cdot - h)) dt + 2\sqrt{h} \int_0^T E_h(\chi) dt \\ &\stackrel{(43)}{\leq} C_0 \max\left\{ \frac{s^2}{\sqrt{h}}, \sqrt{h} \right\} = C_0\sqrt{h}. \end{aligned}$$

For the unrestricted range of $h \leq s \leq \sqrt{h}$, we write $s = Nh + s'$ with $N \in \mathbb{N}$ and $s' \in [0, h)$, use $I(s) \leq I(Nh) + I(s')$, and appeal to (45) for the first contribution and to (42) in the previously treated case of $s' \leq h$ for the second contribution. Finally, for (42) in the remaining case of $s \geq \sqrt{h}$, we write $s = N\sqrt{h} + s'$ with $N \in \mathbb{N}$ and $s' \in [0, \sqrt{h})$, use $I(s) \leq NI(\sqrt{h}) + I(s')$, and appeal to the previously treated case of (42) for both terms.

STEP 3. Proof of the compactness statement. From (38) and thanks to the first part of our assumed bound (18), we learn that $G_{h_0} * \chi_h$ is close to χ_h in $L^\infty((0, T), L^2([0, 1]^d))$ (and thus in $L^1((0, T) \times [0, 1]^d)$), as $h_0 \downarrow 0$, uniformly in $h \downarrow 0$. Hence it remains to argue for fixed $h_0 > 0$ that $\{G_{h_0} * \chi_h\}_{h \downarrow 0}$ is compact in L^1 . Because of the convolution in space, and of the equi-integrability following from $G_{h_0} * \chi_h \in [0, 1]$, this follows from a modulus of continuity in time in L^1 that is uniform in $h \downarrow 0$. Thanks to our assumption (18), this holds for χ_h itself by Step 2; it transmits to $G_{h_0} * \chi_h$ by Jensen's inequality.

STEP 4. Before establishing the exact inequality (19), which will be done at the end of Step 6, it is convenient to first argue that $\nabla\chi$ is a bounded measure, equi-integrable in t , under the mere assumption (20). We focus on $\partial_1\chi$, give ourselves a $\zeta \in C_0^\infty((0, T) \times [0, 1]^d)$ and note that as a consequence of (28) we have

$$\begin{aligned} -c_0\partial_1\zeta &= \lim_{h\downarrow 0} \frac{1}{\sqrt{h}} \int_{z_1>0} (\zeta - \zeta(\cdot + \sqrt{h}z))G_1(z)dz \\ &\stackrel{(26)}{=} \lim_{h\downarrow 0} \frac{1}{\sqrt{h}} \int_{z_1>0} (\zeta - \zeta(\cdot + z))G_h(z)dz \end{aligned}$$

uniformly in time-space, where we write $\zeta(\cdot + z)$ for the space-shifted function $(t, x) \mapsto \zeta(t, x + z)$. Hence it follows from (4) that

$$-c_0 \int \partial_1\zeta\chi dxdt = \lim_{h\downarrow 0} \int_{z_1>0} \zeta \frac{1}{\sqrt{h}} (\chi_h - \chi_h(\cdot - z))G_h(z) dxdt dz$$

and thus

$$c_0 \left| \int \partial_1\zeta\chi dxdt \right| \leq \int_0^T \sup_x |\zeta| dt \liminf_{h\downarrow 0} \operatorname{ess\,sup}_t \frac{1}{\sqrt{h}} \int |\chi_h - G_h * \chi_h| dx.$$

According to (36), the second r. h. s. factor is estimated by $2 \operatorname{ess\,sup}_t E_h(\chi_h)$, the $\liminf_{h\downarrow 0}$ of which is bounded by our assumption (20).

STEP 5. Turning to (21) & (22), it is convenient to have the following equi-integrability of the non-negative

$$\begin{aligned} \rho_h(z, t, x) &:= G_1(z) \frac{1}{\sqrt{h}} ((1 - u_h)(t, x)u_h(t, x - \sqrt{h}z) \\ &\quad + u_h(t, x)(1 - u_h)(t, x - \sqrt{h}z)) \end{aligned}$$

in the (non-compact) variables t and z in the sense of

$$\int \exp\left(\frac{3|z|^2}{8}\right) \rho_h(z, t, x) dx dz \leq 2^{d+2} E_h(u_h(t)),$$

cf. (20). Indeed, we first observe that $G_4(z) = \frac{1}{2^d} \exp(\frac{3|z|^2}{8})G_1(z)$, so that by the scaling (26) we obtain $\int \exp(\frac{3|z|^2}{8})\rho_h dx dz = \frac{2^d}{\sqrt{h}} \int ((1 - u_h)G_{4h} * u_h + u_h G_{4h} * (1 - u_h)) dx$, so that by symmetry of G_{4h} and the definition (15) of E_{4h} , $\int \exp(\frac{3|z|^2}{4})\rho_h dx dz = 2^{d+2} E_{4h}(u_h)$, so that it remains to appeal to (37).

STEP 6. Proof of (21) & (22). We focus on (21); according to Step 5, it is sufficient to treat bounded continuous test functions $\zeta(z, t, x)$, which by linearity and splitting into positive and negative part we may assume to be $[0, 1]$ -valued. The statement splits into the local lower

bound

$$(46) \quad \begin{aligned} & \liminf_{h \downarrow 0} \int \zeta G_1(z) \frac{1}{\sqrt{h}} u_h(1 - u_h)(\cdot - \sqrt{h}z) dx dt dz \\ & \geq \int \zeta G_1(z) (\nu \cdot z)_+ |\nabla \chi| dt dz \end{aligned}$$

and the global upper bound

$$(47) \quad \begin{aligned} & \limsup_{h \downarrow 0} \int G_1(z) \frac{1}{\sqrt{h}} u_h(1 - u_h)(\cdot - \sqrt{h}z) dx dt dz \\ & \leq \int G_1(z) (\nu \cdot z)_+ |\nabla \chi| dt dz. \end{aligned}$$

Indeed, splitting a given test function $\zeta = 1 - (1 - \zeta)$, appealing to linearity and using (46) with ζ replaced by $1 - \zeta \in [0, 1]$, we obtain also the local upper bound.

We note that the global upper bound (47) is nothing else than our assumption (5) (with χ_h replaced by u_h): For the l. h. s. this follows from the the scaling (26), the symmetry of G_h* , and the definition (15) of E_h . For the r. h. s. this follows from (28).

Hence it remains to establish (46) by a typical l. s. c. argument: By Fatou's Lemma, it is enough to establish for fixed z

$$\liminf_{h \downarrow 0} \int \zeta \frac{1}{\sqrt{h}} u_h(1 - u_h)(\cdot - \sqrt{h}z) dx dt \geq \int \zeta (\nu \cdot z)_+ |\nabla \chi| dt.$$

Since by Step 4, we already know that $\nabla \chi$ is a bounded measure, we have $(\nu \cdot z)_+ |\nabla \chi| = (\partial_z \chi)_+$. Hence by the definition of the positive part of a measure, and by the equi-integrability in t established in Step 5, it is enough to establish for any non-negative $\zeta \in C_0^\infty((0, T) \times [0, 1]^d)$

$$\liminf_{h \downarrow 0} \int \zeta \frac{1}{\sqrt{h}} u_h(1 - u_h)(\cdot - \sqrt{h}z) dx dt \geq - \int \partial_z \zeta \chi dx dt.$$

Since by assumption (4), the r. h. s. is the limit of

$$\int \frac{1}{\sqrt{h}} (\zeta - \zeta(\cdot + \sqrt{h}z)) u_h = \int \zeta \frac{1}{\sqrt{h}} (u_h - u_h(\cdot - \sqrt{h}z)),$$

the statement follows from the elementary inequality $u - u' \leq u(1 - u')$ that is valid for any $u, u' \in [0, 1]$.

We note that this argument for (46) did not involve the extra assumption (5) and thus applies also under the assumptions of part i) of the lemma. Statement (46) applied to $\zeta = 1$ yields (19) (for the same reason, given above, that (47) is a mere reformulation of (5)).

PROOF OF PROPOSITION 1.

Note that by definition (17) of d_h and (27) we have

$$\begin{aligned} & \int_h^T \frac{1}{2h^2} d_h^2(\chi_h(t), \chi_h(t-h)) dt \\ &= \int_{(h,T) \times [0,1]^d} \frac{1}{h\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h - \chi_h(\cdot - h))|^2 dx dt, \end{aligned}$$

which motivates to introduce the non-negative (bounded) “dissipation measure” on $(0, T) \times [0, 1]^d$ (after possibly passing to a subsequence)

$$(48) \quad \mu := \lim_{h \downarrow 0} \frac{1}{h\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h - \chi_h(\cdot - h))|^2.$$

In fact, we shall establish (24) in the localized form of

$$(49) \quad c_0 V^2 |\nabla \chi| dt \leq \mu.$$

We now give an outline of the proof, which amounts to a better quantification of Step 2 in the proof of Lemma 3. In deriving this estimate on the distribution $\partial_t \chi$, we work on an intermediate time scale, which we fix to be an (eventually small) fraction of the characteristic spatial scale:

$$(50) \quad \tau := \alpha \sqrt{h} \quad \text{for some fixed } \alpha \in (0, \infty).$$

We consider the corresponding increments and their positive and negative parts

$$(51) \quad \delta \chi := \chi_h - \chi_h(\cdot - \tau) \quad \text{and} \quad \delta \chi_{\pm} := \max\{0, \pm \delta \chi\}.$$

Using $|\delta \chi| = \delta \chi_+ + \delta \chi_-$ we then write

$$(52) \quad \begin{aligned} \frac{1}{\sqrt{h}} |\delta \chi| &= \frac{1}{\sqrt{h}} (\delta \chi_+ G_h * \delta \chi_+ + \delta \chi_+ G_h * (1 - \delta \chi_+) \\ &\quad + \delta \chi_- G_h * \delta \chi_- + \delta \chi_- G_h * (1 - \delta \chi_-)). \end{aligned}$$

In Step 1 we will show, in the sense of closeness between distributions,

$$\frac{1}{\sqrt{h}} |\delta \chi| \approx \frac{1}{\sqrt{h}} (\delta \chi G_h * \delta \chi + \delta \chi_+ G_h * (1 - \delta \chi_+) + \delta \chi_- G_h * (1 - \delta \chi_-)).$$

The main idea is to estimate the first r. h. s. by the dissipation measure μ (in Step 2) and to estimate the two last contributions by the surface measure $|\nabla \chi| dt$ (in Step 3). However, this just suffices to show that $\partial_t \chi$ is absolutely continuous w. r. t. $|\nabla \chi| dt$, as is carried out at the beginning of Step 4 on the level of $O(\alpha)$ as $\alpha \downarrow 0$. In order to retrieve (49), we need the finer estimate in Step 3 and look at level $O(\alpha^2)$.

STEP 1. We claim that the mixed term vanishes distributionally:

$$(53) \quad \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} (\delta \chi_+ G_h * \delta \chi_- + \delta \chi_- G_h * \delta \chi_+) = 0.$$

Spelling out the z -integral, we want to show that the distributional limit of

$$(54) \quad \int G_h(z) \frac{1}{\sqrt{h}} (\delta\chi_+ \delta\chi_-(\cdot - z) + \delta\chi_- \delta\chi_+(\cdot - z)) dz$$

vanishes. Fixing some unit vector ν_0 , we split the expression into

$$(55) \quad \int_{\nu_0 \cdot z \geq 0} G_h(z) \frac{1}{\sqrt{h}} (\delta\chi_+ \delta\chi_-(\cdot - z) + \delta\chi_- \delta\chi_+(\cdot - z)) dz$$

and the analogous expression on $\{\nu_0 \cdot z \leq 0\}$. We note that by definition of (51), we have $\delta\chi_+ = \chi_h(1 - \chi_h)(\cdot - \tau)$ and $\delta\chi_- = \chi_h(\cdot - \tau)(1 - \chi_h)$ and thus $\delta\chi_+ \delta\chi_-(\cdot - z) = \chi_h(1 - \chi_h)(\cdot - \tau)\chi_h(\cdot - \tau - z)(1 - \chi_h)(\cdot - z) \leq (1 - \chi_h)(\cdot - \tau)\chi_h(\cdot - \tau - z)$ and likewise $\delta\chi_- \delta\chi_+(\cdot - z) = \chi_h(\cdot - \tau)(1 - \chi_h)\chi_h(\cdot - z)(1 - \chi_h)(\cdot - \tau - z) \leq (1 - \chi_h)\chi_h(\cdot - z)$. Here, with a slight abuse of notation, $\chi(\cdot - \tau - z)(t, x) = \chi(t - \tau, x - z)$. Hence the distributional limit of (55) is dominated by the one of

$$\int_{\nu_0 \cdot z \geq 0} G_h(z) \frac{1}{\sqrt{h}} ((1 - \chi_h)(\cdot - \tau)\chi_h(\cdot - \tau - z) + (1 - \chi_h)\chi_h(\cdot - z)) dz.$$

Since the first contribution differs from the second one just by a (vanishing) time shift, which we may put into the continuous test function, the distributional limit is equal to the weak limit of

$$2 \int_{\nu_0 \cdot z \geq 0} G_h(z) \frac{1}{\sqrt{h}} (1 - \chi_h)\chi_h(\cdot - z) dz,$$

provided the latter exists. Appealing to the scaling (26), according to (22) in Lemma 3 which we test with the characteristic function of the closed set $\{\nu_0 \cdot z \geq 0\}$, the weak limit of this term is dominated by the measure $2 \int_{\nu_0 \cdot z \geq 0} G_1(z)(\nu \cdot z)_- |\nabla\chi| dt dz$. Treating the contribution of $\{\nu_0 \cdot z \leq 0\}$ in a similar way (exchanging the roles of χ and $1 - \chi$), the weak limit of that contribution is dominated by $2 \int_{\nu_0 \cdot z \leq 0} G_1(z)(\nu \cdot z)_+ |\nabla\chi| dt dz$.

Hence we have shown that the weak limit $\lambda \geq 0$ of (54) satisfies

$$(56) \quad \lambda \leq 2 \left(\int_{\nu_0 \cdot z \geq 0} G_1(\nu \cdot z)_- dz + \int_{\nu_0 \cdot z \leq 0} G_1(\nu \cdot z)_+ dz \right) |\nabla\chi| dt.$$

In particular, we have $\lambda \leq 4c_0 |\nabla\chi| dt$, cf. (28), so that there is a $\theta \in L^1(|\nabla\chi| dt)$ with $\lambda = \theta |\nabla\chi| dt$, which allows us the rewrite (56) as

$$\theta \leq 2 \int_{\nu_0 \cdot z \geq 0} G_1(\nu \cdot z)_- dz + 2 \int_{\nu_0 \cdot z \leq 0} G_1(\nu \cdot z)_+ dz$$

$$|\nabla\chi| dt - \text{a. e. and for all } \nu_0 \in \mathbb{R}^d.$$

An elementary separability argument allows to exchange the order between the \forall , so that we may choose $\nu_0 = \nu$, obtaining $\theta \leq 0 |\nabla\chi| dt$ -a. e. and thus $\lambda \leq 0$, yielding (53).

STEP 2. We claim that in a distributional sense

$$(57) \quad \limsup_{h \downarrow 0} \frac{1}{\sqrt{h}} \delta\chi G_h * \delta\chi \leq \alpha^2 \mu.$$

Indeed, by definition (48), we may split this into

$$(58) \quad \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} (\delta\chi G_h * \delta\chi - |G_{\frac{h}{2}} * \delta\chi|^2) = 0 \quad \text{and}$$

$$(59) \quad \limsup_{h \downarrow 0} \left(\frac{1}{\sqrt{h}} |G_{\frac{h}{2}} * \delta\chi|^2 - \alpha^2 \frac{1}{h\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h - \chi_h(\cdot - h))|^2 \right) \leq 0.$$

We start with (59) and assume w. l. o. g. that $\tau = Nh$ for some $N \in \mathbb{N}$ so that by telescoping and the Cauchy-Schwarz inequality in n

$$\begin{aligned} & \frac{1}{\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h - \chi_h(\cdot - \tau))|^2 \\ & \leq N \sum_{n=0}^{N-1} \frac{1}{\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h(\cdot - nh) - \chi_h(\cdot - (n+1)h))|^2. \end{aligned}$$

Appealing to $N = \frac{\alpha}{\sqrt{h}}$, the r. h. s. may be rewritten as

$$\alpha^2 \frac{1}{N} \sum_{n=0}^{N-1} \frac{1}{h\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h(\cdot - nh) - \chi_h(\cdot - (n+1)h))|^2.$$

Note that this is an average of time shifts of $\alpha^2 \frac{1}{h\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h - \chi_h(\cdot - h))|^2$; because of $Nh = O(\sqrt{h}) = o(1)$ all these time shifts are small, so that the (non-negative) expression has the same (bounded) weak limit as $\alpha^2 \frac{1}{h\sqrt{h}} |G_{\frac{h}{2}} * (\chi_h - \chi_h(\cdot - h))|^2$ itself. This yields (59).

We now turn to (58). By the semi-group property (27) and the symmetry of $G_{\frac{h}{2}} *$, we have for a smooth test function ζ

$$\int \zeta (\delta\chi G_h * \delta\chi - |G_{\frac{h}{2}} * \delta\chi|^2) = - \int [\zeta, G_{\frac{h}{2}} *] \delta\chi G_{\frac{h}{2}} * \delta\chi,$$

where $[\zeta, G_{\frac{h}{2}} *]$ denotes the commutator of multiplying with ζ and convolving with $G_{\frac{h}{2}}$. Hence by the boundedness of $\frac{1}{\sqrt{h}} |G_{\frac{h}{2}} * \delta\chi|^2$ as a sequence of measures, which follows from (59), it is enough to establish

$$\lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int |[\zeta, G_{\frac{h}{2}} *] \delta\chi|^2 = 0.$$

We spell out the integrand:

$$([\zeta, G_{\frac{h}{2}} *] \delta\chi)(t, x) = \int G_{\frac{h}{2}}(z) (\zeta(t, x) - \zeta(t, x - z)) \delta\chi(t, x - z) dz,$$

so that $|[\zeta, G_{\frac{h}{2}} *] \delta\chi|(t, x) \leq \sup |\nabla\zeta| \int G_{\frac{h}{2}}(z) |z| |\delta\chi(t, x - z)| dz$ and thus

$$\frac{1}{\sqrt{h}} \int |[\zeta, G_{\frac{h}{2}} *] \delta\chi|^2 dx \leq \frac{1}{\sqrt{h}} (\sup |\nabla\zeta| \int G_{\frac{h}{2}}(z) |z| dz)^2 \int |\delta\chi|^2 dx dt.$$

By the scaling (26), the r. h. s. is $O(\sqrt{h})$ and thus vanishing.

STEP 3. For given unit vector ν_0 and $V_0 \in (0, \infty)$ we claim that in a distributional sense

$$(60) \quad \limsup_{h \downarrow 0} \frac{1}{\sqrt{h}} \left(\delta\chi_+ G_h * (1 - \delta\chi_+) + \delta\chi_- G_h * (1 - \delta\chi_-) \right. \\ \left. - 2 \int_{z \cdot \nu_0 > \alpha V_0} G_1(z) dz |\delta\chi| \right) \leq 2 \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1(z) |z \cdot \nu| |\nabla\chi| dt.$$

We split this into three steps: 1) The l. h. s. may be substituted according to

$$(61) \quad \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \left(\delta\chi_{\pm} G_h * (1 - \delta\chi_{\pm}) \right. \\ \left. - \int_{z \cdot \nu_0 \geq 0} G_h(z) |\delta\chi_{\pm} - \delta\chi_{\pm}(\cdot - z)| dz \right) = 0.$$

2) The integrand, which is a second-order difference, satisfies the two inequalities

$$(62) \quad \left\{ \begin{array}{l} |\delta\chi_+ - \delta\chi_+(\cdot - z)| + |\delta\chi_- - \delta\chi_-(\cdot - z)| \\ \leq \left\{ \begin{array}{l} |\chi_h - \chi_h(\cdot - z)| + |\chi_h(\cdot - \tau) - \chi_h(\cdot - \tau - z)| \\ |\delta\chi| + |\delta\chi(\cdot - z)| \end{array} \right\}, \end{array} \right.$$

where the first inequality is “space-like” and we use it on the set $\{0 \leq z \cdot \nu_0 \leq \tau V_0\}$, while the second one is “time-like” and we use it on the complement $\{z \cdot \nu_0 > \tau V_0\}$. 3) We finally argue that

$$(63) \quad \limsup_{h \downarrow 0} \frac{1}{\sqrt{h}} \int_{0 \leq z \cdot \nu_0 \leq \tau V_0} G_h(z) (|\chi_h - \chi_h(\cdot - z)| + |\chi_h(\cdot - \tau) \\ - \chi_h(\cdot - z - \tau)|) dz \leq 2 \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1(z) |z \cdot \nu| |\nabla\chi| dt dz,$$

$$(64) \quad \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \left(\int_{z \cdot \nu_0 > \tau V_0} G_h(z) (|\delta\chi| + |\delta\chi(\cdot - z)|) dz \right. \\ \left. - 2 \frac{1}{\sqrt{h}} \int_{z \cdot \nu_0 > \alpha V} G_1(z) dz |\delta\chi| \right) = 0.$$

We start with (62); the second inequality is obvious by the triangle inequality in form of $|\delta\chi_{\pm} - \delta\chi_{\pm}(\cdot - z)| \leq \delta\chi_{\pm} + \delta\chi_{\pm}(\cdot - z)$ and by $\delta\chi_+ + \delta\chi_- = |\delta\chi|$. In view of the definition (51), the first inequality in (62) follows from the elementary inequality

$$(65) \quad |(a - a')_+ - (b - b')_+| + |(a - a')_- - (b - b')_-| \\ \leq |a - b| + |a' - b'|,$$

which can be seen by distinguishing two cases: Case 1): $(a - a')(b - b') \geq 0$, w. l. o. g. by symmetry $(a, a', b, b') \rightsquigarrow (-a, -a', -b, -b')$ we may assume $a - a', b - b' \geq 0$, in which case (65) turns into the obvious

$|(a - a') - (b - b')| \leq |a - b| + |a' - b'|$. Case 2): $(a - a')(b - b') \leq 0$, by the same symmetry we may assume $a - a' \geq 0 \geq b - b'$, in which case (65) turns into the obvious $(a - a') + (b' - b) \leq |a - b| + |a' - b'|$.

We now argue for (63) & (64). Putting the vanishing shift τ in the time variable on the continuous test function, (63) follows once we argue that

$$(66) \quad \begin{aligned} & \limsup_{h \downarrow 0} \frac{1}{\sqrt{h}} \int_{0 \leq z \cdot \nu_0 \leq \tau V_0} G_h(z) |\chi_h - \chi_h(\cdot - z)| dz \\ & \leq \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1(z) |z \cdot \nu| |\nabla \chi| dt dz, \end{aligned}$$

which because of non-negativity implies that the l. h. s. admits a limit as a measure on $(0, T) \times [0, 1]^d$. Appealing to scaling (26) in form of $G_h(z) dz = G_1(\frac{z}{\sqrt{h}}) d\frac{z}{\sqrt{h}}$ and noting that $z \cdot \nu \leq \tau V_0$ is equivalent to $\frac{z}{\sqrt{h}} \cdot \nu \leq \alpha V_0$, cf. (50), (66) follows from taking the sum of (21) and (22), appealing to (40), testing with the characteristic function of the closed set $\{0 \leq z \cdot \nu_0 \leq \alpha V_0\}$, and integrating out z .

We now turn to (64) and note that it follows from

$$\lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int_{z \cdot \nu_0 > \tau V} G_h(z) (|\delta \chi(\cdot - z)| - |\delta \chi|) dz = 0,$$

which testing with $\zeta \in C_0^\infty((0, T) \times [0, 1]^d)$ assumes the form

$$\lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int_{z \cdot \nu_0 > \tau V} G_h(z) (\zeta(\cdot + z) - \zeta) |\delta \chi| dx dt dz = 0.$$

This holds, since the integral can be estimated by

$$(67) \quad \begin{aligned} & \sup |\nabla \zeta| \int \frac{|z|}{\sqrt{h}} G_h(z) |\delta \chi| dx dt dz \\ & \stackrel{(26), (51)}{=} \sup |\nabla \zeta| \int |z| G_1(z) dz \int |\chi_h - \chi_h(\cdot - \tau)| dx dt, \end{aligned}$$

and the last contribution vanishes in the limit since τ does, and since $\{\chi_h\}_{h \downarrow 0}$ is compact in $L^1((0, T) \times [0, 1]^d)$, cf. part i) of Lemma 3.

We finally address (61) and focus on the $+$ -part. We first argue that

$$(68) \quad \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} (\delta \chi_+ G_h * (1 - \delta \chi_+) - (1 - \delta \chi_+) G_h * \delta \chi_+) = 0.$$

Spelling out the z -integral, and using that G_h is even, this follows from

$$(69) \quad \begin{aligned} & \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int G_h(z) \delta \chi_+(1 - \delta \chi_+)(\cdot - z) dz \\ & = \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int G_h(z) \delta \chi_+(\cdot + z)(1 - \delta \chi_+) dz. \end{aligned}$$

Since the second function differs from the first just by a spatial shift of z , the limits coincide provided the l. h. s. limit is finite, which by

the non-negativity of the functions follows if their integral remains bounded. The l. h. s. integral indeed remains bounded since $\delta\chi_+(1 - \delta\chi_+)(\cdot - z) \leq \chi_h(1 - \chi_h)(\cdot - z) + (1 - \chi_h)(\cdot - \tau)\chi_h(\cdot - \tau - z)$, which follows from $\delta\chi_+ = \chi_h(1 - \chi_h)(\cdot - \tau)$, and since the integral of the first summand remains bounded by (21), whereas the integral of the second summand is oblivious to the (vanishing) time shift and thus remains bounded by (22).

Equipped with (68), we now may substitute $\frac{1}{\sqrt{h}}\delta\chi_+G_h * (1 - \delta\chi_+)$ by $\frac{1}{2\sqrt{h}}(\delta\chi_+G_h * (1 - \delta\chi_+) + (1 - \delta\chi_+)G_h * \delta\chi_+)$, which we may write as

$$\frac{1}{2\sqrt{h}} \int G_h(z) |\delta\chi_+ - \delta\chi_+(\cdot - z)| dz,$$

where we used for any $a, b \in \{0, 1\}$ that $a(1 - b) + (1 - a)b = |a - b|$. Hence in order to obtain (61), we need the two function sequences

$$\frac{1}{\sqrt{h}} \int_{\pm z \cdot \nu_0 \geq 0} G_h(z) |\delta\chi_+ - \delta\chi_+(\cdot - z)| dz$$

to have the same limit. Again by the evenness of G_h , these two functions only differ by a spatial shift z . The same argument as for (69) shows that the limits agree.

STEP 4. Conclusion. We start from the identity (52) in form of

$$\begin{aligned} \frac{1}{\sqrt{h}} |\delta\chi| &= \frac{1}{\sqrt{h}} (\delta\chi G_h * \delta\chi \\ &\quad + \delta\chi_+ G_h * \delta\chi_- + \delta\chi_- G_h * \delta\chi_+ \\ &\quad + \delta\chi_+ G_h * (1 - \delta\chi_+) + \delta\chi_- G_h * (1 - \delta\chi_-)), \end{aligned}$$

or rather, using $2 \int_{z \cdot \nu_0 \geq 0} G_1 dz = 1$,

$$\begin{aligned} 2 \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1 dz \frac{1}{\sqrt{h}} |\delta\chi| &= \frac{1}{\sqrt{h}} (\delta\chi G_h * \delta\chi \\ &\quad + \delta\chi_+ G_h * \delta\chi_- + \delta\chi_- G_h * \delta\chi_+ \\ &\quad + \delta\chi_+ G_h * (1 - \delta\chi_+) + \delta\chi_- G_h * (1 - \delta\chi_-) - 2 \int_{z \cdot \nu_0 > \alpha V_0} G_1 dz |\delta\chi|). \end{aligned}$$

We note that by an elementary lower-semi-continuity argument based on the definitions (50) and (51), we have the distributional inequality $\alpha |\partial_t \chi| \leq \liminf_{h \downarrow 0} \frac{1}{\sqrt{h}} |\delta\chi|$, provided the r. h. s. is a finite measure. Hence we obtain from (53), (57), and (60) the distributional inequality

$$\begin{aligned} 2\alpha \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1 dz |\partial_t \chi| &\leq \alpha^2 \mu \\ (70) \qquad \qquad \qquad &+ 2 \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1 |z \cdot \nu| dz |\nabla \chi| dt, \end{aligned}$$

which in particular shows that $\partial_t \chi$ is a measure. Letting $V_0 \uparrow \infty$ and appealing to (28), this yields in particular $\alpha |\partial_t \chi| \leq \alpha^2 \mu + 4c_0 |\nabla \chi| dt$ which we divide by α :

$$|\partial_t \chi| \leq \alpha \mu + \frac{4}{\alpha} c_0 |\nabla \chi| dt.$$

Letting $\alpha \downarrow 0$, we learn from the latter estimate that null sets of $|\nabla \chi| dt$ are null sets of $\partial_t \chi$, so that there exists $V \in L^1(|\nabla \chi| dt)$ such that (23) holds.

Since $|\partial_t \chi| = |V| |\nabla \chi| dt$ is absolutely continuous w. r. t. to $|\nabla \chi| dt$, (70) even holds with μ replaced by its absolutely continuous part μ' w. r. t. $|\nabla \chi| dt$. Writing $\mu' = \theta |\nabla \chi| dt$ with $\theta \in L^1(|\nabla \chi| dt)$, (70) assumes the form

$$2\alpha \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1 dz |V| \leq \alpha^2 \theta + 2 \int_{0 \leq z \cdot \nu_0 \leq \alpha V_0} G_1 |z \cdot \nu| dz \quad |\nabla \chi| dt\text{-a. e.}$$

As in Step 1, a separability argument now allows to choose $\nu_0 = \nu$, so that by radial symmetry of G_1 , the above assumes the form

$$2\alpha \int_{0 \leq z_1 \leq \alpha V_0} G_1 dz |V| \leq \alpha^2 \theta + 2 \int_{0 \leq z_1 \leq \alpha V_0} G_1 z_1 dz \quad |\nabla \chi| dt\text{-a. e.}$$

Dividing by α^2 and momentarily writing $\alpha' := \alpha V_0$, this turns into

$$2 \frac{V_0}{\alpha'} \int_{0 \leq z_1 \leq \alpha'} G_1 dz |V| \leq \theta + 2 \frac{V_0^2}{\alpha'^2} \int_{0 \leq z_1 \leq \alpha'} G_1 z_1 dz \quad |\nabla \chi| dt\text{-a. e.}$$

We now appeal to the limiting relations (which follow from factorizing G_1 into the $(d-1)$ -dimensional standard Gaussian and $G_1^{d=1}$)

$$\begin{aligned} \lim_{\alpha' \downarrow 0} \frac{1}{\alpha'} \int_{0 \leq z_1 \leq \alpha'} G_1 dz &= G_1^{d=1}(0) = c_0, \\ \lim_{\alpha' \downarrow 0} \frac{1}{\alpha'^2} \int_{0 \leq z_1 \leq \alpha'} G_1 z_1 dz &= \frac{1}{2} G_1^{d=1}(0) = \frac{c_0}{2}, \end{aligned}$$

to see that the above turns into

$$2c_0 V_0 |V| \leq \theta + c_0 V_0^2 \quad |\nabla \chi| dt\text{-a. e.}$$

Again, by a separability argument for V_0 , we may assume $V_0 = |V|$ so that the above yields (49) in form of $c_0 V^2 \leq \theta$.

PROOF OF PROPOSITION 2.

STEP 1. Metric slope and functional derivative. We claim the following relation between the metric slope $|\partial E(u)|$ of a functional E on \mathcal{M} , cf. (14), at a configuration u , and its first variation $\delta E(u) \cdot \xi$ in direction of a smooth vector field ξ :

$$(71) \quad \frac{1}{2} |\partial E(u)|^2 \geq \delta E(u) \cdot \xi - \frac{1}{2} (\delta d(u, \cdot)(u) \cdot \xi)^2.$$

As usual, first variation δ is defined by considering the curve $s \mapsto u_s$ of configurations characterized via the transport equation (to be interpreted distributionally or solved explicitly with help of the flow map Φ_s via $u_s \circ \Phi_s = u$)

$$(72) \quad \frac{\partial u_s}{\partial s} + \xi \cdot \nabla u_s = 0 \quad \text{and} \quad u_{s=0} = u,$$

and setting

$$(73) \quad \delta E(u) \cdot \xi := \left. \frac{d}{ds} \right|_{s=0} E(u_s) \quad \text{and} \quad \delta d(u, \cdot)(u) \cdot \xi := \left. \frac{d}{ds} \right|_{s=0} d(u, u_s),$$

with the understanding that both derivatives exist (and define linear functionals in ξ which is the case for $E = E_h$ and $d(u, \cdot) = d_h(u, \cdot)$, cf. Steps 2 and 3). Inequality (71) is then a direct consequence of the definition $|\partial E(u)|$, cf. (13), which yields

$$\begin{aligned} |\partial E(u)| &\geq \limsup_{s \downarrow 0} \frac{(E(u) - E(u_s))_+}{d(u, u_s)} \\ &\geq \frac{\lim_{s \downarrow 0} \frac{1}{s} (E(u_s) - E(u))}{\lim_{s \downarrow 0} \frac{1}{s} d(u, u_s)} = \frac{\delta E(u) \cdot \xi}{\delta d(u, \cdot)(u) \cdot \xi}, \end{aligned}$$

and Young's inequality.

STEP 2. Representation of $\delta E_h(u) \cdot \xi$; we claim:

$$(74) \quad \begin{aligned} \delta E_h(u) \cdot \xi &= \frac{1}{\sqrt{h}} \int \left(\nabla \cdot \xi ((1-u)G_h * u + uG_h * (1-u)) \right. \\ &\quad \left. + u[\xi, \nabla G_h *](1-u) \right), \end{aligned}$$

where $[\xi, \nabla G_h *] = \sum_{i=1}^d [\xi_i, \partial_i G_h *]$ denotes the commutator of multiplying with ξ and convolving with ∇G_h . In checking this formula we may by approximation assume that u is smooth; by definition (72) & (73) of δ we obtain from the definition (15) of E_h

$$\delta E_h(u) \cdot \xi = \frac{1}{\sqrt{h}} \int ((\xi \cdot \nabla u)G_h * u - (1-u)G_h * (\xi \cdot \nabla u)),$$

which by the symmetry of $G_h *$ we rewrite as

$$\delta E_h(u) \cdot \xi = -\frac{1}{\sqrt{h}} \int (\xi \cdot \nabla(1-u)G_h * u + \xi \cdot \nabla u G_h * (1-u)).$$

We write $\xi \cdot \nabla u = \nabla \cdot (u\xi) - u\nabla \cdot \xi$ and $\xi \cdot \nabla(1-u) = \nabla \cdot ((1-u)\xi) - (1-u)\nabla \cdot \xi$, so that (74) reduces to the identity

$$\begin{aligned} & - \int (\nabla \cdot ((1-u)\xi)G_h * u + \nabla \cdot (u\xi)G_h * (1-u)) \\ & = \int u[\xi, \nabla G_h *](1-u), \end{aligned}$$

which follows from integration by parts and the anti-symmetry of $\nabla G_h *$.

STEP 3. Representation of $\delta d_h(u, \cdot)(u) \cdot \xi$:

$$\begin{aligned}
& \frac{1}{2} (\delta d_h(u, \cdot)(u) \cdot \xi)^2 \\
&= \sqrt{h} \int \left(u \xi \cdot \nabla^2 G_h * ((1-u)\xi) - u \xi \cdot \nabla G_h * ((1-u)\nabla \cdot \xi) \right. \\
(75) \quad & \left. + u \nabla \cdot \xi \nabla G_h * ((1-u)\xi) - u \nabla \cdot \xi G_h * ((1-u)\nabla \cdot \xi) \right).
\end{aligned}$$

In the notation of Step 2, $\frac{1}{2} (\delta d_h(u, \cdot)(u) \cdot \xi)^2 = \frac{1}{2} \left(\frac{d}{ds} \Big|_{s=0} d_h(u, u_s) \right)^2 = \frac{1}{4} \frac{d^2}{ds^2} \Big|_{s=0} d_h^2(u, u_s)$, so that by definition (17) of d_h and by (72), we have $\frac{1}{2} (\delta d_h(u, \cdot)(u) \cdot \xi)^2 = \sqrt{h} \int \frac{\partial u_s}{\partial s} \Big|_{s=0} G_h * \frac{\partial u_s}{\partial s} \Big|_{s=0} = \sqrt{h} \int \xi \cdot \nabla u G_h * (\xi \cdot \nabla u)$. Rewriting the second factor $\xi \cdot \nabla u = \nabla \cdot (u\xi) - u \nabla \cdot \xi$ and using the symmetry of $G_h *$, an integration by parts, and $-\nabla u = \nabla(1-u)$, we obtain

$$\begin{aligned}
& \frac{1}{2} (\delta d_h(u, \cdot)(u) \cdot \xi)^2 \\
&= \sqrt{h} \int \left(u \xi \cdot \nabla G_h * (\xi \cdot \nabla(1-u)) + u \nabla \cdot \xi G_h * (\xi \cdot \nabla(1-u)) \right).
\end{aligned}$$

We write $\xi \cdot \nabla(1-u) = \nabla \cdot ((1-u)\xi) - (1-u)\nabla \cdot \xi$ to obtain (75).

STEP 4. Passage to the limit in δE_h ; we claim that

$$(76) \quad \lim_{h \downarrow 0} \int_0^T \delta E_h(u_h) \cdot \xi dt = c_0 \int (\nabla \cdot \xi - \nu \cdot \nabla \xi \nu) |\nabla \chi| dt.$$

According to (74), we may split into two statements. The first statement is

$$\begin{aligned}
& \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int \nabla \cdot \xi \left((1-u_h) G_h * u_h + u_h G_h * (1-u_h) \right) \\
&= 2c_0 \int \nabla \cdot \xi |\nabla \chi| dt,
\end{aligned}$$

which is an immediate consequence of testing (21) & (22) with $\nabla \cdot \xi$, appealing to the scaling (26) and to the formula (28). The second statement is

$$\begin{aligned}
& \lim_{h \downarrow 0} \frac{1}{\sqrt{h}} \int u_h [\xi, \nabla G_h *] (1-u_h) \\
(77) \quad &= -c_0 \int (\nu \cdot \nabla \xi \nu + \nabla \cdot \xi) |\nabla \chi| dt,
\end{aligned}$$

for which we now give the argument. Spelling out

$$\begin{aligned}
& ([\xi, \nabla G_h *] (1-u_h))(t, x) \\
&= \int (\xi(t, x) - \xi(t, x-z)) \cdot \nabla G_h(z) (1-u_h)(t, x-z) dz,
\end{aligned}$$

we see that

$$\begin{aligned}
& |([\xi, \nabla G_h *](1 - u_h))(t, x) \\
& \quad - \int \nabla G_h(z) \cdot \nabla \xi(t, x) z (1 - u_h)(t, x - z) dz| \\
(78) \quad & \leq \frac{1}{2} \sup |\nabla^2 \xi| \int |z|^2 |\nabla G_h(z)| (1 - u_h)(t, x - z) dz.
\end{aligned}$$

Appealing to (26) in form of $\nabla G_h(z) dz = \frac{1}{\sqrt{h}} \nabla G_1(\frac{z}{\sqrt{h}}) d\frac{z}{\sqrt{h}}$, we learn that the limit of the contribution of the main term can be computed by testing (21) with $\frac{\nabla G_1(z) \cdot \nabla \xi(t, x) z}{G_1(z)} = -z \cdot \nabla \xi(t, x) z$ (which is of polynomial growth in z); it assumes the value

$$\int \nabla G_1 \cdot \nabla \xi z (\nu \cdot z)_+ |\nabla \chi| dt dz,$$

which yields (77) by formula (83) below. The contribution of the r. h. s. error term in (78) is vanishing of $O(\sqrt{h})$, as follows from appealing to (21) tested with $\frac{|\nabla G_1(z)| |z|^2}{G_1(z)} = |z|^3$.

STEP 5. Passage to the limit in δd_h ; we claim

$$(79) \quad \lim_{h \downarrow 0} \frac{1}{2} \int_0^T (\delta d_h(u_h, \cdot)(u_h) \cdot \xi)^2 dt = c_0 \int (\xi \cdot \nu)^2 |\nabla \chi| dt.$$

According to (75), this statement may be split into a leading-order statement

$$(80) \quad \lim_{h \downarrow 0} \sqrt{h} \int u_h \xi \cdot \nabla^2 G_h * ((1 - u_h) \xi) dx dt = c_0 \int (\xi \cdot \nu)^2 |\nabla \chi| dt,$$

and the higher-order statements

$$\begin{aligned}
(81) \quad & \int u_h \xi \cdot \nabla G_h * ((1 - u_h) \nabla \cdot \xi) dx dt = O(1), \\
& \int u_h \nabla \cdot \xi \nabla G_h * ((1 - u_h) \xi) dx dt = O(1), \\
& \frac{1}{\sqrt{h}} \int u_h \nabla \cdot \xi G_h * ((1 - u_h) \nabla \cdot \xi) dx dt = O(1).
\end{aligned}$$

The statement (80) itself splits into the main part

$$\lim_{h \downarrow 0} \sqrt{h} \int \xi \cdot (u_h \nabla^2 G_h * (1 - u_h)) \xi dx dt = c_0 \int (\xi \cdot \nu)^2 |\nabla \chi| dt$$

and the higher-order commutator

$$(82) \quad \int u_h \xi \cdot [\xi, \nabla^2 G_h] * (1 - u_h) dx dt = O(1).$$

The main part follows from appealing to (26) in form of $\sqrt{h} \nabla^2 G_h(z) dz = \frac{1}{\sqrt{h}} \nabla^2 G_1(\frac{z}{\sqrt{h}}) d\frac{z}{\sqrt{h}}$, testing (21) with $\frac{\xi(t, x) \cdot \nabla^2 G_1(z) \xi(t, x)}{G_1(z)} = (\xi(t, x) \cdot z)^2 - |\xi(t, x)|^2$, and appealing to formula (84). The estimate of the error

term (82) follows from estimating the integrand by $\sup |\xi| \sup |\nabla \xi| \int |z| |\nabla^2 G_h(z)| u_h(t, x) (1 - u_h)(t, x - z) dz$, and then using the scaling (26) further by

$$\sup |\xi| \sup |\nabla \xi| \int |z| |\nabla^2 G_1(z)| \frac{1}{\sqrt{h}} u_h(t, x) (1 - u_h)(t, x - \sqrt{h}z) dz,$$

so that another application of (21) yields (82). Statement (81) and the other two higher-order estimates follow along the same lines: For instance, the integrand in (81) is $\leq \sup |\xi| \sup |\nabla \cdot \xi| \int |\nabla G_h(z)| u_h(t, x) (1 - u_h)(t, x - z) dz$. By rescaling, $\int |\nabla G_h(z)| u_h(t, x) (1 - u_h)(t, x - z) dz = \int |z| G_1(z) \frac{1}{\sqrt{h}} u_h(t, x) (1 - u_h)(t, x - \sqrt{h}z) dz$, which is $O(1)$ by (21).

STEP 6. Conclusion. By Riesz' representation theorem in $L^2(|\nabla \chi| dt)$ and an approximation argument in the (arbitrary) smooth vector field ξ , the statement of Proposition 2 is a consequence of

$$\liminf_{h \downarrow 0} \int_0^T \frac{1}{2} |\partial E_h(u_h)|^2 dt \geq c_0 \int (\nabla \cdot \xi - \nu \cdot \nabla \xi \nu - (\xi \cdot \nu)^2) |\nabla \chi| dt.$$

The latter follows starting from the inequality (71) for $(E, d, u) = (E_h, d_h, u_h(t))$, integrating in $t \in (0, T)$, and appealing to (76) and (79) to pass to the limit on the r. h. s. .

STEP 7. Two formulas: For any unit vector ν , any matrix A , and any vector ξ , we have

$$(83) \quad - \int \nabla G_1(z) \cdot Az (\nu \cdot z)_+ dz = c_0 (\nu \cdot A \nu + \text{tr} A),$$

$$(84) \quad \int \xi \cdot \nabla^2 G_1(z) \xi (\nu \cdot z)_+ dz = c_0 (\xi \cdot \nu)^2.$$

Since $G_1(z) = -z G_1(z)$, for the first formula we may assume that A is symmetric; by linearity we may assume that $A = e \otimes e$ for some unit vector e ; by radial symmetry of G_1 , it thus remains to show

$$- \int \partial_1 G_1 z_1 (\nu \cdot z)_+ dz = c_0 (\nu_1^2 + 1),$$

which by one integration by parts, taking into account (28), reduces to

$$\int_{\nu \cdot z > 0} G_1 z_1 dz = c_0 \nu_1,$$

which in view of $G_1 z_1 = -\partial_1 G_1 = -\nabla \cdot (G_1 e_1)$ by the divergence theorem reduces to

$$(85) \quad \int_{\nu \cdot z = 0} G_1 = c_0,$$

which follows since $c_0 = G_1^{d-1}(0)$. We now turn to (84). By radial symmetry of G_1 and homogeneity, it suffices to show

$$(86) \quad \int \partial_1^2 G_1(\nu \cdot z)_+ dz = c_0 \nu_1^2,$$

which by two integration by parts reduces to (85).

REFERENCES

- [1] Fred Almgren, Jean E. Taylor, and Lihe Wang. Curvature-driven flows: a variational approach. *SIAM Journal on Control and Optimization*, 31(2):387–438, 1993.
- [2] Luigi Ambrosio, Nicola Gigli, and Giuseppe Savaré. *Gradient flows in metric spaces and in the space of probability measures*. Birkhäuser, 2008.
- [3] Selim Esedoğlu and Felix Otto. Threshold dynamics for networks with arbitrary surface tensions. *Communications on Pure and Applied Mathematics*, 68(5):808–864, 2015.
- [4] Lawrence C. Evans. Convergence of an algorithm for mean curvature motion. *Indiana University Mathematics Journal*, 42(2):533–557, 1993.
- [5] Tim Laux and Felix Otto. Convergence of the thresholding scheme for multiphase mean-curvature flow. *Calculus of Variations and Partial Differential Equations*, 55(5):1–74, 2016.
- [6] Tim Laux and Felix Otto. Brakke’s inequality for the thresholding scheme. *arXiv:1708.03071 [math.AP]*, 2017.
- [7] Stephan Luckhaus and Thomas Sturzenhecker. Implicit time discretization for the mean curvature flow equation. *Calculus of Variations and Partial Differential Equations*, 3(2):253–271, 1995.
- [8] Barry Merriman, James K. Bence, and Stanley J. Osher. Diffusion generated motion by mean curvature. CAM Report 92-18, 1992. Department of Mathematics, University of California, Los Angeles.
- [9] Peter W. Michor and David Mumford. Riemannian geometries on spaces of plane curves. *Journal of the European Mathematical Society*, 8(1):1–48, 2006.
- [10] Tiago Salvador and Selim Esedoğlu. A simplified threshold dynamics algorithm for isotropic surface energies. *Journal of Scientific Computing*, pages 1–22, 2018.