

**Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig**

**Rigidity for the hyperbolic
Monge-Ampere equation**

(revised version: May 2004)

by

Chun-Chi Lin

Preprint no.: 53

2003



RIGIDITY FOR THE HYPERBOLIC MONGE-AMPÈRE EQUATION

CHUN-CHI LIN

ABSTRACT. Some properties of nonlinear partial differential equations are naturally associated with the geometry of sets in the space of matrices. In this paper, we consider the model case when the compact set K is contained in the hyperboloid \mathcal{H}_{-1} , where $\mathcal{H}_{-1} \subset \mathbb{M}_{sym}^{2 \times 2}$, the symmetric 2×2 matrices. The hyperboloid \mathcal{H}_{-1} is generated by two families of rank-one lines and related to the hyperbolic Monge-Ampère equation $\det \nabla^2 u = -1$. For some compact subsets $K \subset \mathcal{H}_{-1}$ containing a rank-one connection, we show the rigidity property of K by imposing proper topology for the convergence of approximate solutions and affine boundary conditions.

1. INTRODUCTION

This paper is motivated by the rigidity problems studied by Chlebík and Kirchheim in [5] and [14]. In connection with mathematical models of crystalline microstructures ([2], [6], [7]), one is interested in exact and approximate solutions of the partial differential relations $\nabla f \in K$ a.e. in Ω , where $f : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a Lipschitz map on a bounded domain Ω and K is a compact set of matrices in $\mathbb{M}^{m \times n}$. We say that the compact set K is rigid for exact solutions if each Lipschitz map f with $\nabla f \in K$ a.e. in Ω is necessarily an affine map. The compact set K is rigid for approximate solutions if, for each convergent subsequence of Lipschitz maps $f^{(j)}$ with $\text{dist}(\nabla f^{(j)}, K) \rightarrow 0$ in measure in Ω , the limit map f is necessarily an affine map. We say that a set K contains a rank-one connection if there are two matrices $A, B \in K$ such that $\text{rk}(A - B) \leq 1$. Whenever a set K contains a rank-one connection, one may construct an exact or approximate solution such that ∇f oscillates strongly. In this situation, which corresponds to a loss of ellipticity, one cannot have a rigidity result for arbitrary solutions. However, one may ask if

Date: May 25, 2004.

2000 Mathematics Subject Classification. 49J10, 74G65, 35L70.

Key words and phrases. rigidity, hyperbolic Monge-Ampère equation.

certain rigidity property is restored when we impose proper topology for the convergence of approximate solutions and suitable (e.g. affine) boundary conditions. An interesting model case is to consider a compact subset K of the set

$$\mathcal{H}_{-D} := \{M \in \mathbb{M}_{sym}^{2 \times 2} : \det(M) = -D\} \subset \mathbb{R}^3, \quad D > 0.$$

Using the coordinates $\begin{pmatrix} Z+X & Y \\ Y & Z-X \end{pmatrix}$ on the symmetric 2×2 matrices, we see that \mathcal{H}_{-D} is a hyperboloid of one sheet in \mathbb{R}^3 . It is generated by two families of straight lines which are exactly the rank-one lines on \mathcal{H}_{-D} . Through each point on \mathcal{H}_{-D} there are exactly two rank-one lines. Thus, \mathcal{H}_{-D} locally behaves in many ways like the linear space of diagonal 2×2 matrices, which is well-understood ([4], [17], [24]). However, \mathcal{H}_{-D} globally exhibits much more subtle behavior (see [14] and the discussion below).

On the other hand, the absence of a rank-one connection does not guarantee rigidity, i.e., the absence of microstructures (see [16] for the survey and examples). Especially, the so called T_4 -configuration consisting of four matrices without a rank-one connection, which was observed independently in different contexts ([20], [1], [19], [3], [24]), has been applied to the construction of solutions of elliptic systems with nowhere C^1 (e.g., see [18]). This reveals the difficulty in deriving the regularity theory for general elliptic systems of PDEs. For more relevant issues, one can find the one-well problem in [13], the N -gradient problem for $N \in \{2, 3, 4\}$ in [2], [23], [5], and the geometric rigidity problem in [11]. Note that none of the sets considered in these papers contains a rank-one connection.

In [5] and [14], Chlebík and Kirchheim have analyzed the behavior of Lipschitz maps with $\nabla f \in \mathcal{H}_{-1}$, i.e.,

$$(1.1) \quad \begin{cases} (\nabla f)^T = \nabla f, \\ \det \nabla f = -1. \end{cases}$$

This is equivalent to the hyperbolic Monge-Ampère equation,

$$(1.2) \quad \det \nabla^2 u = -1 \text{ in } \Omega, \quad u \in W^{2,\infty}(\Omega),$$

where $f = \nabla u$. This equation is locally determined by two functions of one variable. In fact, Eq.(1.2) can be transformed into the linear wave equation by a nonlinear change of variables. To see this, suppose at $x_0 \in \Omega$, $\partial_1^2 u(x_0) \neq 0$, $\partial_2^2 u(x_0) \neq 0$ and

hence $\partial_1 \partial_2 u(x_0) \neq \pm 1$ (this can always be achieved by a suitable rotation of coordinates). Then, we will see in the following that we can use $(f_2(x) - x_1, f_2(x) + x_1)$ as independent variables and $(f_1(x) + x_2, f_1(x) - x_2)$ as dependent variables. Since ∇f is symmetric and $\det \nabla f = -1$, the map

$$f^-(x) := f(x) - \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}$$

satisfies

$$\det \nabla f^- = \det \left(\nabla f - \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right) = 0.$$

Thus $f_1(x) + x_2$ is locally a function of $f_2(x) - x_1$, by the implicit function theorem.

Similar reasoning for

$$f^+(x) := f(x) + \begin{pmatrix} -x_2 \\ x_1 \end{pmatrix}$$

shows that there exist locally two functions d_1 and d_2 such that near x_0 ,

$$(1.3) \quad \begin{pmatrix} f_1(x) + x_2 \\ f_1(x) - x_2 \end{pmatrix} = \begin{pmatrix} d_1(f_2(x) - x_1) \\ d_2(f_2(x) + x_1) \end{pmatrix}.$$

To the author's knowledge, this method was first used by Heinz [12] for the elliptic Monge-Ampère equation. In Schulz's book [22], this method has been called the Legendre-like transformation and applied to derive the regularity for the two-dimensional elliptic Monge-Ampère equations. Further references concerning this method include [9], [21] for the elliptic case, and [15] for the hyperbolic case. For the standard Legendre transformation converting a nonlinear PDE into a linear one, readers are referred to Ch.4 of [8]. Kirchheim in [14] gave an example showing that Eq.(1.3) may fail for $W^{2,\infty}$ solutions of Eq.(1.2) or, equivalently, $W^{1,\infty}$ solutions of Eq.(1.1). However, he showed that the set of so called branch points \bar{x} , for which Eq.(1.3) does not hold in a neighborhood of \bar{x} (after suitable rotation of local coordinates), is discrete. Furthermore, for exact Lipschitz solutions of Eq.(1.1) with affine boundary condition, Kirchheim proved that any compact subset $K \subset \mathcal{H}_{-1}$ is rigid.

In this paper, we consider the perturbed equation of Eq.(1.2),

$$(1.4) \quad \det \nabla^2 u = -\varphi^2 \text{ in } W^{2,\infty}(\Omega),$$

where φ^2 means the square of a Lipschitz continuous function φ with $\|\varphi - 1\|_{W^{1,\infty}}$ sufficiently small. We show that a perturbed version of Eq.(1.3) holds locally if the oscillation of $\nabla^2 u$ is not very large (see Theorem 1). By applying the

Legendre-like transformations, we establish the Bilipschitz maps S, T, h between domains $\Omega, \mathcal{U}_z, \mathcal{U}_\xi$ as shown in Figure 1. Since the map h (or $g = h^{-1}$) satisfies the quasilinear hyperbolic 2×2 system, for a fixed point $z^{(0)} \in \mathcal{U}_z$ (or $\xi^{(0)} = h(z^{(0)}) \in \mathcal{U}_\xi$) and $\pm \in \{+, -\}$, there are exactly two characteristic curves, or so called rarefaction curves, $R_\pm(z^{(0)})$ (or $E_\pm(\xi^{(0)})$) passing through $z^{(0)}$ (or $\xi^{(0)}$) with strictly positive and negative finite slopes. Notice that for each $\pm \in \{+, -\}$ we have $S^{-1}(R_\pm(z^{(0)})) = T^{-1}(E_\pm(\xi^{(0)})) \subset \Omega$, which intersects the lines $\{x_1 = c_1\}$ and $\{x_2 = c_2\}$ transversely for some $(c_1, c_2) \in \Omega$. Denote by $(c_1, w_\pm(z^{(0)})) = S^{-1}(R_\pm(z^{(0)})) \cap \{x_1 = c_1\}$ and $(w_\pm(\xi^{(0)}), c_2) = T^{-1}(E_\pm(\xi^{(0)})) \cap \{x_2 = c_2\}$ the points of intersection in Ω . Thus, for each point $x = S^{-1}(z) = T^{-1}(\xi) \in \Omega$, we can locally define the maps $R : \mathcal{U}_z \rightarrow \mathbb{R}^2$ and $E : \mathcal{U}_\xi \rightarrow \mathbb{R}^2$ by $R(z) = (w_+(z), w_-(z))$ and $E(\xi) = (\eta_+(\xi), \eta_-(\xi))$. Since $w_\pm(\tilde{z}) = w_\pm(z)$ (or $\eta_\pm(\tilde{\xi}) = \eta_\pm(\xi)$) whenever $\tilde{z} \in R_\pm(z)$ (or $\tilde{\xi} \in E_\pm(\xi)$), $w_\pm(z)$ (or $\eta_\pm(\xi)$) are called the Riemann invariants for all $\pm \in \{+, -\}$, in order to be consistent with the terminology of hyperbolic systems in PDE theory. We further define the Bilipschitz map $D : R(\mathcal{U}_z) \rightarrow E(\mathcal{U}_\xi)$ locally by $D(w_+(z), w_-(z)) = (\eta_+(\xi), \eta_-(\xi))$. Since $\bigcup_{\pm, z} S^{-1}(R_\pm(z))$ forms a local coordinate net in Ω and for each fixed \pm and z , $S^{-1}(R_\pm(z))$ intersects the straight lines $\{x_1 = c_1\}$ and $\{x_2 = c_2\}$ transversely, the Bilipschitz map D is a diagonal map, i.e., ∇D is a diagonal 2×2 matrix. By applying the Kirszbraun's Lipschitz extension theorem (see Federer [10] 2.10.43), we establish a global argument from the local one stated as above. The estimate in Corollary 1 is a direct consequence of the diagonal map D and the affine boundary condition. For a nonconvex domain Ω , we refine the above argument to derive the geometric rigidity results for some compact subsets of \mathcal{H}_{-1} with respect to certain approximate solutions of Eq.(1.2) in Theorem 2. The rigidity result could be expected for an elliptic equation. Note, however, that Eq.(1.4) is a nonlinear hyperbolic equation. Since the definition of the so called branch points in [14] strongly used the explicit structure of Eq.(1.1), it is not clear how to generalize Kirchheim's arguments to the perturbed case. We thus leave it as a topic to be studied in the near future.

The following theorem gives the main estimates in this paper.

Theorem 1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded and convex Lipschitz domain. Assume $u \in W^{2,\infty}(\Omega)$ is a solution of Eq.(1.4) with φ satisfying*

$$(1.5) \quad \|\varphi - 1\|_{W^{1,\infty}(\Omega)} \leq \delta < \frac{1}{2}.$$

If we further assume

$$(1.6) \quad \frac{\partial^2 u}{\partial^2 x_1} \leq -m^2, \quad \frac{\partial^2 u}{\partial^2 x_2} \geq m^2, \quad 0 < m^2 < \frac{1}{2},$$

where m is a nonzero constant, then for a given small enough $\varepsilon > 0$, there exists a positive constant $\delta_0 = \delta_0(\varepsilon, m, \text{diam}(\Omega))$ such that the following holds:

If $\delta < \delta_0$, then there exist Bilipschitz maps \tilde{E} , \tilde{R} and D such that ∇D is diagonal, i.e.,

$$D \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} = \begin{pmatrix} d_1(w_1) \\ d_2(w_2) \end{pmatrix},$$

$$\|\tilde{E} - id\|_{W^{1,\infty}(\Omega)} < \varepsilon, \quad \|\tilde{R} - id\|_{W^{1,\infty}(\Omega)} < \varepsilon,$$

and

$$\tilde{E} \begin{pmatrix} \frac{\partial u}{\partial x_1} + x_2 \\ \frac{\partial u}{\partial x_1} - x_2 \end{pmatrix} = D \circ \tilde{R} \begin{pmatrix} \frac{\partial u}{\partial x_2} - x_1 \\ \frac{\partial u}{\partial x_2} + x_1 \end{pmatrix}.$$

This result immediately implies a rigidity estimate for compact perturbations of affine data.

Corollary 1. *Suppose all the hypotheses of Theorem 1 hold. If we further assume $\nabla^2 u = A$ in a neighborhood of $\partial\Omega$, where A is a constant matrix with $\det A = -1$, then*

$$\|\nabla^2 u - A\|_{L^\infty(\Omega)} < \delta \cdot C(m, \text{diam}(\Omega)).$$

More generally, we have the following rigidity result:

Theorem 2. *(i) Suppose Ω is a bounded Lipschitz domain in \mathbb{R}^2 , which is not necessarily simply connected, and K is a compact subset of \mathcal{H}_{-1} with sufficiently small diameter $\text{diam}(K)$. Let $u^{(j)}$ be a sequence of functions in $W^{2,\infty}(\bar{\Omega})$ with*

$$\text{dist}(\nabla^2 u^{(j)}, K) \rightarrow 0 \text{ in } W_0^{1,\infty}(\Omega),$$

and on the boundary $\partial\Omega$,

$$(1.7) \quad \nabla u^{(j)}|_{\partial\Omega} = F, \quad \nabla^2 u^{(j)}|_{\partial\Omega} = \nabla F = A,$$

for some affine map F . Then,

$$\nabla^2 u^{(j)} \rightarrow A \text{ in } L^\infty(\Omega).$$

(ii) The restriction on $\text{diam}(K)$ can be dropped if the sequence of functions are in $C^2(\overline{\Omega})$.

If Ω is simply connected, then the assumption in the first part of Eq.(1.7), $\nabla u^{(j)}|_{\partial\Omega} = F$, is redundant since the connectedness of $\partial\Omega$ implies the assumption up to a constant.

2. PROOF OF THE MAIN RESULTS

To simplify our notation, it will be more convenient to establish the equivalent representative

$$E\left(\begin{matrix} f_1(x) \\ f_2(x) \end{matrix}\right) = D \circ R\left(\begin{matrix} x_1 \\ x_2 \end{matrix}\right),$$

with

$$\left\| \nabla E - \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \right\|_{L^\infty} < \varepsilon, \quad \left\| \nabla R - \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix} \right\|_{L^\infty} < \varepsilon.$$

The maps E and R will be constructed by Riemann invariants. The relation between the various maps and domains which appear in the proof is summarized in Figure 1. We remark that the map Σ exists only when f is invertible. We will, however, not use Σ but work directly with g and h .

$$\begin{array}{ccccc} (f_1, f_2) & \xrightarrow{\Sigma} & (x_1, f_2) \in \mathcal{U}_z & \xrightarrow{R} & (w_1, w_2) \in \mathcal{U}_w \\ & & \nearrow^S & & \downarrow D \\ f \uparrow & & g \uparrow \downarrow h & & \\ (x_1, x_2) \in \Omega & \xrightarrow{T} & (f_1, x_2) \in \mathcal{U}_\xi & \xrightarrow{E} & (\eta_1, \eta_2) \in \mathcal{U}_\eta \end{array}$$

Figure 1
Outline of the changes of variables in the proof

Proof of Theorem 1. We first study the geometry of the restrictions, Eq.(1.4) and (1.6), on $\nabla^2 u$. The norm of a symmetric 2×2 matrix $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ is defined as

$$\|A\| := \sqrt{\frac{1}{2} \sum_{i,j=1}^2 a_{ij}^2}.$$

Applying the coordinates $\begin{pmatrix} Z - X & Y \\ Y & Z + X \end{pmatrix}$ on symmetric 2×2 matrices, we have the form: $\|A\| = \sqrt{X^2 + Y^2 + Z^2}$. Assume $Z_0 \geq 0$ is a fixed constant. Then, Eq.(1.6) means that X satisfies both $X \geq m^2 + Z_0$ and $X \geq m^2 - Z_0$. Thus, $X \geq m^2 + Z_0$. By Eq.(1.4),

$$(2.1) \quad 0 \leq Y^2 = Z_0^2 - X^2 + \varphi^2 \leq Z_0^2 + \varphi^2 - (m^2 + Z_0)^2 = \varphi^2 - 2m^2Z_0 - m^4.$$

By Eq.(1.5) and the same calculation in the case: $Z_0 < 0$, we conclude that

$$(2.2) \quad \left| \frac{\partial^2 u}{\partial x_1 \partial x_2} \right| \leq \sqrt{\frac{1}{4} - m^4}.$$

Furthermore, Z_0 satisfies

$$(2.3) \quad |Z_0| \leq \frac{\varphi^2 - m^4}{2m^2}.$$

Now, Eq.(1.5) implies that $\text{dist}(\nabla^2 u, \mathcal{H}_{-1})$ is finite in the sense of L^∞ -norm. Meanwhile, Eq.(2.3) implies $\nabla^2 u$ lies between the two hyperplanes, $\{(X, Y, Z) : Z = \pm c(m)\}$.

By the geometry of the hyperboloid \mathcal{H}_{-1} , we conclude that there exists a positive constant M , depending on m , such that

$$(2.4) \quad \sup_{i,j,x} \left| \frac{\partial^2 u(x)}{\partial x_i \partial x_j} \right| \leq M < \infty,$$

For simplicity, we abbreviate $\frac{\partial^2 u}{\partial x_i^2}$ and $\frac{\partial f_i}{\partial x_i}$ as $\partial_i^2 u$ and $\partial_i f_i$ respectively in the rest of this paper.

Step 1⁰ (Existence of maps g, h and the hyperbolic system for g) Set $f(x) = \nabla u(x)$. Define the map $S : \Omega \rightarrow \mathbb{R}^2$ by

$$S(x_1, x_2) = (x_1, f_2(x)).$$

Then S is Lipschitz with

$$(2.5) \quad \nabla S = \begin{pmatrix} 1 & 0 \\ \partial_1 f_2 & \partial_2 f_2 \end{pmatrix}, \quad \det \nabla S = \partial_2^2 u \geq m^2 > 0.$$

Thus S is open and $\mathcal{U}_z := S(\Omega)$ is open. Since $x_2 \mapsto f_2(x_1, x_2)$ is strictly monotone and Ω is *convex*, the map S is injective and has a Lipschitz inverse $S^{-1} : \mathcal{U}_z \rightarrow \Omega$ with

$$(2.6) \quad (\nabla S^{-1}) \circ S = \frac{1}{\partial_2 f_2} \begin{pmatrix} \partial_2 f_2 & 0 \\ -\partial_1 f_2 & 1 \end{pmatrix}.$$

Similarly the map

$$T(x_1, x_2) = (f_1(x), x_2)$$

is Bilipschitz with

$$\nabla T = \begin{pmatrix} \partial_1 f_1 & \partial_2 f_1 \\ 0 & 1 \end{pmatrix}, \quad (\nabla T^{-1}) \circ T = \frac{1}{\partial_1 f_1} \begin{pmatrix} 1 & -\partial_2 f_1 \\ 0 & \partial_1 f_1 \end{pmatrix},$$

and

$$\det \nabla T = \partial_1^2 u \leq -m^2 < 0.$$

We set $\mathcal{U}_\xi := T(\Omega)$ and define Bilipschitz maps

$$(2.7) \quad g = S \circ T^{-1}, \quad h = T \circ S^{-1} = g^{-1}.$$

Then

$$(2.8) \quad (\nabla g) \circ T = \frac{1}{\partial_1 f_1} \begin{pmatrix} 1 & -\partial_2 f_1 \\ \partial_1 f_2 & \det \nabla f \end{pmatrix}.$$

Keeping in mind that $\partial_1 f_2 = \partial_2 f_1$ and $\det \nabla f = -\varphi^2$, we deduce that for $g(\xi)$,

$$\frac{\partial}{\partial \xi_2} g + \begin{pmatrix} 0 & 1 \\ \varphi^2(z_1, \xi_2) & 0 \end{pmatrix} \cdot \frac{\partial}{\partial \xi_1} g = 0,$$

for $\xi \in T(\Omega)$ and $z = g(\xi)$. The matrix $\begin{pmatrix} 0 & 1 \\ \varphi^2(z_1, \xi_2) & 0 \end{pmatrix}$ has eigenvalues

$$\lambda^\pm = \pm \varphi(z_1, h_2(z)),$$

with right and left eigenvectors

$$(2.9) \quad r^\pm = (1, \pm \varphi(z_1, h_2(z))), \quad l^\pm = (\pm \varphi(z_1, h_2(z)), 1).$$

In the above, the notation \pm means either $+$ or $-$. From now on, denote the opposite sign of \pm as \mp .

Step 2⁰ (Globally defined Riemann invariants and existence of Bilipschitz maps R and E) For a fixed point $z^{(0)} \in \mathcal{U}_z$ and $\pm \in \{+, -\}$, we define the rarefaction curve $R^\pm(z^{(0)})$ locally in \mathcal{U}_z by

$$(2.10) \quad \begin{cases} \frac{d}{dt} \alpha^{\pm, z^{(0)}}(t) = \left(1, \pm \varphi \left(\alpha_1^{\pm, z^{(0)}}(t), h_2 \left(\alpha^{\pm, z^{(0)}}(t) \right) \right) \right), \\ \alpha^{\pm, z^{(0)}}(0) = z^{(0)}. \end{cases}$$

By basic ODE theory, the Lipschitz condition of φ and h implies the existence of the Lipschitz solutions of Eq.(2.10). We may assume $(c_1, c_2) \in \Omega$, for some constants c_1 and c_2 . We may assume that the rarefaction curves $\bigcup_{z^{(0)}, \pm} \{R^\pm(z^{(0)})\}$ parametrize the domain \mathcal{U}_z , because of the local uniqueness of the solution of Eq.(2.10). Since $\varphi \approx 1$, the curve $R^\pm(z^{(0)})$ intersects the line $\{z_1 = c_1\}$ transversely. Define the Riemann invariants $w_+(z^{(0)})$ and $w_-(z^{(0)})$ by

$$w_\pm(z^{(0)}) := \{\widehat{z}_2 : \text{if } (c_1, \widehat{z}_2) \text{ lies on the curves } R^\pm(z^{(0)})\}.$$

By the definition of $w_{\pm}(z^{(0)})$ and Eqs.(2.10), (2.9), we have

$$\nabla_z w_{\pm} \cdot r^{\pm} = 0, \quad \forall \pm \in \{+, -\}.$$

Let $z^{(0)} = (z_1^{(0)}, z_2^{(0)})$ and $\alpha^{\pm, z^{(0)}}(t)$ be the solution of Eq.(2.10). It is easy to verify that

$$(2.11) \quad w_{\pm}(z^{(0)}) = z_2^{(0)} \mp \int_{c_1}^{z_1^{(0)}} \varphi\left(t, h_2\left(\alpha^{\pm, z^{(0)}}(t)\right)\right) dt.$$

From now on, we identify the notation (w_+, w_-) with (w_1, w_2) . Denote by R the map $R(z) = w$ and the set $R(\mathcal{U}_z)$ by \mathcal{U}_w . Notice, that the Riemann invariants w_{\pm} are defined by the rarefaction curves $\alpha^{\pm, z}(t)$, which are defined in \mathcal{U}_z only. However, by Kirszbraun's Lipschitz extension Theorem (see [10] 2.10.43), we can extend the Lipschitz function

$$\psi(z) := \varphi(z_1, h_2(z))$$

from \mathcal{U}_z to \mathbb{R}^2 such that $\bar{\psi}: \mathbb{R}^2 \rightarrow \mathbb{R}$, with

$$(2.12) \quad Lip(\bar{\psi}) = Lip(\psi), \quad \|\bar{\psi} - 1\|_{L^\infty} < \delta \cdot (\text{diam}(\mathcal{U}_z) + 1).$$

Therefore, the rarefaction curves $\alpha^{\pm, z}(t)$ can be globally defined in \mathbb{R}^2 by

$$(2.13) \quad \frac{d}{dt} \alpha^{\pm, z}(t) = (1, \pm \bar{\psi}(\alpha^{\pm, z}(t))).$$

Hence, the Riemann invariants w_{\pm} are well defined on \mathcal{U}_z .

Now, we show that the map R is Bilipschitz. From Eqs.(2.11) and (2.13), for any $z^{(0)}, z^{(1)} \in \mathcal{U}_z$, we have

$$(2.14) \quad \begin{aligned} w^{\pm}(z^{(1)}) - w^{\pm}(z^{(0)}) &= \left(z_2^{(1)} - z_2^{(0)}\right) \mp \left(z_1^{(1)} - z_1^{(0)}\right) \mp \int_{z_1^{(1)}}^{z_1^{(1)}} \left[\bar{\psi}\left(\alpha^{\pm, z^{(1)}}(t)\right) - 1\right] dt \\ &\quad \mp \int_{c_1}^{z_1^{(0)}} \left[\bar{\psi}\left(\alpha^{\pm, z^{(1)}}(t)\right) - \bar{\psi}\left(\alpha^{\pm, z^{(0)}}(t)\right)\right] dt. \end{aligned}$$

By Eq.(1.5),

$$(2.15) \quad \left| \int_{z_1^{(0)}}^{z_1^{(1)}} \left[\bar{\psi}\left(\alpha^{\pm, z^{(1)}}(t)\right) - 1\right] dt \right| < \delta \cdot (\text{diam}(\mathcal{U}_z) + 1) \cdot |z^{(1)} - z^{(0)}|.$$

By Eqs.(2.2), (1.6), (2.5), (2.6), (2.7), (2.8), (2.12), we have

$$(2.16) \quad \text{diam}(\mathcal{U}_z) \leq \sup |\nabla S| \cdot \text{diam}(\Omega) \leq C(m) \cdot \text{diam}(\Omega),$$

$$Lip(\bar{\psi}) = Lip(\psi) \leq Lip(\varphi) \cdot Lip(S^{-1}) \leq Lip(\varphi) \cdot C(m),$$

and

$$(2.17) \quad \frac{1}{|z^{(1)} - z^{(0)}|} \left| \int_{c_1}^{z_1^{(0)}} \left[\bar{\psi} \left(\alpha^{\pm, z^{(1)}}(t) \right) - \bar{\psi} \left(\alpha^{\pm, z^{(0)}}(t) \right) \right] dt \right| \leq \text{diam}(\mathcal{U}_z) \cdot \text{Lip}(\bar{\psi}) \cdot \sup_t \frac{|\alpha^{\pm, z^{(1)}}(t) - \alpha^{\pm, z^{(0)}}(t)|}{|z^{(1)} - z^{(0)}|}.$$

By standard ODE theory, we have

$$(2.18) \quad \lim_{z^{(1)} \rightarrow z^{(0)}} \left(\sup_t \frac{|\alpha^{\pm, z^{(1)}}(t) - \alpha^{\pm, z^{(0)}}(t)|}{|z^{(1)} - z^{(0)}|} \right) \leq \exp(\text{Lip}(\bar{\psi}) \cdot \text{diam}(\mathcal{U}_z)).$$

By the definition of the map R and the estimates in Eqs.(2.14)~(2.18), one can verify that we have the estimate,

$$(2.19) \quad \left\| \nabla R - \begin{pmatrix} -1 & 1 \\ 1 & 1 \end{pmatrix} \right\|_{L^\infty(\overline{\mathcal{U}_z})} < \delta \cdot C_1(m, \text{diam}(\Omega)).$$

If we assume $\varepsilon = \delta \cdot C_1$ is sufficiently small, then R is Bilipschitz. By standard ODE theory, we have the local existence and uniqueness for the solutions of Eq.(2.10). This implies that R is injective. Thus R^{-1} exists and we may treat ξ as the function of w by setting $\xi = h \circ R^{-1}(w)$.

Step 3⁰ (Properties of the diagonal map D between two pairs of Riemann invariants) For fixed $\xi^{(0)} := h(z^{(0)})$ and $\pm \in \{+, -\}$, denote the Lipschitz curve $h \circ \alpha^{\pm, z^{(0)}}(t)$ in \mathcal{U}_ξ as $E_\pm(\xi^{(0)})$. Let

$$\beta^{\pm, \xi^{(0)}}(t) := h \circ \alpha^{\pm, z^{(0)}}(t).$$

Then, by Eq.(2.7), (2.8) and (2.10), one can verify that

$$(2.20) \quad \frac{d}{dt} \beta^{\pm, \xi^{(0)}}(t) = (\nabla h) \left(\alpha^{\pm, z^{(0)}}(t) \right) \cdot \frac{d}{dt} \alpha^{\pm, z^{(0)}}(t) = \frac{\pm \varphi - \partial_1 f_2}{\partial_2 f_2} \begin{pmatrix} \mp \varphi \\ 1 \end{pmatrix}.$$

The assumptions in Eqs.(1.5), (2.2) and (1.6) imply that $\frac{\pm \varphi - \partial_1 f_2}{\partial_2 f_2}$ is bounded, nonzero and stays one-signed. One can verify that the Lipschitz curve $E^\pm(\xi^{(0)})$ can be described by $\gamma^{\pm, \xi^{(0)}}(s)$, which satisfies

$$\begin{cases} \frac{d}{ds} \gamma^{\pm, \xi^{(0)}}(s) = \left(\mp \varphi \left(g_1 \left(\gamma^{\pm, \xi^{(0)}}(s) \right), s \right), 1 \right), \\ \gamma^{\pm, \xi^{(0)}}(0) = \xi^{(0)}. \end{cases}$$

Define the quantity η_\pm (or called the Riemann invariant) on \mathcal{U}_ξ by

$$\eta_\pm(\xi^{(0)}) := \left\{ \widehat{\xi}_1 : \left(\widehat{\xi}_1, c_2 \right) \text{ lies on the curves } E^\pm(\xi^{(0)}) \right\}.$$

It is easy to verify that

$$(2.21) \quad \eta_{\pm}(\xi^{(0)}) = \xi_1^{(0)} \pm \int_{c_2}^{\xi_2^{(0)}} \varphi\left(g_1\left(\gamma^{\pm, \xi^{(0)}}(s)\right), s\right) ds.$$

By a similar argument as in *Step* ϱ^0 , one can verify that the map E is well defined on \mathcal{U}_{ξ} and Bilipschitz with

$$(2.22) \quad \left\| \nabla E - \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \right\|_{L^{\infty}(\overline{\mathcal{U}_{\xi}})} < \delta \cdot C_1(m, \text{diam}(\Omega)).$$

The Bilipschitz map R maps the rarefaction curves α^{\pm} to the lines $\{w_{\pm} = \text{constant}\}$. Similarly, E maps the curves $\beta^{\pm} = h \circ \alpha^{\pm}$ to the lines $\{\eta_{\pm} = \text{constant}\}$. Hence $D := E \circ h \circ R^{-1}$ is Bilipschitz which maps $\{w_1 = \text{constant}\}$ to $\{\eta_1 = \text{constant}\}$ and $\{w_2 = \text{constant}\}$ to $\{\eta_2 = \text{constant}\}$. Therefore, $D(w) = (d_1(w_1), d_2(w_2))$ and $E \circ h = D \circ R$. Recall the definition of h , we find $\eta = D(w)$ on \mathcal{U}_w and

$$(2.23) \quad \nabla D = \begin{pmatrix} \frac{\partial \eta_1}{\partial w_1} & \frac{\partial \eta_1}{\partial w_2} \\ \frac{\partial \eta_2}{\partial w_1} & \frac{\partial \eta_2}{\partial w_2} \end{pmatrix} = \begin{pmatrix} \dot{d}_1(w_1) & 0 \\ 0 & \dot{d}_2(w_2) \end{pmatrix},$$

where \dot{d}_i denotes the derivative of the functions of one-variable d_i .

Let $\tilde{R} := R \circ L_z$ and $\tilde{E} := E \circ L_{\xi}$, where

$$L_z : \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \mapsto \begin{pmatrix} (z_2 - z_1)/2 \\ (z_2 + z_1)/2 \end{pmatrix},$$

$$L_{\xi} : \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} \mapsto \begin{pmatrix} (\xi_1 + \xi_2)/2 \\ (\xi_1 - \xi_2)/2 \end{pmatrix}.$$

Then the proof is finished. \square

Proof of Corollary 1. From the proof in Theorem 1, it is obvious that for each fixed $x^{(0)} \in \Omega$ there exist two points $x^{(+)}, x^{(-)}$ on $\partial\Omega$ such that $S(x^{(\pm)}) \in \alpha^{\pm, S(x^{(0)})}$. Since $\frac{\pm\varphi - \partial_1 f_2}{\partial_2 f_2}$ is bounded and stays one-signed, one can simply verify that the approximate tangent lines of the Lipschitz curve $S^{-1}\left(\alpha^{\pm, S(x^{(0)})}\right)$ are never vertical, horizontal or changing the sign of slope in the chosen coordinates of Ω . From Figure 1 and Theorem 1, one can further verify that

$$(2.24) \quad \left\| \nabla D \circ R \circ S - \begin{pmatrix} \frac{\partial_1 f_2 + 1}{\partial_2 f_2} & 0 \\ 0 & \frac{\partial_1 f_2 - 1}{\partial_2 f_2} \end{pmatrix} \right\|_{L^{\infty}(\Omega)} < \delta \cdot C_2(m, \text{diam}(\Omega)).$$

By Eq. (2.23), (2.24) and triangle inequality, we obtain the estimates

$$\left| \frac{\partial_1 f_2 \pm 1}{\partial_2 f_2}(x^{(0)}) - \frac{\partial_1 f_2 \pm 1}{\partial_2 f_2}(x^{(\mp)}) \right| < \delta \cdot 2C_2(m, \text{diam}(\Omega)),$$

for each $\pm \in \{+, -\}$. The proof is then finished by applying the affine boundary condition. \square

Proof of Theorem 2. (i) For each $f^{(j)}$, we want to apply the argument in the proof of Theorem 1 and Corollary 1. Thus, we need to check if the slightly different assumptions in Theorem 2 would cause problems. When j is sufficiently large and $\text{diam}(K)$ is sufficiently small, the assumptions in Eq.(2.2) and (1.6) are fulfilled by a suitable rotation of coordinates in Ω . Notice that, here, we apply the property: both $\|\nabla^2 u\|$ and $\det \nabla^2 u$ are invariant under rotations of the coordinates of Ω . Due to the nonlinear change of variables associated to $f^{(j)}$, we have the maps S_j and T_j , which denote the maps S and T respectively in Figure 1. However, without assuming the convexity of Ω , the inverse map S_j^{-1} or T_j^{-1} may not exist. This problem can be fixed by the affine boundary condition in Eq.(1.7). Namely, we have the Lipschitz extension of $f^{(j)}$ by defining

$$\bar{f}^{(j)}(x) := \begin{cases} f^{(j)}(x) & , x \in \Omega, \\ F(x) & , x \in \mathbb{R}^2 \sim \Omega. \end{cases}$$

Since $\nabla \bar{f}^{(j)}$ is symmetric and uniformly bounded on \mathbb{R}^2 , there exists an entire function $\bar{u}^{(j)}$ so that $\nabla \bar{u}^{(j)} = \bar{f}^{(j)}$. It is easy to verify that $\bar{u}^{(j)}$ satisfies the assumptions in Eqs.(1.5), (2.2), (1.6), and thus both $\partial_2^2 \bar{u}^{(j)}$ and $\partial_1^2 \bar{u}^{(j)}$ are strictly positive and negative. This guarantees the existence of both the inverse maps of \bar{S}_j and $\bar{T}_j : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, which are maps induced from $\nabla \bar{u}^{(j)}$ as shown in Figure 1. Therefore, both the inverse maps S_j^{-1} and T_j^{-1} exist.

The proof is finished by following the argument in Theorem 1 and Corollary 1.

(ii) Since Ω is bounded, we can choose a domain of rectangle in \mathbb{R}^2 to contain Ω . Namely, for each fixed $k \in \{1, 2\}$, we can define the finite numbers,

$$x_k^{\text{inf}}(\Omega) := \inf_{x \in \Omega} \{x_k : x = (x_1, x_2) \in \Omega\}; \quad x_k^{\text{sup}}(\Omega) := \sup_{x \in \Omega} \{x_k : x = (x_1, x_2) \in \Omega\}.$$

Thus, $\Omega \subset [x_1^{\text{inf}}(\Omega), x_1^{\text{sup}}(\Omega)] \times [x_2^{\text{inf}}(\Omega), x_2^{\text{sup}}(\Omega)]$. Let $t > 0$ and define the domains

$$\Omega_t := \{x = (x_1, x_2) \in \Omega : x_2 > x_2^{\text{sup}}(\Omega) - t\}.$$

By choosing a subsequence, we may suppose as $j \rightarrow \infty$, $\text{dist}(\nabla^2 u^{(j)}, K)$ decreases monotonically in the sense of $W^{1, \infty}(\bar{\Omega})$. For given $\varepsilon > 0$ and j , let

$$V_j^\varepsilon := \left\{ x \in \Omega : \|\nabla^2 u^{(j)} - A\|_{L^\infty(\Omega)} < \varepsilon \right\}.$$

Since $u^{(j)} \in C^2(\bar{\Omega})$, V_j^ε is an open set. Let

$$t_j^\varepsilon := \sup \{t : \Omega_t \subset V_j^\varepsilon\}, \quad \text{and } \mathcal{I}_j^\varepsilon := (0, t_j^\varepsilon) \subset (0, \infty).$$

Now we claim: for given $\varepsilon > 0$, there exists $J^\varepsilon \in \mathbb{N}$ such that

$$\|\nabla^2 u^{(j)} - A\|_{L^\infty(\Omega)} < \varepsilon, \text{ if } j \geq J^\varepsilon.$$

This is equivalent to showing that for fixed $\varepsilon > 0$, $\mathcal{I}_j^\varepsilon = (0, \infty)$ for some sufficiently large j . Since for each j , $u^{(j)} \in C^2(\overline{\Omega})$, thus $\mathcal{I}_j^\varepsilon$ is open and nonempty. Therefore, we only need to show that for some sufficiently large j , $\mathcal{I}_j^\varepsilon$ is also closed in $(0, \infty)$. Assume on the contrary that $\sup_j t_j^\varepsilon \not\leq x_2^{\sup}(\Omega) - x_2^{\inf}(\Omega)$. In other words, for any sufficiently large j , there exists $a^{(j)} = (a_1^{(j)}, a_2^{(j)}) \in \Omega$ with $a_2^{(j)} = x_2^{\sup}(\Omega) - t_j^\varepsilon$ such that

$$(2.25) \quad \|\nabla^2 u^{(j)}(a^{(j)}) - A\|_{L^\infty(\Omega)} \geq \varepsilon.$$

Since the slope of the curve $S^{-1}(\alpha^{\pm, S(a^{(j)})})$ is one-signed for each $\pm \in \{+, -\}$, the curve $S^{-1}(\alpha^{\pm, S(a^{(j)})})$ must intersect $\partial\Omega_{t_j^\varepsilon} \sim \{(x^1, x^2) \in \Omega : x_2 = t_j^\varepsilon\}$, which is a subset of $\partial\Omega$.

Now, applying the argument in Theorem 2 (i) on $\Omega_{t_j^\varepsilon}$, we obtain contradiction from the conclusion of Theorem 2 (i). \square

Acknowledgements. I would like to thank Bernd Kirchheim for introducing the problem, Stefan Müller for revising the manuscript and improving the presentation of the proofs, and the Max Planck Institute for Mathematics in the Sciences, Leipzig/Germany for support during this research. In addition, I appreciate the referee for the comments and suggestions used to revise this paper, and Simon Morgan for his help in revising.

REFERENCES

- [1] R. Aumann and S. Hart, *Bi-convexity and bi-martingales*, Israel J. Math. 54 (1986), no. 2, 159–180.
- [2] J. M. Ball and R. D. James, *Fine phase mixtures as minimizers of energy*, Arch. Rat. Mech. Anal. **100** (1987), 13–52.
- [3] E. Casadio-Tarabusi, *An algebraic characterization of quasi-convex functions*, Ricerche Mat. 42 (1993), no. 1, 11–24.
- [4] N. Chaudhuri and S. Müller, *Rank-one convexity implies quasiconvexity on certain hypersurfaces*, MPI Preprint, 68/2002.
- [5] M. Chlebík and B. Kirchheim, *Rigidity for the four gradient problem*, Journal für die Reine und Angewandte Mathematik **551** (2002), pp.1–9.
- [6] M. Chipot and D. Kinderlehrer, *Equilibrium configurations of crystals*, Arch. Rational Mech. Anal. **103** (1988), 237–277.
- [7] G. Dolzmann, *Variational Methods for Crystalline Microstructure–Analysis and Computation*, Springer Lecture Notes in Mathematics 1803 (2003).

- [8] L. C. Evans, *Partial differential equations*, Graduate Studies in Mathematics, 19. American Mathematical Society, Providence, RI, 1998.
- [9] L. C. Evans and R. F. Gariepy, *On the partial regularity of energy-minimizing, area-preserving maps*, Calc. Var. Partial Differential Equations 9 (1999), no. 4, 357–372.
- [10] H. Federer, *Geometric Measure Theory*, Springer-verlag, New York, 1969.
- [11] G. Friesecke, R. D. James and S. Müller, *A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity*, Comm. Pure Appl. Math. 55 (2002), no. 11, 1461–1506.
- [12] E. Heinz, *Über die Lösungen der Minimalflächengleichung*, Nach. Akad. Wissensch. in Göttingen Math.-Phys. Kl. II (1952), 51-56.
- [13] D. Kinderlehrer, *Remarks about equilibrium configurations of crystals*, in: Material instabilities in continuum mechanics and related mathematical problems (J.M. Ball, ed.), Oxford University Press, 1988, 217-242.
- [14] B. Kirchheim, *Habilitation thesis*, University of Leipzig, 2001.
- [15] O. I. Mokhov and Y. Nutku, *Bianchi transformation between the real hyperbolic Monge-Ampère equation and the Born-Infeld equation*, Lett. Math. Phys. 32 (1994), no. 2, 121–123.
- [16] S. Müller, *Variational models for microstructure and phase transitions*, Calculus of variations and geometric evolution problems (Cetraro, 1996), 85–210, Lecture Notes in Math., 1713, Springer, Berlin, 1999.
- [17] S. Müller, *Rank-one convexity implies quasiconvexity on diagonal matrices*, Internat. Math. Res. Notices 1999, no. 20, 1087–1095.
- [18] S. Müller and V. Šverák, *Convex integration for Lipschitz mappings and counterexamples to regularity*, Ann. of Math. (2) 157 (2003), no. 3, 715–742.
- [19] V. Nesi and G.W. Milton, *Polycrystalline configurations that maximize electrical resistivity*, J. Mech. Phys. Solids 39 (1991), no. 4, 525–542.
- [20] V. Scheffer, *Regularity and irregularity of solutions to nonlinear second order elliptic systems of partial differential equations and inequalities*, Dissertation, Princeton University, 1974.
- [21] R. Schoen and J. Wolfson, *Minimizing volume among Lagrangian submanifolds*, Differential equations: La Pietra 1996 (Florence), 181–199, Proc. Sympos. Pure Math., 65, Amer. Math. Soc., Providence, RI, 1999.
- [22] F. Schulz, *Regularity theory for quasilinear elliptic systems and Monge-Ampère equations in two dimensions*, Lecture Notes in Mathematics, 1445. Springer-Verlag, Berlin, 1990.
- [23] V. Šverák, *On regularity for the Monge-Ampère equations*, Preprint, Heriot-Watt University, 1991.
- [24] L. Tartar, *Some remarks on separately convex functions*, in: *Microstructure and phase transitions*, IMA Vol. Math. Appl. 54 (D. Kinderlehrer, R. D. James, M. Luskin and J.L. Ericksen, eds.), Springer, 1993, 191-204.

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTR. 22-26, LEIPZIG D-04103, GERMANY

Current address: Department of Mathematics, National Taiwan Normal University, Taipei 116, Taiwan

E-mail address: chunlin@math.ntnu.edu.tw