

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

Geometric Rigidity of Conformal
Matrices

by

Daniel Faraco and Xiao Zhong

Preprint no.: 33

2004



Geometric Rigidity of Conformal Matrices

Daniel Faraco* Xiao Zhong†

Abstract

We provide a geometric rigidity estimate à la Friesecke-James-Müller for conformal matrices. Namely, we replace $SO(n)$ by an arbitrary compact subset of conformal matrices, bounded away from 0 and invariant under $SO(n)$, and rigid motions by Möbius transformations.

1 Introduction

This paper is concerned with the so-called geometric rigidity estimates for conformal matrices. Recently, Friesecke, James and Müller developed a successful new approach to the classical problem of dimension reduction in nonlinear elasticity [7]. A fundamental ingredient was the following rigidity estimate for the group of special orthogonal matrices of \mathbb{R}^n , $SO(n) = \{A \in M^{n \times n} : A^t A = I, \det A = 1\}$.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain and $n \geq 2$. There exists a constant $C_1 = C_1(\Omega)$ with the property that for each $v \in W^{1,2}(\Omega, \mathbb{R}^n)$, there exists $R \in SO(n)$ such that*

$$\|Dv - R\|_{L^2(\Omega)} \leq C_1 \|\text{dist}_{SO(n)}(Dv)\|_{L^2(\Omega)}. \quad (1.1)$$

Theorem 1.1 has been used after in a number of related problems concerning dimension reduction, e.g [4], [9], [16] and [17]. In all the applications it is crucial that the dependence between the left and right hand side is linear and that the v is any general Sobolev mapping (the classical result of John [13] gives an $L^2 - L^\infty$ estimate valid for locally Bi-Lipschitz maps). Theorem 1.1 makes quantitative the following classical result of Reshetnyak [18] for sequences.

*Max-Planck-Institute for Mathematics in the Sciences, Inselstr. 22-26, 04103 Leipzig, Germany. faraco@mis.mpg.de

†Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), FIN-40014 University of Jyväskylä, Finland. zhong@maths.jyu.fi

Theorem 1.2. *Let $\{v_j\} \in W^{1,2}(\Omega)$ be a weakly convergent sequence in $W^{1,2}$. Then there exists $R \in SO(n)$ such that*

$$\lim_{j \rightarrow \infty} \|\text{dist}_{SO(n)}(Dv_j)\|_{L^2(\Omega)} = 0 \Rightarrow \lim_{j \rightarrow \infty} \|Dv_j - R\|_{L^2(\Omega)} = 0. \quad (1.2)$$

A natural question raised by Theorem 1.1 is to determine when a qualitative rigidity theorem like Theorem 1.2 can be in fact made quantitative in the sense of Theorem 1.1. That is, let $E \subset M^{n \times n}$ be such that for a weakly convergent sequence $\{v_j\} \in W^{1,2}(\Omega)$ it holds that

$$\lim_{j \rightarrow \infty} \|\text{dist}_E(Dv_j)\|_{L^2(\Omega)} = 0 \Rightarrow \lim_{j \rightarrow \infty} \|Dv_j - D\varphi\|_{L^2(\Omega)} = 0 \quad (1.3)$$

where $\varphi \in W^{1,2}(\Omega, \mathbb{R}^n)$ is such that $D\varphi \in E$ a.e.

Then, does there exist a constant $C(\Omega)$ such that for every $v \in W^{1,2}(\Omega, \mathbb{R}^n)$ there exists φ with $D\varphi \in E$ and

$$\|Dv - D\varphi\|_{L^2(\Omega)} \leq C(\Omega) \|\text{dist}_E(Dv)\|_{L^2(\Omega)}? \quad (1.4)$$

Canonical examples of set satisfying (1.3) are compact subsets of the n dimensional conformal matrices

$$CO_+(n) = \{A \in M^{n \times n} : A = \rho R, \text{ where } \rho \in \mathbb{R}_+ \text{ and } R \in SO(n)\}.$$

In this work, we show that if E is a compact subset of $CO_+(n)$, invariant under $SO(n)$, and with $0 \notin E$ a quantitative rigidity estimate holds (see Theorem 1.4). Before stating the result, we need to recall that for $E \subseteq CO_+(n)$, the solutions to the differential inclusion

$$D\varphi \in E, \quad \varphi \in W^{1,2}(\Omega) \quad (1.5)$$

are described by the Liouville Theorem.

Theorem 1.3 (Liouville Theorem). *Let $\Omega \subset \mathbb{R}^n, n \geq 3$ and let $\varphi \in W^{1,n}(\Omega, \mathbb{R}^n)$ such that*

$$D\varphi(x) \in CO_+(n) \text{ a.e. } x \in \Omega. \quad (1.6)$$

Then,

$$\varphi(x) = b + Ax, \text{ or } \varphi(x) = b + AR \frac{x - a}{|x - a|^2} \quad (1.7)$$

where $b \in \mathbb{R}^n, a \in \mathbb{R}^n \setminus \Omega, A \in CO_+(n)$ and $R = \text{diag}(1, \dots, -1)$.

The Liouville Theorem has a long history. Liouville established it for C^3 mappings in [15], Gehring for homeomorphisms in $W^{1,n}(\Omega, \mathbb{R}^n)$ in [5] and Reshetnyak removed the injectivity assumption in [18]. In fact, $W^{1,n}$ is not the borderline case. Iwaniec [11] proved that there exists a critical threshold $p_n < n$ such that Liouville theorem holds for mappings in $W^{1,p}(\Omega, \mathbb{R}^n)$, $p \geq p_n$. Moreover, Iwaniec and Martin [12] proved that if $n = 2m$ and m an integer then $p_n = m$ and the result is optimal. For n odd p_n is conjectured to be also $\frac{n}{2}$.

Geometrically, the Liouville theorem relates $CO_+(n)$ with the special Möbius group \mathcal{M}_n . The Möbius group $\mathbf{Möb}(n)$ is the group generated by reflections on spheres and hyperplanes. Then \mathcal{M}_n consists of Möbius transformations preserving orientation (see [1] for an introduction of the geometry of the Möbius group and its discrete subgroups). It turns out that any Möbius transform can be represented as the composition of an affine mapping and an inversion respect to a sphere. Analytically, this yields the formula (1.7).

We are interested in compact subsets of $CO_+(n)$ invariant under $SO(n)$ and finitely connected. Let us introduce the notation

$$\begin{aligned} E^{m,M} &= (CO_+(n) \cap B(0, M)) \setminus B(0, m), \\ mSO(n) &= \{mR : R \in SO(n)\}. \end{aligned} \quad (1.8)$$

Then our conditions on E imply that it can be represented by,

$$E = \cup_{i=1}^{n_1} E^{m_i, M_i} \cup_{i=1}^{n_2} m_i SO(n). \quad (1.9)$$

In addition, since the sets E are compact the solutions to (1.5) are in particular Möbius transforms. We denote them by $\mathcal{M}_n^E(\Omega)$. An interesting new feature respect to the other nonlinear sets for which quantitative rigidity estimates are available (e.g [2]) is that $\mathcal{M}_n^E(\Omega)$ contains non affine solutions. We are now in the position to state the rigidity estimate:

Theorem 1.4. *Let $E \subset CO_+(n)$ be compact, finitely connected, with $0 \notin E$ and such that*

$$SO(n)E = E.$$

Let $\Omega' \Subset \Omega \subset \mathbb{R}^n$, $n \geq 3$ and Ω be a bounded domain. Then,

- i) There exists a constant $C_2 = C_2(E, \Omega', \Omega)$ such that for any $v \in W^{1,2}(\Omega, \mathbb{R}^n)$ there exists $\varphi \in \mathcal{M}_n^E(\Omega', \mathbb{R}^n)$ such that*

$$\int_{\Omega'} |D\varphi - Dv|^2 \leq C_2 \int_{\Omega} \text{dist}_E^2(Dv). \quad (1.10)$$

ii) Let Ω be Lipschitz. Then, a constant $C_3 = C_3(E, \Omega)$ such that

$$\int_{\Omega} |D\varphi - Dv|^2 \leq C_3 \int_{\Omega} \text{dist}_E^2(Dv)$$

exists if and only if $E = \cup_{i=1}^{n^2} m_i SO(n)$.

Our approach to Theorem 1.4 is essentially a combination of ideas of [7] with those developed by Reshetnyak in his study of the stability of Liouville theorem respect to different parameters and classes of functions (See [20], in particular Theorems 3.2 and 5.2). The paper is organized as follows. In Section 2 we state some basic facts about Möbius mappings and solutions to elliptic equations. Section 3 is devoted to prove Theorem 1.4 in the special case where $v = \varphi \in \mathcal{M}_n$. We provide an example showing that, if $\mathcal{M}_n^E(\Omega)$ contains non affine mappings, a global estimate like in Theorem 1.4 (ii) does not hold even in this simplified setting.

Section 4 constitutes the crux of the paper. We prove Theorem 1.4 for Ω' a ball. The proof works because E is related to elliptic equations globally and locally. Globally, there exists a smooth uniformly convex mapping $F : \mathbb{R}^n \rightarrow \mathbb{R}$ such that for every $A \in E$

$$F(A_i) = \det(A), \tag{1.11}$$

where A_i is any row of the matrix A . The existence of such a mapping F enable us to write

$$v = w + z,$$

where each of the coordinates z_i satisfies the equation

$$\text{div}(DF(Dz_i)) = 0,$$

and $\int_{\Omega} |\nabla w|^2 \leq C \int_{\Omega} \text{dist}_E^2(Dv)$. Hence, it suffices to prove Theorem 1.4 for such a mapping z . Next, the regularity theory of elliptic equations implies that z_i enjoys a priori estimates. In particular, the modulus of continuity of Dz is uniformly bounded. This allows the use of a compactness argument to deduce that it is essentially enough to prove Theorem 1.4 for mappings z such that

$$\|\text{dist}(Dz, I)\|_{L^\infty(\Omega')} \ll 1 \tag{1.12}$$

and the modulus of continuity of Dz is uniformly bounded. In this situation we can use the local equation which is given by the tangent plane to $CO_+(n)$. We proceed by adapting the ideas of Reshetnyak [20] to our situation. Beside a Korn type inequality for the tangent plane to $CO_+(n)$, a degree argument

involving the exponential map of \mathcal{M}_n is needed to choose the Möbius mapping closest to z . In this way, one obtains a mapping $\varphi' \in \mathcal{M}_n(\Omega')$ satisfying (1.10). However, these arguments do not imply that $\varphi' \in \mathcal{M}_n^E(\Omega')$. The desired mapping φ is obtained by applying section 3 to φ' . Let us remark that this is the only moment along the whole proof where the $SO(n)$ invariance of E is used. The last section is devoted to prove Theorem 1.4 for an arbitrary $\Omega' \Subset \Omega$.

2 Notation and preliminaries

We will denote by C_1, C_2, \dots, C_n constants which will be used during the whole paper, whereas c_1, c_2, \dots, c_n will be used for different constants within the same proof.

Concerning sets in \mathbb{R}^n , we will use B for balls and $B(a, r)$ where we want to specify the center a and the radius r . For balls centered at the origin we use $B_r = B(0, r)$. Given a ball $B(a, r)$, $hB = B(a, hr)$. For a measurable set L , $|L|$ denotes its Lebesgue measure. Let $K \subset \mathbb{R}^n$ and $s \in \mathbb{R}$. Then $K_s = \{y \in \mathbb{R}^n : \text{dist}(y, K) \leq s\}$.

Let $A = (a_{ij}) \in M^{n \times n}$ then $|A|$ stands for the operator norm. Given a closed set $E \in M^{n \times n}$ $\text{dist}(A) = \inf_{B \in E} |A - B|$. Let us remark the choice of the operator norm in our definition of the distance is motivated to simplify the constants in section 3 but of course any other norm would do.

Let E be the set in Theorem 1.4. Since $0 \notin E$ and E is compact there exists $0 < m < M < \infty$ such that

$$E \subset E^{m, M} \tag{2.1}$$

where $E^{m, M}$ was introduced in (1.9).

We will use the notation E' for an auxiliary set different E but satisfying the assumptions of Theorem 1.4. In particular it will satisfy (2.1) for some other numbers m' and M' .

Given a closed set $\Omega \subset \mathbb{R}^n$, $\mathcal{M}_n(\Omega)$ are Möbius transforms which are finite in Ω , $\mathcal{M}_n(\Omega_1, \Omega_2)$ stands for Möbius transform mapping Ω_1 onto Ω_2 . Let us recall the notation

$$\mathcal{M}_n^E(\Omega) = \{\varphi \in W^{1,2}(\Omega) : D\varphi(x) \in E \text{ a. e. } x \in \Omega\}.$$

We discuss now several basic properties of $\mathcal{M}_n(\Omega)$ and more precisely of $\mathcal{M}_n^E(B)$. In particular we show in the first lemma that mappings $\varphi \in \mathcal{M}_n^E(B)$ can be handled in an uniform way. The main reason is that for $\varphi \in \mathcal{M}_n^E(B)$ $\varphi^{-1}(\infty)$, the center of the sphere associated to φ , is bounded away from B independently of φ . For the notation, recall that for $\varphi \in \mathcal{M}_n$, $\varphi(B)$ is a ball.

Lemma 2.1. *Let $\varphi \in \mathcal{M}_n^E(B)$ and let m, M be the constants in (2.1). Then the following three properties hold:*

1. $m^n \leq \frac{|\varphi(B)|}{|B|} \leq M^n$.
2. $\varphi \in \mathcal{M}_n(h_0 B)$ where $h_0 = h_0(\frac{M}{m}) > 1$.
3. Let $s < 1$. Then there exists a number $h_1 = h_1(s, E) < 1$ such that $\varphi(sB) \subset h_1\varphi(B)$ and $s\varphi(B) \subset \varphi(h_1B)$.

Proof: Since for $A \in CO_+(n)$, $|A|^n = \det(A)$, it follows that

$$|\varphi(B)| = \int_B J_\varphi = \int_B |D\varphi|^n$$

(1) follows from (2.1).

We prove (2) for $\varphi \in \mathcal{M}_n^E(B)$ not affine. It follows from (1.7) that for such φ there exists $r \in \mathbb{R}$ and $a \in \mathbb{R}^n$ such that $|D\varphi(x)| = r^2|x-a|^{-2}$. Thus, $\max_B |D\varphi| = r^2 \text{dist}_B(a)^{-2}$ and $\min_B |D\varphi| = r^2(\text{dist}_B(a) + \text{diam}(B))^{-2}$. Since $\varphi \in \mathcal{M}_n^E(B)$ we have that

$$\frac{r^2 \text{dist}_B(a)^{-2}}{r^2(\text{dist}_B(a) + \text{diam}(B))^{-2}} \leq \frac{M}{m},$$

i.e

$$\text{dist}_B(a) \geq \frac{\text{diam}(B)}{\sqrt{\frac{M}{m}} - 1}.$$

Hence $h_0 = 1 + 2\frac{1}{\sqrt{\frac{M}{m}} - 1}$.

For (3), we firstly observe there is no loss of generality in assuming that $B = B_1$ and $\varphi(B_1) = B_1$. The general case follows by considering similarities $T_B, T_{\varphi(B)}$ such that $T_B(B_1) = B$ and $T_{\varphi(B)}(B_1) = \varphi(B)$. Then the mapping $\tilde{\varphi} = T_{\varphi(B)}^{-1} \circ \varphi \circ T_B$ satisfies that $\tilde{\varphi}(B_1) = B_1$ and $m^2 \leq |D\tilde{\varphi}| \leq M^2$. It is easy to check that if the thesis holds for $\tilde{\varphi}$ it also holds for φ .

Therefore there is no loss of generality assuming $\varphi \in \mathcal{M}_n^E(B_1, B_1)$. Since $\frac{1}{M} \leq |D\varphi^{-1}| \leq \frac{1}{m}$, (2) implies that $\varphi^{-1} \in \mathcal{M}_n(h_0 B_1)$ for the same h_0 . Thus, $\varphi(\infty) = b$, with $|b| \geq h_0$.

On the other hand, $\varphi(B_1) = B_1$ implies that φ fixes S^{n-1} . Thus, it conjugates with the mapping $R = \frac{x}{|x|^2}$ (see [1, Theorem 3.2.4]). i.e $\varphi(x) = R \circ \varphi \circ R(x)$. Putting $x = 0$ yields $\varphi(0) = R \circ \varphi \circ R(0) = R(b) = \frac{b}{|b|^2}$. Recall

now that $\mathcal{M}_n(B_1, B_1)$ is the group of isometries of (B_1, d) , where d denotes the hyperbolic metric of B_1 (see [1, chapter 3]). Thus, by triangle inequality,

$$d(\varphi(x), 0) \leq d(\varphi(x), \varphi(0)) + d(\varphi(0), 0) = d(x, 0) + d\left(\frac{b}{|b|^2}, 0\right). \quad (2.2)$$

The hyperbolic distance is related with the Euclidean distance by the formula $d(x, 0) = \log\left(\frac{1+|x|}{1-|x|}\right)$. Therefore we deduce from (2.2) that if $x \in sB_1$ and $\varphi \in \mathcal{M}_n^E(B_1, B_1)$, $|\varphi(x)|$ satisfies the inequality

$$\frac{|\varphi(x)| + 1}{|\varphi(x)| - 1} \leq \frac{(s+1)(h_0+1)}{(s-1)(h_0-1)}.$$

Hence h_1 in the claim (3) of the Lemma is implicitly defined by

$$\frac{h_1 + 1}{h_1 - 1} \leq \frac{(s+1)(h_0+1)}{(s-1)(h_0-1)}.$$

The assertion $s\varphi(B) \subset \varphi(h_1B)$ is equivalent to $\varphi^{-1}(sB) \subset h_1B$ and hence it follows from the above reasoning. \square

The following proposition relies on the fact that \mathcal{M}_n is a finite dimensional manifold and $\mathcal{M}_n(hB)$ is a compact manifold. Thus all metrics are equivalent.

Proposition 2.2 ([20, Chapter 4. Lemma 2.5]). *Let $B = B(a, r)$ and $\varphi, \psi \in \mathcal{M}_n(hB)$ with $h > 1$. Then there exists a constant $C_4 = C_4(h)$ such that for any $x, y \in B$ we have that*

$$|D\varphi(x) - D\psi(x)| \leq \frac{C_4}{|B|} \int_B |D\varphi - D\psi| \quad (2.3)$$

and,

$$|D\varphi(x) - D\varphi(y)| \leq \frac{1}{r} C_4 \max_B |D\varphi| |x - y|. \quad (2.4)$$

The next lemma states that derivatives of Möbius transforms are like constants in the following sense: If they are sufficiently close in a ball, they are also close in a bigger ball.

Lemma 2.3 ([20, Chapter 4. Lemma 4.1]). *Let $\varphi \in \mathcal{M}_n$ such that the inequality*

$$|\varphi(x) - x| \leq r\epsilon$$

holds for all $x \in B(a, r)$, where $\epsilon > 0$. Let $h > 0$. Then there exist constants $\alpha = \alpha(h)$ and $C_5 = C_5(h)$ such that if $\epsilon < \alpha$ then the inequality

$$|D\varphi(x) - I| \leq C_5\epsilon$$

holds for all $x \in hB$.

We conclude the section by introducing the elliptic equations needed in Section 4 and DeGiorgi-Nash's Theorem on the regularity of the solutions.

Definition 2.4. Let $F : \mathbb{R}^n \rightarrow \mathbb{R}$ be a convex function. A mapping $z = (z_1, \dots, z_n) : \Omega \rightarrow \mathbb{R}^n$ is said to be F -harmonic if each of the coordinates z_i satisfies that

$$\operatorname{div}(DF(Dz_i)) = 0 \text{ in } \Omega.$$

Equivalently, z_i minimizes $\int_{\Omega} F(Dv)$ respect to its own boundary values.

Proposition 2.5 ([3],[6]). Let z be an F -harmonic mapping in B . Let $F : \mathbb{R}^n \rightarrow \mathbb{R}$ be a C^∞ uniformly convex and such that $|DF(A)| \leq C_F|A|$, $|D^2F| \leq C_F$ for $C_F > 0$. Let $0 \leq h < 1$. Then, $z \in C^\infty(hB)$ and there exists a number $0 < \alpha < 1$, $\alpha = \alpha(h, F)$ and a constant $C_6 = C_6(h, F)$ such that

$$[Dz]_{C^\alpha(hB)} \leq C_6 \left(\int_B |Dz|^2 \right)^{\frac{1}{2}}.$$

3 \mathcal{M}_n^E as a subset of \mathcal{M}_n

In this section we prove Theorem 1.4 for $v \in \mathcal{M}_n(\Omega) \setminus \mathcal{M}_n^E(\Omega')$. Firstly, we reduce the situation to the case where E is connected in Proposition 3.1. The essential point in the proof is Proposition 2.2 stating that if $\varphi \in \mathcal{M}_n^{E'}$ then $D\varphi$ is Lipschitz and the Lipschitz constant depends only on E' . Thus, there is no loss of generality assuming $E = E^{m.M}$. We treat this case in Proposition 3.2. The essential observation for the proof of Proposition 3.2 is that if $\varphi \notin \mathcal{M}_n^E(\Omega')$, then there exists a ball $\tilde{B} \subset \Omega \setminus \Omega'$ with $|\tilde{B}|^{\frac{1}{n}} \approx \operatorname{dist}(\Omega', \partial\Omega)$ such that for $x \in \tilde{B}$, $D\varphi(x) \notin E$. This, and that for $\varphi \in \mathcal{M}_n^E$ the oscillation of $|D\varphi|$ is uniformly controlled, provides us a constant $C = C(\Omega', E)$ such that

$$\min \left\{ \int_{\Omega'} \operatorname{dist}_{\operatorname{MSO}(n)}^2(D\varphi), \int_{\Omega'} \operatorname{dist}_{\operatorname{MSO}(n)}^2(D\varphi) \right\} \leq C \int_{\tilde{B}} \operatorname{dist}_E^2(D\varphi).$$

Therefore, we can conclude by means of Theorem 1.1.

If one tries to follow a similar scheme for proving an estimate up to the boundary one faces the situation of the Example 3.3 presented at the end of the section.

Proposition 3.1. *Let E, Ω', Ω as in Theorem 1.4 and let $\{E_i\}_{i=1}^I$ be the connected components of E . Then, there exists $C_7 = C_7(E, \Omega', \Omega)$ such that for every $\varphi \in \mathcal{M}_n(\Omega)$*

$$\min_i \left\{ \int_{\Omega'} \text{dist}_{E_i}^2(D\varphi) \right\} \leq C_7 \int_{\Omega} \text{dist}_{\cup_i E_i}^2(D\varphi). \quad (3.1)$$

Proof: By induction, it is enough to prove the theorem for E having two connected components $E = E_1 \cup E_2$. Let $m_1 = \min_{E_1} |A| \leq \max_{E_1} |A| = m_2 < \min_{E_2} |A| = M_1 \leq \max_{E_2} |A| = M_2$ and $\rho = M_1 - m_2 = \text{dist}(E_1, E_2)$ (Here we use the invariance of E under $\text{SO}(n)$). We firstly observe that there exist constants $c_1 = c_1(E)$, $c_2 = c_2(E)$ such that if either $|A| \geq 2M_2$ or $|A| \leq \frac{1}{2m_1}$ then

$$\text{dist}_{E_i}(A) \leq c_1 \text{dist}_E(A), \quad (3.2)$$

and $\text{dist}_E(A) \geq c_2$.

Let $\Omega^E = \{x \in \Omega' : \frac{1}{2m_1} \leq |D\varphi(x)| \leq 2M_2\}$. Suppose that $|\Omega^E| \leq \frac{1}{2}|\Omega|$. Then we have that $2|\Omega \setminus \Omega^E| \geq |\Omega|$ and therefore

$$\begin{aligned} \int_{\Omega} \text{dist}_{E_1}^2(D\varphi) &\leq c_1 \int_{\Omega \setminus \Omega^E} \text{dist}_E^2(D\varphi) dx + 4M_2^2 |\Omega| \\ &\leq c_1 \int_{\Omega \setminus \Omega^E} \text{dist}_E^2(D\varphi) dx + \frac{8M_2^2}{c_2} c_2 |\Omega \setminus \Omega^E| \\ &\leq c_3 \int_{\Omega \setminus \Omega^E} \text{dist}_E^2(D\varphi), \end{aligned}$$

and (3.1) holds. Hence, we can assume that $|\Omega^E| \geq \frac{1}{2}|\Omega|$.

Now, if

$$\text{dist}_{E_1}(D\varphi(x)) \geq \frac{\rho}{2},$$

for every $x \in \Omega_E$ it would follow that

$$\text{dist}_{E_2}(D\varphi(x)) \leq c_4 \text{dist}_E(D\varphi),$$

with $c_4 = \frac{4}{\rho} M_2$. Together with (3.2) we would have that

$$\text{dist}_{E_2}(D\varphi(x)) \leq \max\{c_1, c_4\} \text{dist}_E(D\varphi)$$

for all $x \in \Omega$ and (3.1) would be trivial.

Since the same argument works if we exchange E_1 and E_2 we are left to the case where there exist two points $x_1, x_2 \in \Omega^E$ such that

$$\text{dist}_E(D\varphi(x_1)) = \text{dist}_{E_1}(D\varphi(x_1)) \leq \frac{\rho}{2},$$

$$\text{dist}_E(D\varphi(x_2)) = \text{dist}_{E_2}(D\varphi(x_2)) \leq \frac{\rho}{2}.$$

Then, since Ω is connected we can assume Ω' to be connected. Thus, there exists $x_0 \in \Omega'$ with $|D\varphi(x_0)| = \frac{M+m}{2}$ and consequently $\text{dist}_{E_i}(D\varphi(x_0)) = \frac{\rho}{2}$ for $i = 1, 2$. Since $\Omega' \Subset \Omega$, there is no loss of generality assuming the existence of $r_1 = r_1(\Omega', \Omega)$ such that $B(x_0, r_1) \subset \Omega'$. Furthermore, since $2|\Omega^E| \geq |\Omega|$ and Ω is bounded there exists $r_2 = r_2(\Omega, n)$ such that $B(x_0, r_2) \subset \Omega^E$.

Now it follows from Proposition 2.2 that $D\varphi|_{B(x_0, r_2)}$ is Lipschitz with a constant $L(E, r_2)$. Thus, there is new $r_3 = r_3(\Omega, \Omega', E)$ such that on $B(x_0, r_3)$, $\text{dist}_E(D\varphi(x)) \geq \frac{\rho}{4}$. But now this implies that

$$\begin{aligned} \int_{\Omega} \text{dist}_E^2(D\varphi) &= \int_{\Omega \setminus \Omega^E} \text{dist}_E^2(D\varphi) + \int_{\Omega^E} \text{dist}_E^2(D\varphi) \\ &\geq \int_{\Omega \setminus \Omega^E} \text{dist}_E^2(D\varphi) + \int_{B(x_0, r_3)} \text{dist}_E^2(D\varphi) \\ &\geq \int_{\Omega \setminus \Omega^E} \text{dist}_E^2(D\varphi) dx + c_5 \rho^2 r_3^n \geq c_6 \int_{\Omega'} \text{dist}_{E_1}^2(D\varphi), \end{aligned}$$

where $c_6 = \max\{\frac{c_5 \rho^2 r_3^n}{M^2 |\Omega'|}, c_1^2\}$. The proof is concluded. \square

Proposition 3.2. *Let Ω', Ω and E as in Theorem 1.4. Let $E \subset E'$. Then, there exists a constant $C_8 = C_8(\Omega, \Omega', E')$ such that for every $\varphi \in \mathcal{M}_n^{E'}(\Omega)$ there exists $\varphi_{\Omega'} \in \mathcal{M}_n^E(\Omega')$ satisfying*

$$\int_{\Omega'} |D\varphi - D\varphi_{\Omega'}|^2 \leq C_8 \int_{\Omega} \text{dist}_E^2(D\varphi). \quad (3.3)$$

Proof: By Theorem 1.1 and Proposition 3.1 it suffices to prove the thesis for

$$E = E^{m, M}$$

$0 < m < M < \infty$.

Let $\varphi \in \mathcal{M}_n^{E'}(\Omega)$ and suppose that $\varphi|_{\Omega'} \notin \mathcal{M}_n^E(\Omega')$. Then either $\max_{\Omega'} |D\varphi| > M$ or $\min_{\Omega'} |D\varphi| < m$. We consider the two possibilities separately.

Case $\max_{\Omega'} |D\varphi| > M$.

Let $d = \text{dist}(\Omega', \partial\Omega)$. We will deduce (3.3) from the next two claims:

There exists $c_1 = c_1(\Omega, \Omega', E)$ such that

$$(\max_{\Omega'_{\frac{d}{2}}} |D\varphi| - M)^2 \leq c_1 \int_{\Omega} \text{dist}_E^2(D\varphi). \quad (3.4)$$

There exists $c_2 = c_2(\Omega, \Omega')$ such that if $\min_{\Omega'} |D\varphi| < M$ we have that

$$(\max_{\Omega'_{\frac{d}{2}}} |D\varphi| - M) \geq c_2(M - \min_{\Omega'} |D\varphi|). \quad (3.5)$$

Let us supposed we have proved (3.4) and (3.5) and deduce the thesis from them. Indeed, (3.4) and (3.5) imply that if $x \in \Omega'$ and $|D\varphi(x)| \leq M$, then it holds that

$$\text{dist}_{MSO(n)}^2(D\varphi(x)) \leq c_3 \int_{\Omega} \text{dist}_E^2(D\varphi),$$

with $c_3 = \frac{c_1}{c_2^2}$. On the other hand if $|D\varphi(x)| \geq M$, then $\text{dist}_{MSO(n)}^2(D\varphi(x)) = \text{dist}_E^2(D\varphi(x))$. Hence,

$$\int_{\Omega'} \text{dist}_{MSO(n)}^2(D\varphi) \leq c_3 \int_{\Omega} \text{dist}_E^2(D\varphi) \quad (3.6)$$

and thus (3.3) would follow from Theorem 1.1 with $\varphi_{\Omega'} = Ax$ where $A \in MSO(n)$.

To prove the claims (3.4) and (3.5) we need further notation. Since $\varphi \in \mathcal{M}_n^{E'}(\Omega)$ we have that $|D\varphi(x)| = r^2|x - a|^{-2}$ where $r \in \mathbb{R}$ and $a \in \mathbb{R}^n \setminus \Omega$. Let $x_0 \in \Omega'$ be such that $|D\varphi(x_0)| = \max_{\Omega'} |D\varphi| > M$ and let $L(t) : \mathbb{R} \rightarrow \mathbb{R}^n$ be defined by $L(t) = x_0 + t \frac{a-x_0}{|a-x_0|}$. By triangle inequality if $x \in B(L(\frac{d}{4}), \frac{d}{4})$ then $|D\varphi(x)| \geq M$. Thus for such an x

$$\text{dist}_E(D\varphi(x)) = |D\varphi(x)| - M = \text{dist}_{MSO(n)}(D\varphi(x)). \quad (3.7)$$

Therefore we can apply Theorem 1.1 to φ in $B(L(\frac{d}{4}), \frac{d}{4})$. Together with (3.7) it yields an $R_\varphi \in SO(n)$ such that

$$\int_{B(L(\frac{d}{4}), \frac{d}{4})} |D\varphi - MR_\varphi|^2 \leq c_4 \int_{B(L(\frac{d}{4}), \frac{d}{4})} \text{dist}_E^2(D\varphi). \quad (3.8)$$

Since $\varphi \in \mathcal{M}_n(\Omega)$, $\varphi \in \mathcal{M}_n(B(L(\frac{d}{4}), 4\frac{d}{4}))$. Thus, we can apply Proposition 2.2 to obtain

$$\|D\varphi - MR_\varphi\|_{L^\infty(B(L(\frac{d}{4}), \frac{d}{4}))} \leq \frac{C_4(4)}{|B(L(\frac{d}{4}), \frac{d}{4})|} \int_{B(L(\frac{d}{4}), \frac{d}{4})} |D\varphi - MR_\varphi|. \quad (3.9)$$

Combining (3.8), (3.9) and triangle inequality yields

$$(|D\varphi(L(\frac{d}{2}))| - |MR_\varphi|)^2 \leq c_1 \int_{\Omega} \text{dist}_E^2(D\varphi), \quad (3.10)$$

with $c_1 = \frac{c_4 C_4(4)}{|B(L(\frac{d}{4}), \frac{d}{4})|}$. We have obtained the desired (3.4).

For (3.5), we observe that the function $f(t) = |D\varphi(L(t))|$ is convex and increasing for $t < d$. Let $|D\varphi(x_1)| = \min_{\Omega'} |D\varphi(x)|$. Then $|D\varphi(x_1)| = f(t_1)$ where $t_1 = |x_1 - a| - |a - x_0|$. Since $f(0) > M$, $t_1 < 0$ and by triangle inequality $-t_1 \leq \text{diam}(\Omega')$. Now since f is convex it holds that

$$f\left(\frac{d}{2}\right) - f(0) \geq \frac{d}{-2t_1}(f(0) - f(t_1)). \quad (3.11)$$

Replacing f by its values, we see that (3.11) is indeed (3.5) with $c_2 = \frac{d}{2\text{diam}(\Omega')}$. Therefore the proof is concluded in the case $\max_{\Omega'} |D\varphi| > M$.

Case $\min_{\Omega'} |D\varphi| < m$.

By considering the inverse mapping φ^{-1} we will reduce the situation to the case $\max_{\Omega'} |D\varphi| \geq M$. Since $\varphi \in \mathcal{M}_n^{E'}(\Omega)$ we can apply Proposition 2.1 to find a constant $s = s(E', \Omega', \Omega)$ such that $\text{dist}(\varphi(\Omega'), \partial\varphi(\Omega)) \geq s$. Hence we can argue as in the previous case with φ^{-1} in the place of φ and $\frac{1}{m}$ in the place of M . We obtain that if $\inf_{\Omega'} |D\varphi(x)| < m$, there exists $c_5 = c_5(E, s)$, $R \in SO(n)$ and $B \subset \varphi(\Omega) \setminus \varphi(\Omega')$ with $|B| \geq c_6 s^n$ such

$$\int_{\varphi(\Omega')} \left| \frac{1}{m}R - D\varphi^{-1} \right|^2 \leq c_5 \int_B \text{dist}_{\frac{1}{m}SO(n)}^2(D\varphi^{-1}) \quad (3.12)$$

and for all $x \in B$ $|D\varphi^{-1}(x)| \geq \frac{1}{m}$. Now recall that if $\varphi \in \mathcal{M}_n^{E'}(\Omega)$ $m^n \leq J_\varphi \leq M^n$ and that for $\rho \in \mathbb{R}$ $\text{dist}_{\rho SO(n)}(D\varphi(x)) = |\rho - |D\varphi(x)||$. Therefore we can change variables in 3.12 to obtain that

$$\int_{\Omega'} |mR^{-1} - D\varphi|^2 \leq c_7 \int_{\varphi^{-1}(B)} \text{dist}_{mSO(n)}^2(D\varphi) \quad (3.13)$$

with $c_7 = c_7(E, E', \Omega', \Omega)$. We also used that $|A - B| = |A||B||A^{-1} - B^{-1}|$. Finally for $x \in \varphi^{-1}(B)$ $|D\varphi(x)| < m$ and hence $\text{dist}_{mSO(n)}^2(D\varphi(x)) = \text{dist}_E^2(D\varphi(x))$. Thus, the desired estimate

$$\int_{\Omega'} |mR^{-1} - D\varphi|^2 \leq c_7 \int_{\varphi^{-1}(B)} \text{dist}_E^2(D\varphi) \quad (3.14)$$

follows and the proof is concluded. \square

In the following example $E = E^{m,M}$.

Example 3.3. *There exists a sequence $\varphi_j \in \mathcal{M}_n(B_1)$ such that*

$$\lim_{j \rightarrow \infty} \frac{\inf_{\psi \in \mathcal{M}_n^E(B_1)} \int_{B_1} |D\varphi_j - D\psi|^2}{\int_{B_1} \text{dist}_E^2(D\varphi_j)} = \infty.$$

Proof:

Let $\varphi = r \frac{x - \lambda e_1}{|x - \lambda e_1|^2}$ with λ and r chosen so that $|D\varphi(e_1)| = M$ and $|D\varphi(-e_1)| = m$ where $e_1 = (1, 0, \dots, 0)$. Let us consider the sequence $\varphi_j = r \frac{x - (\lambda - \frac{1}{j})e_1}{|x - (\lambda - \frac{1}{j})e_1|^2}$ and set $t_j = |D\varphi_j(e_1)| - M$. On one hand, we have that

$$\begin{aligned} \int_{B_1} \text{dist}_E^2(D\varphi_j) &\leq t_j^2 |\{x \in B_1 : \text{dist}_E^2(D\varphi_j) > 0\}| \\ &= t_j^2 |\{x \in B_1 : |D\varphi_j| \geq M\}|. \end{aligned} \quad (3.15)$$

On the other hand, let $\psi \in \mathcal{M}_n^E(B_1)$ and φ_j defined as above with $\frac{1}{j} \leq \frac{\lambda-1}{2}$. Then φ_j and $\psi \in \mathcal{M}_n(hB)$ for some $h > 1$, h not depending of j . Thus, Proposition 2.2 yields a constant $c_1 = C_4(h)$ such that

$$\begin{aligned} t_j^2 &= (|D\varphi_j(e_1)| - M)^2 \leq |D\varphi_j(e_1) - D\psi(e_1)|^2 \\ &\leq c_1 \int_{B_1} |D\varphi_j - D\psi|^2. \end{aligned} \quad (3.16)$$

If we put (3.15) and (3.16) together, we obtain that

$$\frac{\inf_{\psi \in \mathcal{M}_n^E(B_1)} \int_{B_1} |D\varphi_j - D\psi|^2}{\int_{B_1} \text{dist}_E^2(D\varphi_j)} \geq \frac{c_1}{|\{x \in B_1 : |D\varphi_j| \geq M\}|},$$

which proves the claim. \square

4 Proof of Theorem 1.4 in a ball

The starting point for the proof of Theorem 1.1 in [7] is that thanks to the structure of the set $SO(n)$ it suffices to prove Theorem 1.1 for mappings whose components are uniformly Lipschitz harmonic functions. We start along the same lines than in [7] since E enjoys a nice structure as well.

4.1 Reduction to Lipschitz mappings

Proposition 4.1. *Let $E \subset E^{m,M}$. There exists constants $C_9 = C_9(n)$, $C_{10} = C_{10}(M)$ such that if $v \in W^{1,2}(B)$ there exists $v_M \in W^{1,\infty}(\Omega)$ such that*

- $\|Dv_M\|_{L^\infty(B)} \leq C_9 M$ and
- $\int_B |Dv - Dv_M|^2 \leq C_{10} \int_B \text{dist}_E^2(Dv)$.

Proof: The appendix A.1 in [7] yields v_M satisfying

- $\|Dv_M\|_{L^\infty(B)} \leq c_1 M$
- $\int_B |Dv - Dv_M|^2 \leq c_2 \int_{\{x \in B: |Dv| \geq 2M\}} |Dv|^2$.

Since if $|A| \geq 2M$, $|A| \leq 2\text{dist}_E(A)$ we are done. \square

4.2 Reduction to an F-harmonic mapping

As $SO(n)$ is related to harmonic functions so is $CO_+(n)$ to n-harmonic functions. However since the n -harmonic equation is degenerate the arguments below yield a wrong exponent in (4.6). Instead we take advantage of E being a compact subset of $CO_+(n)$ bounded away from 0. Then we can modify $|\cdot|^n$ near 0 and ∞ to relate E to minimizers of a variational problem $\int_\Omega F(Dv)$, where F has the appropriate growth.

Throughout the section the n -tuple of n -vectors (A_1, A_2, \dots, A_n) stands for the matrix with rows (A_1, A_2, \dots, A_n) and $(\text{Cof}(A))_{ij} = \partial_{ij} \det(A)$.

Lemma 4.2. *There exists a function $\psi : \mathbb{R} \rightarrow \mathbb{R}$ such that the function $F(A) : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by $F(A) = \psi(|A|)$ satisfies $DF(A) \leq C_F(1 + |A|)$ and $\frac{1}{C_F} \leq D^2F(A) \leq C_F$ with $C_F = C_F(E)$. Moreover, for every $A \in E$ it holds that $(DF(A_1), DF(A_2), \dots, DF(A_n)) = \text{Cof}(A)$. Equivalently $\text{Cof}(A) = D\tilde{F}$ where $\tilde{F}(A) = \sum_{i=1}^n F(A_i)$.*

Proof: Let $F(A) = \psi(|A|)$. Then a direct calculation gives that

$$\begin{aligned} DF(A) &= \psi'(|A|) \frac{A}{|A|}, & D^2F(A) &= \psi''(|A|) \left(\frac{A}{|A|} \otimes \frac{A}{|A|} \right) \\ & & &+ \frac{\psi'(|A|)}{|A|} \left(I - \frac{A}{|A|} \otimes \frac{A}{|A|} \right). \end{aligned} \quad (4.1)$$

If we put

$$\psi_1(x) = n(n-1) \left(\int_0^x \int_0^y z^{n-2} \chi_{(0,M)}(z) + (1 - \chi_{(0,M)}(z)) M^{n-2} dz dy \right),$$

$\psi_1(x) = x^n \chi_{0,M} + ((1 - \chi_{(0,M)}(x)) ax^2 + bx + c)$ with $a, b \geq 0$ and it satisfies the claims of the Lemma 4.2 except in a neighborhood of the origin. Then we consider $\psi_2(x) = \max\{(\frac{m_1}{2})^{n-2} x^2, \psi_1(x)\}$. This new function satisfies the claims but it is not smooth. We repair this by replacing ψ_2 by a smooth approximation of it in a neighborhood of $\frac{m_1}{2}$. Let r and ϵ be small numbers,

and $\varphi \in C_0^\infty(\frac{m_1}{2} - 2r, \frac{m_1}{2} + 2r)$ $\varphi = 1$ on $(\frac{m_1}{2} - r, \frac{m_1}{2} + r)$ and let ρ_ϵ be an approximation of the identity. Then setting

$$\psi_{\epsilon,r} = \varphi(\rho_\epsilon * \psi_2) + (1 - \varphi)\psi_2$$

for ϵ and r small enough it is not hard to check that $F(A) = \psi_{\epsilon,r}(|A|)$ satisfies all the conditions concerning regularity in the statement of the lemma.

Finally, if $A = (A_1, A_2, \dots, A_n) \in CO_+(n)$, $\det(A) = |A|^n = |A_i|^n$ where $|A_i|$ is the Euclidean norm of the vector A_i . Thus, if $A \in E$ we have that $\tilde{F}(A) = n \det(A)$ for $A \in E$ and that $D\tilde{F}(A) = (DF(A_1), DF(A_i), DF(A_n)) = \text{Cof}(A)$ as claimed. \square

Set $h(A) = D\tilde{F}(A) - \text{Cof}(A)$.

Lemma 4.3. *Let $v \in W^{1,2}(B, \mathbb{R}^n)$. Then*

$$v = w + z,$$

where z is an F -harmonic mapping in B and $w \in W_0^{1,2}(B, \mathbb{R}^n)$ is such that

$$\int_B |Dw|^2 \leq C_F \int_B |h(Dv)|^2.$$

Proof: To obtain the decomposition we solve the following system

$$\begin{cases} \text{Div}(D\tilde{F}(Dz)) = 0 & \text{in } B \\ z = v & \text{on } \partial B. \end{cases} \quad (4.2)$$

Here the operator Div means taking the divergence of the rows of matrix. Observe that by the definition of \tilde{F} the system decouples so that in fact we are solving n scalar equations. In the case $D\tilde{F}(A) = A$, (4.2) is equivalent to $\text{Div}(Dw) = \text{Div}(h(Dv))$ for $w = v - z$. In this nonlinear situation (4.2) means that

$$\begin{cases} \text{Div}(A(x, Dw)) = -\text{Div}(h(Dv)) & \text{in } B \\ w = 0 & \text{on } \partial B, \end{cases} \quad (4.3)$$

where $A(x, \xi) = D\tilde{F}(Dv(x) - \xi) - D\tilde{F}(Dv(x))$ is an \mathcal{A} -harmonic type of operator. From the uniform convexity of F , and the subsequent strong monotonicity of DF it follows that

$$\int_B |Dw|^2 \leq C_F \int_B \langle D\tilde{F}(Dv - Dw) - D\tilde{F}(Dv), Dw \rangle. \quad (4.4)$$

Plugging (4.3) into (4.4) and using Hölder's inequality we obtain

$$\int_B |Dw|^2 \leq C_F \int_B |h(Dv)|^2, \quad (4.5)$$

as desired. \square

Remark 4.4. Let v be CM -Lipschitz. Since h is uniformly Lipschitz in $B(0, 2M)$, we have that $|h(Dv)| \leq c_1 \text{dist}_E(Dv)$ and therefore Lemma 4.3 reads as

$$\int_B |Dw|^2 \leq C_{11} \int_B |\text{dist}_E(Dv)|^2. \quad (4.6)$$

In addition,

$$\int_B |Dz|^2 \leq \int_B |Dv|^2 + |\text{dist}_E(Dv)|^2.$$

Now if v is C_9M -Lipschitz $\text{dist}_E(Dv) \leq c_2(M + m)$ and thus

$$\int_B |Dz|^2 \leq C_{12},$$

with $C_{11} = C_{11}(M)$, $C_{12} = C_{12}(M)$.

4.3 Compactness

In the case of $SO(n)$ it is proved in [7] the existence of a constant C such that for any harmonic function z , there is $R \in SO(n)$ such that

$$\|R - Dz\|_{L^\infty(hB)} \leq C \left(\int_B \text{dist}_{SO(n)}^2(Dz) \right)^{\frac{1}{4}}, \quad (4.7)$$

for $h < 1$. In our setting, we do not know how to obtain this type of explicit estimate. However, the regularity of F -harmonic functions allows to use a compactness argument to prove an expression like (4.7), where $(\int_B \text{dist}_{SO(n)}^2(Dz))^{\frac{1}{4}}$ is replaced by $\rho_1(\int_B \text{dist}_E^2(Dz))$. Here, $\rho_1(\epsilon)$ is an unknown increasing function that at least converges to 0 as ϵ goes to 0.

Lemma 4.5. *Let $h < 1$ and B_1 the unit ball. There exists a function $\rho_1(\epsilon)$ with $\lim_{\epsilon \rightarrow 0} \rho_1(\epsilon) = 0$ such that for every F -harmonic mapping z there exists, $\varphi \in \mathcal{M}_n^E(B_1)$ satisfying*

$$\|D\varphi - Dz\|_{L^\infty(hB_1)} \leq \rho_1 \left(\int_{B_1} \text{dist}_E^2(Dz) \right). \quad (4.8)$$

Proof: Let

$$\delta(z) = \inf_{\varphi \in \mathcal{M}_n^E(B_1)} \|D\varphi - Dz\|_{L^\infty(hB_1)}, \text{ and}$$

$$\rho_1(\epsilon) = \sup\{\delta(z) : z \text{ F-harmonic}, \int_{B_1} \text{dist}_E^2(Dz) \leq \epsilon\}.$$

Clearly ρ_1 is a positive bounded decreasing function whence $\lim_{\epsilon \rightarrow 0} \rho_1(\epsilon) = \eta$ exists. Let us argue by contradiction by assuming that $\eta > 0$. This means that there exists a sequence of F-harmonic mappings $\{z_j\}$ such that $\delta(z_j) \geq \frac{\eta}{2}$. But

$$\int_{B_1} \text{dist}_E^2(Dz_j) \leq \frac{1}{j}, \quad (4.9)$$

i.e. $\lim_{j \rightarrow 0} \int_{B_1} \text{dist}_E^2(Dz_j) = 0$. Since E is a compact set it follows that $\int_{B_1} |Dz_j|^2 \leq c_1$, $c_1 = c_1(E)$. Therefore there is no loss of generality assuming that $Dz_j \rightarrow D\varphi$ weakly in $L^2(B_1)$ for some $\varphi \in W^{1,2}(\Omega)$. Furthermore, by Proposition 2.5 and Ascoli-Arzelà theorem Dz_j converge to $D\varphi$ uniformly on compact subsets of B_1 . This together with (4.9) this yields that $D\varphi \in E$ a.e.x. Given that E is bounded, $D\varphi \in W^{1,\infty}(B_1)$ and therefore, by Theorem 1.3, $\varphi \in \mathcal{M}_n^E(B_1)$. Finally, $Dz_j \rightarrow D\varphi$ uniformly in hB_1 , which contradicts that $\eta > 0$ and proves the claim. \square

4.4 Linearization and Local estimate

We start this section by recalling how the geometry of $CO_+(n)$ and of \mathcal{M}_n are related (a detailed account of this is given in [20]). We denote the tangent plane to $CO_+(n)$ at the identity matrix by $TCO_+(n)$. It can be seen that

$$A \in TCO_+(n) \iff \frac{A + A^t}{2} = \frac{1}{n} \text{Tr}(A)I.$$

Thus,

$$\Theta(A) = \left| \frac{A + A^t}{2} - \frac{1}{n} \text{Tr}(A)I \right|$$

is equal to the distance from A to $TCO_+(n)$. It follows from the definition of Θ that the set of functions $\{u : Du \in TCO_+(n)\}$ is a finite dimensional vector space of dimension $d = \frac{(n+2)(n+1)}{2}$. We denote it by Σ_n . Now recall that \mathcal{M}_n is a Lie group isomorphic to the classical group of matrices $SO_o(n+2, 1)$. Alternatively one can arrive to Σ_n as the Lie Algebra of \mathcal{M}_n isomorphic to

$so(n+2, 1)$. Let $\{u_i\}_{i=1}^d$ be a basis of Σ_n . Then we consider the projection $\Pi_{\Sigma_n, B_1} : W^{1,p}(B_1) \rightarrow \Sigma_n$ defined by

$$f \rightarrow \sum_{i=1}^d u_i \frac{1}{|B_1|} \int_{B_1} \langle f, u_i \rangle. \quad (4.10)$$

We will need the following Korn type of inequality first proved by Reshetnyak [19]. We refer to [20] for an extended discussion about this type of linear operators.

Theorem 4.6 ([20, Chapter 3, Theorem 3.2]). *Let $1 < p < \infty$. Then There exists a constant $C_{13} = C_{13}(p)$ such that for every function $g \in W^{1,p}(B, \mathbb{R}^n)$ we have that*

$$\|D(g) - D\Pi_{\Sigma_n}(g)\|_{L^p(B)} \leq C_{13} \|\Theta(D(g))\|_{L^p(B)}.$$

The following proposition and corollary originate from Reshetnyak's study of the stability of the Möbius group in terms of quasiconformality when the distortion is closed to 1 [20]. It is based on the following consideration: If a mapping g happens to satisfy that $\Pi_{\Sigma_n}(g) = 0$, then Theorem 4.6 implies that $\|D(g)\|_{L^p(B)} \leq \|\Theta(D(g))\|_{L^p(B)}$. We would like to apply this argument to the mapping $f - x$. Of course, $\Pi_{\Sigma_n}(f - x)$, does not need to be 0. In Proposition 4.7 we use an homotopy argument to prove that if $f - x$ is small, we can find a desired φ so that $\Pi_{\Sigma_n}(f \circ \varphi - x) = 0$. Let us remark that for this it is not needed to control $Df - I$. However to replace $\text{dist}_{TCO_+(n)}(A - I)$ by $\text{dist}_{CO_+(n)}(A)$, it is needed that $DA - I$ is small, and hence we need to have control over $D(f \circ \varphi) - I$. It turns out that this is rather complex in the quasiconformal setting [20] but it is much simpler in our situation. This is due to the fact that the modulus of continuity of Df can be assumed to be bounded independently of f .

In the Corollary 4.8 we use the Taylor expansion of $\text{dist}_{CO_+(n)}^2$ to express the local estimate in the form needed for the next section.

Proposition 4.7. *Let $0 \leq h \leq 1$, γ_1, γ_2 positive constants, $B = B(a, r)$ an arbitrary ball and $f \in C^{1,\alpha}((1+h)B)$ such that*

$$[Df]_{C^\alpha((1+h)B)} \leq \gamma_1 r^{-\alpha}.$$

Then there exists a positive number $\delta_1(h, \gamma_2)$, a non increasing function $\rho_2(\epsilon)$ with $\lim_{\epsilon \rightarrow 0} \rho_2(\epsilon) = 0$ and a constant C_{14} such that, if

$$\|Df - I\|_{L^\infty((1+h)B)} \leq \epsilon \leq \delta_1$$

then, there exists $\varphi \in \mathcal{M}_n(2B)$ satisfying the next three properties:

1. $\|D(f \circ \varphi - x)\|_{L^2(B)} \leq C_{14} \|\Theta(D(f \circ \varphi - x))\|_{L^2(B)}$,
2. $\|D(f \circ \varphi - x)\|_{L^\infty(B)} \leq \rho_2(\epsilon)$ and
3. $\|\varphi - x\|_{W^{1,\infty}(2B)} \leq r\gamma_2$.

The function ρ_2 depends on the modulus of continuity of Df and $C_{14} = C_{13}(2)$, where C_{13} was introduced in Theorem 4.6.

Proof:

Since both the assumptions and the claim depend on Df , there is no loss of generality assuming that $f(0) = 0$. We can also assume that $B = B_1$. The case of an arbitrary ball follows by composing with similarities.

For every continuous mapping we define $\Lambda_f : \Sigma_n \rightarrow \Sigma_n$ by $u \rightarrow \Pi_{\Sigma_n}(f \circ \varphi_u - x)$ where $\Pi_{\Sigma_n} = \Pi_{\Sigma_n, B_1}$ is the projection from $W^{1,p}(B_1, \mathbf{R}^n) \rightarrow \Sigma_n$ defined in (4.10) and $\varphi_u = \exp(u)$. Here \exp is the exponential mapping from Σ_n to \mathcal{M}_n (Observe that using the isomorphism between \mathcal{M}_n and $SO_o(n+2, 1)$ it is not hard to obtain an explicit expression of \exp . Denoting by F the isomorphism between \mathcal{M}_n and $SO_o(n+2, 1)$, $\exp(u) = F^{-1}(\exp(dF(u))$, where \exp is the matrix exponential in $M^{n+2 \times n+2}$). The following properties of the exponential will be useful:

- a) Let us declare $M(s) = \max\{\|\varphi_u(x) - x\|_{W^{1,\infty}(2B)} : u \in B(0, s) \subset \Sigma_n\}$. Then, since all the topologies in \mathcal{M}_n are the same,

$$\lim_{s \rightarrow 0} M(s) = 0. \quad (4.11)$$

- b) Let us state $\Lambda = \Lambda_f$. One of the properties defining the exponential mapping is that $\frac{d}{dt}\varphi_{tu}(x)|_0 = u(x)$. It is easy to see that this implies that for any

$$h \in \Sigma_n, \quad \Lambda'(0)h = h.$$

Hence, the inverse function theorem tells us that there exists $s_0 > 0$ such that Λ maps $B(0, s_0)$ homeomorphically into Σ_n . In particular, $\deg_\Lambda(0, B(0, s_0))$, the topological degree of Λ at zero in $B(0, s_0)$, is 1.

Due to (4.11) we can assume without loss of generality that

$$M(s_0) \leq \gamma_2. \quad (4.12)$$

Let $s \leq s_0$, then we have that

$$\begin{aligned} |\Lambda_f(u) - \Lambda(u)|_{|B(0,s)} &= |\Pi_{\Sigma_n}(f \circ \varphi_u - \varphi_u)| \\ &\leq \|\Pi_{\Sigma_n}\| \|\varphi_u\| \|Df - I\|_{L^\infty((1+h)B)} \leq c_1 \epsilon, \end{aligned} \quad (4.13)$$

where $\|\cdot\|$ stands for the operator norm of Π_{Σ_n} and

$$c_1 = \|\Pi_{\Sigma_n}\| \max_{x \in B, u \in B(0, s_0)} |\varphi_u(x)|. \quad (4.14)$$

By (4.12) we have that $c_1 \leq \|\Pi_{\Sigma_n}\|(1 + M(s_0)) \leq \|\Pi_{\Sigma_n}\|(1 + \gamma_2)$.

Let us declare

$$m(s) = \min_{\partial B(0, s)} |\Lambda(u)|.$$

Since Λ is an homeomorphism in $B(0, s_0)$ and $\Lambda(0) = 0$ we have that $0 < m(s)$ for every $r < s_0$. Moreover, since $\varphi_u \rightarrow x$ uniformly

$$\lim_{s \rightarrow 0} m(s) = 0. \quad (4.15)$$

Then we define s_ϵ by

$$s_\epsilon = \inf\{s : m(s) \geq c_1\epsilon + \epsilon\}.$$

where c_1 is defined in (4.14). Then, (4.15) implies that

$$\lim_{\epsilon \rightarrow 0} s_\epsilon \rightarrow 0. \quad (4.16)$$

Now, the definition of s_ϵ and (4.13) yield that if $s_\epsilon \leq s_0$, then

$$t\Lambda + (1 - t)\Lambda_f(u) \neq 0 \text{ for } u \in \partial B(0, s_\epsilon). \quad (4.17)$$

Thus, $\deg(0, B(0, s_\epsilon))$ remains constant through the homotopy (This is standard see [10] chapter 14 or [14]).

Let us declare $\delta_1 = \frac{m(s_0)}{c_1 + 1}$. If $\epsilon < \delta_1$, then $s_\epsilon < s_0$ and thus, $\deg_{\Lambda_f}(0, B(0, s_\epsilon)) = \deg_\Lambda(0, B(0, s_\epsilon)) = 1$. Therefore, there exists $u \in B(0, s_\epsilon)$ such that $\Lambda_f(u) = 0$. When we apply Theorem 4.6 to $f \circ \varphi_u - x$, (1) follows.

To prove (2) we observe that

$$\begin{aligned} |D(f \circ \varphi_u(x)) - I| &\leq |Df(x) - I| + |Df(x)||D\varphi_u(x) - I| \\ &\quad + |D\varphi_u(x)||Df(\varphi_u(x)) - Df(x)|. \end{aligned}$$

By definition we have that $|Df(x) - I| \leq \epsilon$, that $|D\varphi_u(x) - I| \leq M(s_\epsilon)$ and that $|\varphi_u(x) - x| \leq M(s_\epsilon)$. Hence, using the Hölder regularity of Df we obtain that

$$|D(f \circ \varphi_u(x) - x)| \leq \epsilon + (1 + \epsilon)M(s_\epsilon) + \gamma_1((1 + M(s_\epsilon))M(s_\epsilon)^\alpha) = \rho_2(\epsilon).$$

Finally, (4.16) and (4.11) imply that $\lim_{\epsilon \rightarrow 0} \rho_2(\epsilon) = 0$ as desired. The proposition is proved since $M(s_0) \leq \gamma_2$ by (4.12), which yields (3). \square

Corollary 4.8 (Local estimate). *Let $0 < h \leq 1$, $\gamma_1, \gamma_2 > 0$ and $B(a, r)$ a ball. Then there exists $\delta_2 = \delta_2(h, \gamma_1, \gamma_2, \alpha)$ and C_{15} such that if f satisfies*

1. $[Df]_{C^\alpha((1+h)B)} \leq \gamma_1 r^{-\alpha}$ and
2. $\|Df - I\|_{L^\infty((1+h)B)} \leq \delta$,

then there exists $\varphi \in \mathcal{M}_n(2B)$ with $\|\varphi(x) - x\|_{W^{1,\infty}(2B)} \leq r\gamma_2$ and

$$\int_B |D(f \circ \varphi) - I|^2 \leq C_{15} \int_B \text{dist}_{CO_+(n)}^2(D(f \circ \varphi)).$$

Proof:

The corollary is proved by plugging the results from the previous proposition into the Taylor expansion of $\text{dist}_{CO_+(n)}$. By Taylor we have that

$$|\text{dist}_{CO_+(n)}^2(A)| = c_1 |\Theta(A - I)|^2 + O(|A - I|^4).$$

Therefore,

$$|\Theta(A - I)|^2 \leq c_2 |\text{dist}_{CO_+(n)}^2(A)| + O(|A - I|^4). \quad (4.18)$$

We first require that $\delta_2 \leq \delta_1$ to apply Proposition 4.7 to the mapping f and find the desired φ . Using Proposition 4.7 (1), putting $A = D(f \circ \varphi)$ in (4.18), and rearranging the resulting expression yield

$$\begin{aligned} \int_B |D(f \circ \varphi) - I|^2 &\leq c_3 \int_B \text{dist}_{CO_+(n)}^2(D(f \circ \varphi)) \\ &\quad + c_4 \int_B |D(f \circ \varphi) - I|^4. \end{aligned} \quad (4.19)$$

We finally impose that δ also satisfies $c_4 \rho_2^4(\delta) \leq \frac{1}{2} \rho_2^2(\delta)$. Then Proposition 4.7 (2) implies that $c_4 |D(f \circ \varphi) - I|^4 \leq \frac{1}{2} |D(f \circ \varphi) - I|^2$. Hence, we can absorb $c_4 \int_B |D(f \circ \varphi) - I|^4$ to the left hand side to conclude that

$$\int_B |D(f \circ \varphi) - I|^2 \leq c_5 \int_B \text{dist}_{CO_+(n)}^2(D(f \circ \varphi)), \quad (4.20)$$

as desired. \square

4.5 Theorem 1.4 in a Ball

In this section we put all the information obtained before together to obtain the Theorem 1.4 in a ball. We state it as a separate proposition for later uses.

Proposition 4.9. *Let $E \subset CO_+(n)$ be compact, finitely connected with*

$$SO(n)E = E,$$

and $0 \notin E$. Let B a ball. Then there exists a constant $C_{16} = C_{16}(E)$ such that for any $v \in W^{1,2}(4B, \mathbb{R}^n)$ there exists $\varphi \in \mathcal{M}_n^E(B, \mathbb{R}^n)$ such that

$$\int_B |D\varphi - Dv|^2 \leq C_{16} \int_{4B} \text{dist}_E^2(Dv). \quad (4.21)$$

The sketch of the proof is the following: Let

$$\epsilon = \int_{4B} \text{dist}_E^2(Dv).$$

Using Subsections 4.1,4.2 one finds that it is enough to prove the proposition for z an F -harmonic mapping such that Dz is uniformly Hölder in $B' \subset 4B$. Next section 4.3 yields $\varphi_1 \in \mathcal{M}_n^E(4B)$ such that

$$\|D\varphi_1 - Dz\|_{L^\infty(B')} \leq \rho_1(\epsilon).$$

Then, changing variables with φ_1 we discover that the mapping $f = z \circ \varphi_1^{-1}$ fulfills the condition required in Corollary 4.8, the local estimate, in a ball $B'' \subset \varphi_1(B')$. Therefore Corollary 4.8 yields φ_2 such that

$$\begin{aligned} \int_{B'''} |D(z \circ \varphi_1^{-1} \circ \varphi_2) - I|^2 \leq \\ C \int_{B'''} \text{dist}_{CO_+(n)}^2(Dz \circ \varphi_1^{-1} \circ \varphi_2). \end{aligned} \quad (4.22)$$

where $B''' \subset B''$. Setting $\varphi_3 = \varphi_2^{-1} \circ \varphi_1$, we can change variables by φ_3 finding that

$$\int_{2B} |Dz - D\varphi_3|^2 \leq C \int_{4B} \text{dist}_{CO_+(n)}^2(Dz). \quad (4.23)$$

where $2B \subset \varphi_3^{-1}(B''')$. The existence of the balls B', B'', B''' fulfilling the properties used above is essentially a consequence of the Lemma 2.1.

Finally, since φ_3 might be in $\mathcal{M}_n(2B) \setminus \mathcal{M}_n^E(B)$ we use Proposition 3.2 to find $\varphi_4 \in \mathcal{M}_n^E(B)$ such that

$$\int_B |D\varphi_3 - D\varphi_4|^2 \leq C \int_{4B} \text{dist}_{CO_+(n)}^2(D\varphi_3).$$

From here, the proof is concluded by reiterative use of triangle inequality.

We now give the detailed argument.

Proof of Proposition 4.9 By a change of variables with a similarity it is enough to prove the estimate in $B_1 \subset 4B_1$. We start by using the Lemma 2.1 to select the appropriate balls. Let $0 < h_0, h_1, h_2 < 1$ be such for any $\varphi \in \mathcal{M}_n^E(4B)$ the following inclusions hold:

$$\begin{aligned} \varphi(2B) &\subset h_0\varphi(4B), & (h_0 + 3h_1)\varphi(4B) &\subset \varphi(4B) \text{ and} \\ (h_0 + 2h_1)\varphi(4B) &\subset \varphi(h_2 4B). \end{aligned} \quad (4.24)$$

Next, we reduce to a smooth mapping. Firstly we apply Proposition 4.1 to v in $4B$ to obtain a C_9M Lipschitz mapping v_M . We prove Proposition 4.9 for v_M . By the Lipschitz bound we have that

$$\int_B |Dv_M - m|^2 \leq c_1,$$

$c_1 = c_1(E)$. Hence we can assume that

$$\int_{4B} \text{dist}_E(Dv_M)^2 \leq \epsilon_0, \quad (4.25)$$

for $0 < \epsilon_0$ to be determined later. For mappings v_M such that 4.25 does not hold, 4.21 is fulfilled with $\varphi(x) = mx$ and $C_{16} = \frac{1}{\epsilon_0}c_1$.

Let z be the F-harmonic mapping obtained by applying Lemma 4.3 to v_M in $4B$. Since $\int_{4B} |Dz|^2 \leq C_{12}$ by Remark 4.4, Proposition 2.5 yields that for any $h < 1$

$$[Dz]_{C^\alpha(h_2 4B)} \leq c_2, \quad (4.26)$$

with $c_2 = c_2(C_6, C_{12}, h)$. Furthermore, the inequality $\text{dist}_E^2(M_1) \leq c_3(\text{dist}_E^2(M_1 + M_2) + |M_2|^2)$, (4.6) and (4.25) imply that

$$\int_{4B} \text{dist}_E^2(Dz) \leq c_4\epsilon_0.$$

Next, we apply Lemma 4.5 to find the existence of $\varphi_1 \in \mathcal{M}_n^E(4B)$ such that

$$\|Dz - D\varphi_1\|_{L^\infty(h_2 4B)} \leq \rho_1(\epsilon), \quad (4.27)$$

where $\epsilon = \int_{4B} \text{dist}_E^2(Dz)$.

We want to translate these estimates to the domain $\varphi_1(4B)$. It follows from Lemma 2.1 that

$$c_5 < |\varphi_1(4B)| \leq c_6, \quad (4.28)$$

$c_5 = c_5(E), c_6 = c_6(E)$. Define $r_1 > 0$ by,

$$|B(0, 1)|r_1^n = c_6 \quad (4.29)$$

to guarantee that $\text{diam}(\varphi_1(4B)) \leq 2r_1$.

Using that $\varphi \in \mathcal{M}_n^E(4B)$ we deduce

$$\begin{aligned} \|D(z \circ \varphi_1^{-1}) - I\|_{L^\infty(\varphi_1(h_2 4B))} &\leq \|D\varphi_1^{-1}\|_{L^\infty((\varphi_1(h_2 4B)))} \|Dz - D\varphi_1\|_{L^\infty(h_2 4B)} \\ &\leq \frac{1}{m} \rho_1(\epsilon). \end{aligned} \quad (4.30)$$

In addition, by (4.26) we have

$$\begin{aligned} [D(z \circ \varphi_1^{-1})]_{C^\alpha(\varphi_1(h_2 4B))} &\leq \|D\varphi_1^{-1}\|_{L^\infty((\varphi_1(h_2 4B)))} [Dz]_{C^\alpha(h_2 4B)} \\ &\leq \frac{c_7}{m}, \end{aligned}$$

with $c_7 = c_7(h_2, M)$. Furthermore, by Lemma 2.1, Proposition 2.2 and (4.28), $D\varphi_1^{-1}$ is uniformly continuous in $\varphi_1(h_2 4B)$. Hence, there exists $c_8 = c_8(h_2, E)$ such that

$$[D(z \circ \varphi_1^{-1})]_{C^\alpha(\varphi_1(h_2 4B))} \leq c_8 \leq c_9 r_1^{-\alpha} \quad (4.31)$$

by 4.29.

Since by (4.24) $(h_0 + 2h_1)\varphi_1(4B) \subset \varphi_1(h_2 4B)$, we intend to apply Corollary 4.8 with $B = (h_0 + h_1)\varphi_1(4B)$, $f = z \circ \varphi_1$, $h = h_1$, $\gamma_1 = c_9$ and $\gamma_2 = \frac{h_1}{r_1}$. Now, by (4.25) we can impose for ϵ_0 that $\rho_1(\epsilon_0) \leq m\delta(E, h_1)$. Thus, (4.30) and (4.31) imply that all the requirements of Corollary 4.8 are fulfilled. Hence, we are provided with $\varphi_2 \in \mathcal{M}_n$ such that

$$\begin{aligned} \int_{(h_0+h_1)\varphi_1(4B)} |D(z \circ \varphi_1^{-1} \circ \varphi_2) - I|^2 &\leq c_{10} \int_{(h_0+h_1)\varphi_1(4B)} \text{dist}_{CO_+(n)}^2(Dz \circ \varphi_1^{-1} \circ \varphi_2). \end{aligned} \quad (4.32)$$

with $c_{10} = C_{15}$. Let $\varphi_3 = \varphi_2^{-1} \circ \varphi_1$. By Proposition 4.7 (3) and our choice of γ_2 we have

$$\|\varphi_2 - x\|_{W^{1,\infty}((h_0+h_1)\varphi_1(4B))} \leq h_1. \quad (4.33)$$

In addition, $\varphi_1 \in \mathcal{M}_n^E(h_2 4B)$. It follows that $\varphi_3^{-1} = \varphi_1^{-1} \circ \varphi_2 \in \mathcal{M}_n^{E'}((h_0 + h_1)\varphi_1(4B))$ for some $E' = E'(E)$. Therefore, there exist constants $c_{11} = c_{11}(E)$, $c_{12} = c_{12}(E)$ such that

$$c_{11} J_{\varphi_3^{-1}}(x) = c_{11} |D\varphi_3^{-1}(x)|^n \leq |D\varphi_3^{-1}(x)|^2 \leq c_{12} |D\varphi_3^{-1}(x)|^n = c_{12} J_{\varphi_3^{-1}}(x).$$

Hence if we change variables in (4.32) by φ_3^{-1} we obtain

$$\int_{\varphi_3^{-1}((h_0+h_1)\varphi_1(4B))} |Dz - D\varphi_3|^2 \leq c_{13} \int_{\varphi_3^{-1}((h_0+h_1)\varphi_1(4B))} \text{dist}_{CO_+(n)}^2(Dz)$$

where $c_{13} = c(c_{11}, c_{12})$. Now we investigate what is $\varphi_3^{-1}((h_0 + h_1)\varphi_1(4B))$. By (4.33) we have that that

$$h_0\varphi_1(4B) \subset \varphi_2((h_0 + h_1)\varphi_1(4B)) \subset (h_0 + 2h_1)\varphi_1(4B).$$

On the other hand, $\varphi_1 \in \mathcal{M}_n^E(4B)$ so (4.24) implies that

$$2B \subset \varphi_1^{-1}(h_0\varphi_1(4B)) \subset \varphi_3^{-1}((h_0 + h_1)\varphi_1(4B)) \subset 4B.$$

Thus,

$$\int_{2B} |Dz - D\varphi_3|^2 \leq c_{13} \int_{4B} \text{dist}_{CO+(n)}^2(D(z)) \quad (4.34)$$

as desired.

The last obstruction is that $\varphi_3 = \varphi_2^{-1} \circ \varphi_1$ is not necessarily in $\mathcal{M}_n^E(2B)$ due to φ_2^{-1} . However (4.33) implies that $\varphi_3 \in \mathcal{M}_n^{E'}(2B)$ for E' depending on E . Since $\text{dist}_E^2(D\varphi_3) \leq \text{dist}_E^2(Dz) + |Dz - D\varphi_3|^2$ we can apply Lemma 3.2 to find $\varphi \in \mathcal{M}_n^E(B)$ such that

$$\begin{aligned} \int_B |D\varphi - Dz|^2 &\leq \int_B |D\varphi - D\varphi_3|^2 + |Dz - D\varphi_3|^2 \\ &\leq c_{14} \int_{4B} \text{dist}_E^2(Dz), \end{aligned} \quad (4.35)$$

where $c_{14} = 2c_{13} + C_8$.

Lastly, we translate the estimate from z to v .

$$\begin{aligned} \int_B |Dv - D\varphi_4|^2 &\leq C \int_B |Dv - Dv_M|^2 \\ &\quad + \int_B |Dv_M - Dz|^2 + \int_B |Dz - D\varphi_4|^2, \end{aligned} \quad (4.36)$$

Therefore, using again that $\text{dist}_E^2(A) \leq \text{dist}_E^2(B) + |A - B|^2$ for $A, B \in M^{n \times n}$ together with Proposition 4.1, Remark 4.4 and the estimate (4.35) we can conclude that

$$\int_B |Dv - D\varphi_4|^2 \leq c_{15} \int_{4B} \text{dist}_E^2(Dv), \quad (4.37)$$

where $c_{15} = c_{15}(C_{10}, C_F, c_{14})$. \square

5 Proof of Theorem 1.4

The arguments in Section 4 can be modified to be applied to a compact set Ω' instead of a ball, but several technical problems arise. Therefore, we

have preferred to keep Section 4 simple and prove Theorem 1.4 for a generic compact set by a covering argument. The idea is as follows: We can suppose that the compact set Ω' is a finite union of overlapping balls $\{B^i\}$. Therefore we can apply Proposition 4.9 to each B^i . This yields a family of Möbius transforms $\{\varphi_i\}$ such that $D\varphi_i$ is close to Dv in B^i . This implies that if we take two balls B^i, B^j $D\varphi_i$ is close to $D\varphi_j$ in $B^i \cap B^j$. The fundamental property of Möbius mappings used here is Lemma 2.3. Lemma 2.3 says that if we choose the parameters correctly, $D\varphi_j$ is close to $D\varphi_i$ not only in $B^i \cap B^j$ but in the whole of Ω' . Therefore if we choose a ball B^0 , $D\varphi_0$ is close to $D\varphi_i$ for any B^i such that B^i intersects B^0 . Moreover, given an arbitrary ball B' we can link it with B^0 by a finite chain of pairwise intersecting balls. It is not hard to see that it follows that the Möbius map φ' corresponding to B' is close to φ_0 in Ω' as well. Therefore all the φ_i are close to each other in Ω' and hence any of them is close to v in the whole Ω' . The following proposition shows that this can be made rigorous.

Proposition 5.1. *Let $\Omega' \Subset \Omega \subset \mathbb{R}^n$ with Ω connected. Then there exists a constant $C_{17} = C_{17}(E, \Omega', \Omega)$ and $E' \subset CO_+(n)$, E' depending on Ω' and E , such that for any $v \in W^{1,2}(\Omega, \mathbb{R}^n)$ there exists $\varphi \in \mathcal{M}_n^{E'}(\Omega)$ such that*

$$\int_{\Omega'} |D\varphi - Dv|^2 \leq C_{17} \int_{\Omega} \text{dist}_{E'}^2(Dv). \quad (5.1)$$

Proof:

Firstly, we observe that it is enough to prove the estimate assuming that

$$\int_{\Omega} \text{dist}_E^2(Dv) \leq \epsilon_0 \quad (5.2)$$

as in the proof of Proposition 4.9.

Let $r = \text{dist}(\Omega', \Omega)$. Since Ω' is compact there exists a family of balls $\{B(x_i, \frac{r}{16})\}_{i=1}^n$ such that $\Omega' \subset \cup_{i=1}^n B(x_i, \frac{r}{16})$ with $n = n(\Omega')$. Since Ω' can be assumed to be connected for each x_i there exist x'_i, y_i such that $B(y_i, \frac{r}{16}) \subset B(x_i, \frac{r}{8}) \cap B(x'_i, \frac{r}{8})$. We set $B^i = B(x_i, \frac{r}{8})$. Then $\Omega' \subset \cup_{i=1}^n B^i$ and $8B^i \subset \Omega$. In addition given any two balls $B^0, B' \in \{B^i\}_{i=1}^n$ there exist two chains of balls $\{B^j\}_{j=1}^m \subset \{B^i\}_{i=1}^n$ and, $\{P_j\}_{j=1}^m$ such that $B^1 = B^0, B^m = B'$, $2P_j \subset 2B^j \cap 2B^{j+1}$ and $|P_j| \geq cr^n$.

We apply Proposition 4.21 to $2B^0$ to find the existence of $\varphi_0 \in \mathcal{M}_n^E(2B^0)$ such that

$$\int_{2B^0} |Dv - D\varphi_0|^2 \leq c_1 \int_{8B^0} \text{dist}_E^2(Dv),$$

$c_1 = C_{16}$. We plan to show that $\varphi_0 \in \mathcal{M}_n^E(\Omega')$ and that

$$\int_{B'} |Dv - D\varphi_0|^2 \leq c_2 \int_{\Omega} \text{dist}_E^2(Dv). \quad (5.3)$$

Clearly, the thesis follows from (5.3) since $B' \in \{B^i\}_{i=1}^n$ was arbitrary.

Application of Proposition 4.21 in each of the balls $2B^j$, yields $\varphi_j \in \mathcal{M}_n^E(B^j)$ such that

$$\int_{2B^j} |Dv - D\varphi_j|^2 \leq \int_{\Omega} \text{dist}_E^2(Dv). \quad (5.4)$$

Then it is easy to see that proving (5.3) is equivalent to proving that

$$\int_{B'} |D\varphi_j - D\varphi_{j-1}|^2 \leq c_1 \int_{\Omega} \text{dist}_E^2(Dv), \quad (5.5)$$

for every $j = 1 \dots m$.

Set $\theta_j = \varphi_{j-1}^{-1} \circ \varphi_j$. Then (5.4) and Proposition 2.2 imply that for $x \in P_j$

$$\begin{aligned} \max_{\Omega'} \{|D\theta_j(x) - I|, |D\theta_j^{-1}(x) - I|\} &\leq c_3 |D\varphi_j - D\varphi_{j-1}|^2 \\ &\leq c_4 \int_{\Omega} \text{dist}_E^2(Dv) \end{aligned} \quad (5.6)$$

where in the first inequality we have used that $\varphi_j, \varphi_{j-1} \in \mathcal{M}_n^E(P_j)$.

Let $h = h(\Omega', \Omega)$ be such that for every $B, \tilde{B} \in \{B_i\}_{i=1}^n$ $2B \subset h\tilde{B}$. Let $\alpha(h)$ be given by Lemma 2.3. Then we firstly imposed that $\epsilon_0 c_4 \leq \alpha(h)$. Thus, by (5.6) and Lemma 2.3 we have that for $x \in B'$

$$|D\theta_j(x) - I|^2 \leq c_5 \int_{\Omega} \text{dist}_E^2(Dv), x \in B', \quad (5.7)$$

where $c_5 = C_5 c_4$, C_5 being defined in Lemma 2.3. To estimate $|D\varphi_j - D\varphi_{j-1}|$ by $c_6 |D\theta_j(x) - I|$ with $c_6 = c_6(E)$ it is enough that $\varphi_j, \varphi_{j-1} \in \mathcal{M}_n^{E'}(\Omega')$ for a set E' depending only on E . To achieve this we further require that $\epsilon_0 C_5 \leq \frac{1}{\text{diam}(\Omega)2n}$. This yields,

$$\max\{|\varphi_{j-1}^{-1} \circ \varphi_j(x) - x|, |\varphi_j^{-1} \circ \varphi_{j-1}(x) - x|\} < \frac{1}{4n}.$$

Therefore, $\varphi_j((\frac{3}{2} - \frac{1}{4n})B') \subset \varphi_{j-1}(\frac{3}{2}B') \subset \varphi_j((\frac{3}{2} + \frac{1}{4n})B')$. Thus, for every j $\varphi_m(\frac{5}{4}B') \subset \varphi_j(\frac{3}{2}(B')) \subset \varphi_m(2B')$. In particular this means that φ_j are finite in $(\frac{3}{2}B')$. Proposition 2.2 implies that for all x in B'

$$\begin{aligned} |D\varphi_j(x)| &\leq \frac{c_6}{|B'|} \int_{B'} |D\varphi| \leq c_6 \left(\frac{|\varphi_j(B')|}{|B'|} \right)^{\frac{1}{n}} \\ &\leq c_6 \left(\frac{|\varphi_m(2B')|}{|B'|} \right)^{\frac{1}{n}} \leq c_7. \end{aligned}$$

where $c_6 = C_4$, C_4 being from Proposition 2.2 and $c_7 = 2Mc_6$. We use here that $\varphi_m \in \mathcal{M}_n^E(2B')$. We can argue similarly for φ_j^{-1} obtaining that $\varphi_j \in \mathcal{M}_n^{E'}(B')$ as desired. Then, (5.3) follows from (5.7). The Proposition is proved. \square

Finally, Theorem 1.4 (i) is proved by Proposition 5.1 and Proposition 3.2. (ii) is proved combining Example 3.3 and Theorem 1.1. \square

Acknowledgments

We thank Stefan Müller for bringing the problem to our attention and for many interesting discussions and suggestions. This work was partially supported by was partially supported by the EU Research Training Networks *HYperbolic and Kinetic Equations*, contract HPRN-CT-2002-00282 and *Phase Transitions in crystalline solids*, contract FMRX-CT 98-0229. Part of this research took place while D.F was visiting Wuhan Institute of Physics and Mathematics, the Chinese Academy of Sciences in April, 2003. He would like to thank the institute for the hospitality.

References

- [1] Beardon, A. F. The geometry of discrete groups. Corrected reprint of the 1983 original. Graduate Texts in Mathematics, 91. Springer-Verlag, New York, (1995).
- [2] Chaudhuri, N; Müller, S. Rigidity estimate for two incompatible wells. *Calc. Var. Partial Differential Equations* 19 (2004), no. 4, 379–390.
- [3] De Giorgi, E. Sulla differenziabilità e l’analiticità delle estremali degli integrali multipli regolari. (Italian) *Mem. Accad. Sci. Torino. Cl. Sci. Fis. Mat. Nat.* (3) 3 (1957) 25–43.
- [4] Dal Maso, G; Negri, M; Percivale, D. Linearized elasticity as Γ -limit of finite elasticity. *Set-Valued Anal.* 10 (2002), 165-183.
- [5] Gehring, F. Rings and quasiconformal mappings in space. *Trans. Amer. Math. Soc.* 103 (1962) 353–393.
- [6] Giusti, E. *Direct methods in the calculus of variations.* World Scientific Publishing Co., Inc., River Edge, NJ, 2003.

- [7] Friesecke, G; James, R.D; Müller, S; Rigorous derivation of nonlinear plate theory and geometric rigidity. *C. R. Math. Acad. Sci. Paris* 334 (2002), no. 2, 173–178.
- [8] Friesecke, G; James, R.D, Müller, S. A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity. *Comm. Pure Appl. Math.* 55 (2002), 1461-1506.
- [9] Friesecke, G; James, R.D; Mora, M.G; Müller, S. Derivation of nonlinear bending theory for shells from three dimensional nonlinear elasticity by Gamma-convergence. *C. R. Acad. Sci. Paris, Sér. I* 336 (2003), 697-702.
- [10] Heinonen, J; Kilpeläinen, T; Martio, O. Nonlinear potential theory of degenerate elliptic equations. *Oxford Mathematical Monographs.* Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, (1993).
- [11] Iwaniec, T. P -harmonic tensors and quasiregular mappings. *Ann. of Math.* (2) 136 (1992), no. 3, 589–624.
- [12] Iwaniec, T; Martin, G. Quasiregular mappings in even dimensions. *Acta Math.* 170 (1993), no. 1, 29–81.
- [13] John, F. Rotation and strain. *Comm. Pure Appl. Math.* 14 1961 391–413.
- [14] Lloyd, N G. A survey of degree theory: basis and development. *IEEE Trans. Circuits and Systems* 30 (1983), no. 8, 607–616.
- [15] Liouville, J. Théorème sur l'équation $dx^2+dy^2+dz^2 = \lambda(d\alpha^2+d\beta^2+d\gamma^2)$. *J.Math. Pures Appl.*, 15, no 1, 103 (1850)
- [16] Mora, M.G; Müller, S. Derivation of the nonlinear bending-torsion theory for inextensible rods by Γ -convergence. *Calc. Var. Partial Differential Equations* 18 (2003), no. 3, 287–305.
- [17] Mora, M.G; Müller, S. A nonlinear model for inextensible rods as a low energy Gamma-limit of three-dimensional nonlinear elasticity. Preprint MPI-MIS n. 13/2003, Leipzig (Germany) (To appear in *Annales de l'Institut Henri Poincaré, Analyse Non Linéaire.*)
- [18] Reshetnyak, J. G. Liouville's conformal mapping theorem under minimal regularity hypotheses. (Russian) *Sibirsk. Mat. Ž.* 8 (1967) 835–840.

- [19] Reshetnyak, J. G. Estimates for certain differential operators with finite-dimensional kernel. (Russian) *Sibirsk. Mat. Ž.* 11 (1970) 414–428.
- [20] Reshetnyak, J. G. Stability theorems in geometry and analysis. Translated from the 1982 Russian original by N. S. Dairbekov and V. N. Dyatlov, and revised by the author. Translation edited and with a foreword by S. S. Kutateladze. *Mathematics and its Applications*, 304. Kluwer Academic Publishers Group, Dordrecht, (1994).