

Max-Planck-Institut  
für Mathematik  
in den Naturwissenschaften  
Leipzig

A variational model for dislocations in the line  
tension limit

by

*Adriana Garroni and Stefan Müller*

Preprint no.: 76

2004





# A variational model for dislocations in the line tension limit

Adriana Garroni  
Dipartimento di Matematica  
Università di Roma “La Sapienza”  
P.le Aldo Moro 3  
00185 Roma  
ITALY  
garroni@mat.uniroma1.it

Stefan Müller  
Max-Planck Institut für  
Mathematik in den Naturwissenschaften  
Inselstr. 22-26  
D-04103 Leipzig  
GERMANY  
Stefan.Mueller@mis.mpg.de

## Abstract

We study the interaction of a singularly perturbed multiwell energy (with an anisotropic nonlocal regularizing term of  $H^{1/2}$  type) and a pinning condition. This functional arises in a phase field model for dislocations which was recently proposed by Koslowski, Cuitiño and Ortiz but is also of broader mathematical interest. In the context of the dislocation model we identify the  $\Gamma$ -limit of the energy in all scaling regimes for the number  $N_\varepsilon$  of obstacles. The most interesting regime is  $N_\varepsilon \approx |\ln \varepsilon|/\varepsilon$ , where  $\varepsilon$  is a nondimensional length scale related to the size of the crystal lattice. In this case the limiting model is of line tension type. One important feature of our model is that the set of energy wells is periodic and hence not compact. A key ingredient in the proof is thus a compactness estimate (up to a single translation) for finite energy sequences, which generalizes earlier results of Alberti, Bouchitté and Seppecher for the two-well problem with an  $H^{1/2}$  regularization.

Keywords: Dislocations, Phase Transitions, Capacity,  $\Gamma$ -convergence, Line Tension

Mathematical Subjects Classification: 82B26, 31C15 49J45

## 1 Introduction

We study the functional

$$E_\varepsilon(u) = \frac{1}{\varepsilon} \int_Q W(u) dx + \iint_{Q \times Q} K_\nu(x-y) |u(x) - u(y)|^2 dx dy, \quad (1)$$

subject to the pinning condition

$$u = 0 \quad \text{on } B(x_\varepsilon^i, R\varepsilon) = B_{R\varepsilon}^i, \quad \text{for } i = 1, \dots, N_\varepsilon. \quad (2)$$

Here  $Q = (-1/2, 1/2)^2$  is the unit square,  $W$  is one-periodic, non-negative and vanishes exactly on the integers  $\mathbf{Z}$ , a typical choice being

$$W(u) = \text{dist}^2(u, \mathbf{Z}), \quad (3)$$

and the nonlocal part of the energy behaves like the  $H^{1/2}$  norm, i.e.  $K_\nu(z) \approx |z|^{-3}$ . Finally  $\varepsilon > 0$  is a small parameter and we study the limit  $\varepsilon \rightarrow 0$  (after suitable rescaling).

The above functional with the choice (3) has recently been proposed by Koslowski, Cuitiño and Ortiz as a phase-field model for dislocations. In this setting  $\varepsilon$  is the ratio of the spacing of the lattice planes and the size of the physical domain under consideration. In this context our main achievement is that we identify the relevant scaling regimes for the number of obstacles  $N_\varepsilon$  and the corresponding  $\Gamma$ -limits of the (suitably scaled) energy  $E_\varepsilon$ . Specifically if  $N_\varepsilon \approx \varepsilon^{-1} |\log \varepsilon|$  we show that the limit of  $E_\varepsilon / (\varepsilon N_\varepsilon)$  is the so called line-tension limit, i.e. the limit functional is defined on the space  $BV(Q; \mathbf{Z})$ , and is given by  $\int_{S_u} \gamma(n) d\mathcal{H}^1 + \int_Q D_\nu(u, B_R) dx$ , where  $S_u$  is the jump set of  $u$ , with normal  $n$ , and  $D_\nu(u, B_R)$  represents the limiting contribution of the obstacles  $B_{R\varepsilon}^i$  (see Theorem 10 below for a precise statement). If  $N_\varepsilon \ll \varepsilon^{-1} |\log \varepsilon|$  the line energy contribution dominates, only constant functions  $u \equiv a$ , with  $a \in \mathbf{Z}$ , are admissible in the limit and their energy is given by  $D(a, B_R)$ . If  $N_\varepsilon \gg \varepsilon^{-1} |\log \varepsilon|$  then the line energy becomes negligible, finite energy sequences may only converge weakly and the limit energy is given by the convex envelope  $\int_Q D^{**}(u, B_R) dx$ , where  $u$  may now take values in  $\mathbf{R}$ , see Corollary 11 below for a precise statement.

From a more general mathematical point of view the problem we consider combines two features which have been very extensively studied in the last years. The first one is the interaction of singularly perturbed multiwell energies and higher order regularisations. Beginning with the work of Modica and Mortola [14, 15] a large body of work has concentrated on multiwell energies with a compact set of energy minimizing states and a local regularization given by the Dirichlet integral (see e.g. [13, 9, 7, 5] and the extensive list of references therein).

The more delicate case of a regularizing term which corresponds to the  $H^{1/2}$  norm and which requires a logarithmic rescaling was studied by Alberti, Bouchitté and Seppecher [3]. For nonlocal terms with general anisotropic, but regular, kernels see [2, 1].

The second feature is the interaction of a pinning condition like (2) and Dirichlet-type energies. Since the work of Marchenko and Kruslov [12] and of Cioranescu and Murat [4] this interaction has attracted a lot of attention (see for instance [6] for many further references). Roughly speaking, a general theme of this large body of work is that for well separated obstacles the limiting problem has no constraint like (2) but involves an extra energy contribution of the form  $\int a(x)|u|^2 dx$  where  $a(x)$  can be viewed as a local 'capacity density' (and where the appropriate notion of 'capacity' is related to the Dirichlet-type energy). Of course  $a$  may be singular or degenerate and instead of  $a(x)dx$  one may obtain more general measures which are no longer absolutely continuous with respect to Lebesgue measure.

Our problem combines both features. We show that the contributions of the singular perturbation (leading to a line energy) and of the pinning constraint are essentially additive. The interaction of pinning and the multiwell structure is reflected in the structure of the limiting pinning energy  $\int_Q D_\nu(u, B_R) dx$  which is no longer a quadratic expression in  $u$ .

Compared with most previous work on singularly perturbed multiwell problems our problem involves an important additional difficulty. The set of wells, i.e. the zero set of  $W$  is no longer compact. Hence it is not clear that sequences with finite (rescaled) energy are bounded in  $L^1$ . One crucial ingredient in our analysis is a uniform  $L^2$  estimate (up to translation by integers) for sequences for which the rescaled energy  $E_\varepsilon / |\log \varepsilon|$  is bounded (see Theorems 12 and 13 below).

## 1.1 A quick review of the phase field theory of Koslowski, Cuitiño and Ortiz

Here we briefly discuss the interpretation of the different terms in (1) and the pinning condition. We refer to [10] for a detailed discussion of the model and to [8] for a discussion of the non-dimensional version of the energy (1). The setting is that of continuum crystal elasticity, with small strains.

Consider the displacement field  $U : \mathbf{R}^3 \rightarrow \mathbf{R}^3$  of an infinity elastic crystal. A specific assumption of the model in [10] is that one allows crystal slip only along the single plane  $x_3 = 0$ . Moreover one assumes that only a single slip system is active, i.e. the jump  $[U]$  of the displacement  $U$  across the slip plane is of the form

$$[U](x_1, x_2) = u(x_1, x_2)\mathbf{b}\mathbf{e}, \quad (4)$$

where  $b = |\mathbf{b}|$  is the length of the Burgers vector  $\mathbf{b}$  of the active slip system and  $\mathbf{e} = \mathbf{b}/b$  its direction. We choose coordinates such that  $\mathbf{e}$  is the first coordinate direction. Given  $u$  the associated elastic energy is obtained by minimizing the usual linear elastic energy away from the slip plane, i.e. by minimizing

$$\int_{\mathbf{R}^3 \setminus \{x_3=0\}} \mu |e(U)|^2 + \frac{\lambda}{2} |\text{tr } e(U)|^2 dx_1 dx_2 dx_3, \quad (5)$$

where  $e(U) = \frac{1}{2}(\nabla U + \nabla U^T)$  is the symmetrized displacement gradient. We now suppose that  $u : \mathbf{R}^2 \rightarrow \mathbf{R}$  is periodic with periodic cell  $Q = (-1/2, 1/2)^2$  so that  $u$  can be viewed as a function on the torus  $T^2$ . Then the elastic energy per period is obtained by minimizing

$$\int_{T^2 \times \mathbf{R}} \mu |e(U)|^2 + \frac{\lambda}{2} |\text{tr } e(U)|^2 dx_1 dx_2 dx_3.$$

This minimization can be carried out by considering the Fourier series of  $u$

$$u(x) = \sum_{k \in (2\pi\mathbf{Z})^2} \hat{u}(k)e^{ik \cdot x}, \quad \hat{u}(k) = \int_{T^2} u(x)e^{-ik \cdot x} dx$$

and this yields

$$E_{\text{elastic}}(u) = \frac{\mu b^2}{4} \sum_{k \in (2\pi\mathbf{Z})^2} m_\nu(k) |\hat{u}(k)|^2,$$

where  $m_\nu(k) = \frac{k_2^2}{|k|} + \frac{1}{1-\nu} \frac{k_1^2}{|k|}$  and where  $\nu \in (-1, 1/2)$  is the Poisson's ratio which is given by  $\nu = \frac{\lambda}{2(\lambda+\mu)}$ .

In real space we have

$$E_{\text{elastic}}(u) = \frac{\mu b^2}{2} \int_{T^2} \int_{T^2} K_\nu(x-y) |u(x) - u(y)|^2 dx dy, \quad (6)$$

and the kernel  $K_\nu$  is given by the Fourier series with coefficients  $-\frac{1}{4}m_\nu(k)$ .

Now we turn to the local contribution in  $E_\varepsilon$ . If  $u$  is a (constant) integer then the jump  $[U]$  is a lattice vector and hence the crystal lattice is not perturbed in the immediate neighbourhood of the slip plane. If  $u$  is not an integer there is an additional distorsion of the lattice near the slip plane and in the Peierls-Nabarro theory one models this by an extra energy contribution

$$E_{\text{Peierls}}(u) = \int_{T^2} W(u) dx,$$

where  $W$  is a one-periodic, non-negative functions which vanishes exactly on  $\mathbf{Z}$ . If one fixes a specific form of  $W$ , e.g.  $W(u) = A \text{dist}^2(u, \mathbf{Z})$ , or  $W = \frac{A}{\pi^2} \sin^2(\pi u)$  one can relate the coefficient  $A$  to the shear modulus  $\mu$  and the properties of the crystal lattice by considering very small shear deformations. This line of reasoning leads to the expression  $A = \mu b^2 / (2d)$ , where  $d$  is distance between two neighbouring slip planes.

Dividing the sum of  $E_{\text{Peierls}}$  and  $E_{\text{elastic}}$  by  $\mu b^2 / 2$  we arrive to the energy  $E_\varepsilon$  in (1) with  $W$  given by (3) and  $\varepsilon = d$ . Thus  $\varepsilon$  is proportional to the lattice spacing. If one starts more generally from the situation where  $u$  is periodic with periodic cell  $(-l/2, l/2)^2$  and rescales to the unit cell one finds similarly that  $\varepsilon = d/l$ . Instead of periodic boundary conditions for  $u$  one can also consider other boundary conditions, see also the next subsection.

The regions  $B_{R\varepsilon}^i$  represent obstacles (e.g. inclusions of another material) which restrain slip. The condition (2) represents the limiting case of infinitely strong obstacles which permit no slip at all.

One can also consider obstacles of finite strength where slip is only possible under sufficiently strong loading. In this case one drops the condition (2) and instead adds an additional term

$$E_{\text{obstacle}} = \sum_i \int \lambda_0 \frac{1}{\varepsilon} \psi\left(\frac{x - x_\varepsilon^i}{\varepsilon}\right) |u| dx \quad (7)$$

to the energy (see Section 4.3 of [10], and the Appendix of [8] for further details). In [8] we have shown (in the dilute limit) that the consideration of obstacles of finite strength leads to a  $\Gamma$ -limit which has exactly the same form as for infinitely strong obstacles. The difference is that now the limiting pinning energy density  $D_\nu(a, B_R)$  no longer has quadratic growth in  $a$ , but only grows linear in  $a$ , the limiting slope being related to the effective strength.

Finally one can easily include a forcing term

$$- \int_{T^2} S^\varepsilon u dx$$

in the energy, where  $S^\varepsilon$  is the resolved shear stress. In view of the results in Section 3 a natural scaling assumption on the applied force is

$$\frac{1}{\varepsilon N^\varepsilon} S^\varepsilon \rightarrow S \quad \text{in } L^2(T^2).$$

In this case the corresponding  $\Gamma$ -limit simply contains the additional term  $-\int_{T^2} S u dx$ . If  $S^\varepsilon \ll \varepsilon N^\varepsilon$  then the applied force disappears in the limit. If  $S^\varepsilon \gg \varepsilon N^\varepsilon$  then the applied force dominates in the limit. In the case  $\varepsilon N^\varepsilon \ll |\log \varepsilon|$  there can still be an interesting interaction between the applied force and the line tension term in the limit, at least if the total applied force  $\int_{T^2} S^\varepsilon dx$  vanishes or converges to zero sufficiently fast as  $\varepsilon \rightarrow 0$ .

## 1.2 Possible generalizations

For the rest of this paper we consider the functional (1) with the specific choice  $W(u) = \text{dist}^2(u, \mathbf{Z})$ , the specific periodic kernel (6) discussed in the previous subsection and the hard pinning condition (2). Many of the results can, however, easily be extended to a more general setting. We briefly discuss some possible generalizations, roughly in the order of increasing difficulty.

- *More general local functions  $W$ .* The results can easily be extended to general periodic continuous functions  $W$  which are minimized exactly on  $\mathbf{Z}$ . For the crucial compactness result one can even allow certain nonperiodic  $W$  as long as their minimizers remain discrete and  $W$  does not degenerate too much near  $\pm\infty$ . As discussed in Remark 8 below the limiting energy does not depend on  $W$  but just on the spacing of its zeroes.
- *Other boundary conditions, other kernels  $K$ .* The results can be extended to other boundary conditions (for a sufficiently smooth bounded domain  $\Omega \subset \mathbf{R}^2$ ). The corresponding three-dimensional elastic energy in the cylinder  $\Omega \times \mathbf{R}$  has the same singular behaviour as in the periodic case. For Neumann (natural) boundary conditions one obtains e.g. the expression  $\int_\Omega \int_\Omega K(x, y) |u(x) - u(y)|^2 dx dy$  where the kernel  $K(x, y)$  is smooth and behaves asymptotically for  $y \rightarrow x$  as  $\Gamma_\nu$ , i.e.  $K(x, y) \approx \Gamma_\nu(x - y)$ . One can also consider abstract (sufficiently smooth) kernels  $K(x, y)$  as long as they behave like  $L(x - y)$  for  $y \rightarrow x$ , where  $L$  is a positive function which is homogeneous of degree  $-3$ .
- *General dimensions.* We focus on the case  $n = 2$  because it arises naturally in the dislocation model which motivated this work. The results can, however, easily be extended to general dimensions, using a kernel which behaves like the kernel of the  $H^{1/2}$  norm, i.e.  $K(x, y) \approx |x - y|^{-n-1}$ . In this case the compactness result gives an estimate in  $L^{n/(n-1)}$  (for  $n \geq 2$ ). The case  $n = 1$  was studied by Kurzke [11]. In this case the optimal estimate is in the Orlicz space  $e^L$ .

- *Soft pinning.* As discussed in the previous subsection one can replace the 'hard' pinning condition (2) by a penalty term (7). For the subcritical scaling this is discussed in the appendix of [8] and the argument can be extended to the general setting.
- *Different obstacles and varying obstacle densities.* It is not necessary to assume that the obstacles have all the same size  $R\varepsilon$  (or the same strength in the case of soft obstacles). Instead one can consider obstacles of varying size  $B_i = B(x_i, R_i\varepsilon)$ . Also the density of obstacles need not be constant. In this case we expect that the penalty term due to pinning is of the form  $\Lambda(x)D_\nu(u(x), B_1)$  where  $\Lambda(x)$  represents the local 'capacity density' (appropriately scaled with  $N_\varepsilon$ ). This is briefly discussed in Remark 13 in [8]. Such extensions are well known in the context of competition between pinning and a local Dirichlet energy. To identify the precise assumptions for such a result to hold in the present context, including a suitable weakening of the condition on equipartition, will require some technical work.

### 1.3 Outline

In Section 2 we briefly review some known results, in particular the properties of the periodic kernel  $K_\nu$  and its relation with a similar  $-3$  homogeneous kernel  $\Gamma_\nu$ , the convergence result in the dilute case  $N_\varepsilon \leq C/\varepsilon$  and the definition of the dislocation capacity  $D_\nu(a, B_R)$ , and finally the results of Alberti, Bouchitté and Seppecher for the competition between a two-well energy and the  $H^{1/2}$  norm under logarithmic rescaling, leading to a line tension limit.

In Section 3 we describe our main results. To emphasize the underlying mathematical structure we state them separately for the functional with and without pinning. The next three sections are devoted to the proof of the result in the critical scaling regime. The central compactness estimate is discussed in Section 4 and the upper and lower bound are discussed in Sections 5 and 6, respectively. In the last section we sketch how the result can be extended to the sub- and supercritical scaling. While the argument in the subcritical scaling is a straightforward extension of the results in the dilute case [8] using the compactness estimate, the lower bound in the supercritical case requires some care. We no longer have compactness but we can still show that the oscillation of  $u$  is small on a scale which is large compared to the typical spacing of the pinning sites and this suffices to conclude.

## 2 Review of some known results

### 2.1 Properties of the anisotropic kernel

Here we review some properties of the nonlocal term in the energy, see [10, 8] for further details and proofs. We start from the expression

$$\iint_{T^2 \times T^2} K_\nu(x-y)|u(x)-u(y)|^2 dx dy.$$

for the (suitably normalized) elastic energy introduced above. The Fourier coefficients of the kernel  $K_\nu$  are given by

$$-\frac{1}{4} \left( \frac{k_2^2}{|k|} + \frac{1}{1-\nu} \frac{k_1^2}{|k|} \right),$$

where  $-1 < \nu < \frac{1}{2}$  is the Poisson's ratio. Since the Fourier coefficients of  $K_\nu(t)$  are bounded from above and below by a multiple of  $|k|$  the kernel is equivalent to the  $H^{\frac{1}{2}}$ -kernel, i.e.

$$\frac{1}{2}[u]_{H^{\frac{1}{2}}(T^2)} \leq \iint_{T^2 \times T^2} K_\nu(x-y)|u(x)-u(y)|^2 dx dy \leq [u]_{H^{\frac{1}{2}}(T^2)}, \quad (8)$$

where  $[u]_{H^{\frac{1}{2}}(T^2)}$  denotes the  $H^{\frac{1}{2}}(T^2)$  seminorm defined by

$$[u]_{H^{\frac{1}{2}}(T^2)}^2 = \sum_{k \in (2\pi\mathbf{Z})^2} |k| |\hat{u}(k)|^2. \quad (9)$$

In real space this seminorm can be written as

$$\frac{1}{2}[u]_{H^{\frac{1}{2}}(T^2)}^2 = \iint_{T^2 \times T^2} K(x-y)|u(x) - u(y)|^2 dx dy, \quad (10)$$

where the kernel  $K(t)$  is defined by  $\widehat{K}(k) = -\frac{1}{4}|k|$  and satisfies the following properties:

- i)  $K(t) = O(|t|^{-3})$  as  $|t| \rightarrow 0$ ,
- ii)  $K(t)$  is periodic, i.e. is defined in  $T^2$ .

The kernel  $K_\nu(t)$  also satisfies properties i) and ii). For small  $t$  the term  $K_\nu(t)$  is well approximated by the homogeneous and positive kernel

$$\Gamma_\nu(t) = \frac{1}{8\pi(1-\nu)|t|^3} \left( \nu + 1 - 3\nu \frac{t_2^2}{|t|^2} \right), \quad (11)$$

which is the inverse Fourier transform of  $-\frac{1}{4} \left( \frac{\lambda_2^2}{|\lambda|} + \frac{1-\nu}{1-\nu} \frac{\lambda_1^2}{|\lambda|} \right)$ . The following precise relation between  $K_\nu$  and  $\Gamma_\nu$  can be easily established by the Poisson summation formula.

**Proposition 1** *There exists a constant  $C > 0$  such that*

$$|\Gamma_\nu(t) - K_\nu(t)| \leq C$$

on  $\{t \in \mathbf{R}^2 : |t_i| \leq 3/4\}$ . Moreover  $K_\nu$  is positive.

**Remark 2** By Proposition 1 using the homogeneity of  $\Gamma_\nu$  we deduce that for every  $\delta > 0$

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^3 K_\nu(\varepsilon t) = \Gamma_\nu(t)$$

uniformly on  $\{t \in \mathbf{R}^2 : |t| \leq \delta\}$ .

**Remark 3** All the results proved here and in [8] are still true for a more general positive kernel  $\mathcal{K}(t)$  equivalent to the  $H^{\frac{1}{2}}$  kernel satisfying (i) and Proposition 1,  $\Gamma$  being the homogeneous function of degree  $-3$  defined by  $\Gamma(t) = \lim_{\varepsilon \rightarrow 0} \varepsilon^3 \mathcal{K}(\varepsilon t)$ .

Since  $[\cdot]_{H^{\frac{1}{2}}}$  is a trace seminorm we can deduce a Poincaré type inequality for functions in  $H^{\frac{1}{2}}(T^2)$  (see [8], Proposition 3).

**Proposition 4** *There exists a constant  $C_0$  such that for every  $u \in H^{\frac{1}{2}}(T^2)$ , with  $u = 0$  on  $E \subseteq T^2$ , we have*

$$\int_{T^2} |u|^2 dx \leq C_0 \left( 1 + \frac{1}{\text{Cap}(E \times \{0\})} \right) [u]_{H^{\frac{1}{2}}(T^2)}^2, \quad (12)$$

where  $\text{Cap}(E \times \{0\})$  denote the harmonic capacity of  $E \times \{0\}$  as a subset of  $\mathbf{R}^3$ .

**Remark 5** Given an arbitrary  $H^{\frac{1}{2}}(Q)$  function we can extend it by reflection to a periodic function on the square of size 2,  $Q_2$ , and applying the above inequality we get that there exists a constant  $C_1$  such that

$$\int_Q |u|^2 dx \leq C_1 \left( 1 + \frac{1}{\text{Cap}(E \times \{0\})} \right) \iint_{Q \times Q} \frac{|u(x) - u(y)|^2}{|x - y|^3} dx dy. \quad (13)$$

## 2.2 Subcritical density of obstacles and the dislocation capacity

In [8] we studied the  $\Gamma$ -convergence of the variational model described in the introduction for the dilute case, i.e. in the regime  $\varepsilon N_\varepsilon \approx 1$  (we will see in Corollary 11 below that the results actually hold as long as  $\varepsilon N_\varepsilon \ll |\log \varepsilon|$ ). We will always assume that the obstacles  $B_{R\varepsilon}^i$  in (2) are uniformly distributed and well separated in the following sense. For every subset  $E$  of  $(-\frac{1}{2}, \frac{1}{2})^2$  we denote by  $\mathcal{I}_\varepsilon(E) := \{i \in \mathcal{I}_\varepsilon : x_\varepsilon^i \in E\}$  and we require the following.

- (*Uniform distribution*) There exists a constant  $L > 0$  such that

$$|\#\mathcal{I}_\varepsilon(Q') - N_\varepsilon|Q'| \leq L \quad (14)$$

for every open square  $Q' \subset (-\frac{1}{2}, \frac{1}{2})^2$ ;

- (*Separation*) There exists  $\beta < 1$  such that

$$\text{dist}(x_\varepsilon^i, x_\varepsilon^j) > 6\varepsilon^\beta \quad (15)$$

for every  $i, j \in \mathcal{I}_\varepsilon$ ,  $i \neq j$ , and for every  $\varepsilon \in (0, \varepsilon_0)$ ;

In [8] we proved that there exists a function defined on integers and denoted by  $D_\nu(a, B_R)$  such that the following result holds.

**Theorem 6** *Assume  $\varepsilon N_\varepsilon \rightarrow \Lambda$  and that the discs  $B_{R\varepsilon}^i$  are uniformly distributed and well separated. Then the functional  $\mathcal{F}_\varepsilon(u) := E_\varepsilon(u)/N_\varepsilon\varepsilon$ , i.e.*

$$\mathcal{F}_\varepsilon(u) = \begin{cases} \frac{1}{N_\varepsilon\varepsilon^2} \int_{T^2} \text{dist}^2(u, \mathbf{Z}) dx + \frac{1}{N_\varepsilon\varepsilon} \iint_{T^2 \times T^2} K_\nu(x-y) |u(x) - u(y)|^2 dx dy & \text{if } u \in H^{\frac{1}{2}}(T^2), \\ & u = 0 \text{ on } \bigcup_i B_{R\varepsilon}^i \\ +\infty & \text{otherwise,} \end{cases}$$

$\Gamma$ -converges, with respect to the strong  $L^2$  topology, to the functional

$$\mathcal{F}(u) = \begin{cases} D_\nu(u, B_R) & \text{if } u = \text{const.} \in \mathbf{Z}, \\ +\infty & \text{otherwise.} \end{cases} \quad (16)$$

The limit  $D_\nu(u, B_R)$  can be characterized by means of the following cell problem formula

$$D_\nu(a, B_R) := \inf \left\{ \int_{\mathbf{R}^2} \text{dist}^2(\zeta, \mathbf{Z}) dx + \iint_{\mathbf{R}^2 \times \mathbf{R}^2} \Gamma_\nu(x-y) |\zeta(x) - \zeta(y)|^2 dx dy : \right. \quad (17)$$

$$\left. \zeta = a \text{ on } B_R, \zeta \in L^4(\mathbf{R}^2) \right\}$$

More generally for every integer  $a$  and for every open bounded set  $\Omega$  we can define a set function that we call the  $H^{\frac{1}{2}}$ -dislocation capacity of an open set  $E$  with respect to  $\Omega$  at the integer level  $a \in \mathbf{Z}$ , as follows

$$D_\nu(a, E, \Omega) := \inf \left\{ \int_{\mathbf{R}^2} \text{dist}^2(\zeta, \mathbf{Z}) dx + \iint_{\mathbf{R}^2 \times \mathbf{R}^2} \Gamma_\nu(x-y) |\zeta(x) - \zeta(y)|^2 dx dy : \right. \quad (18)$$

$$\left. \zeta = a \text{ on } E, \zeta = 0 \text{ on } \mathbf{R}^2 \setminus \Omega \right\}$$

where  $E$  is an open subset of  $\Omega$ . In (17) and (18) the infimum is attained and the minimum is called the  $H^{\frac{1}{2}}$ -dislocation capacity potential of  $E$ .

In [8] we proved that  $D_\nu(a, \cdot)$  (and  $D_\nu(a, \cdot, \Omega)$ ) is actually a Choquet capacity and moreover it is quadratic in  $a$ , as  $a$  goes to infinity. More precisely, for every bounded open set  $E$  there exist two constants,  $C_1$  and  $C_2$ , such that

$$C_1 a^2 \leq D_\nu(a, E) \leq C_1(a^2 + 2a^{3/2}) + C_2 a \quad (19)$$

(see [8], Proposition 8). Another fact which is crucial in using  $D_\nu(a, B_R)$  as a cell-problem formula in the study of the asymptotic behaviour of  $\mathcal{F}_\varepsilon$  is the following convergence property

$$\lim_{T \rightarrow \infty} D_\nu(a, E, B_T) = D_\nu(a, E). \quad (20)$$

### 2.3 The $H^{1/2}$ regularisation and logarithmic rescaling

Alberti, Bouchitté and Seppecher studied the functional  $E_\varepsilon$  (without the pinning condition) for a two-well potential  $W$  and the  $H^{1/2}$  nonlocal energy.

**Theorem 7 ([3])** *The functional*

$$J_\varepsilon(u) = \begin{cases} \frac{1}{|\log \varepsilon| \varepsilon} \int_Q \text{dist}^2(u, \{0, 1\}) dx + \frac{1}{|\log \varepsilon|} \iint_{Q \times Q} \frac{|u(x) - u(y)|^2}{|x - y|^3} dx dy & \text{if } u \in H^{\frac{1}{2}}(Q), \\ +\infty & \text{otherwise.} \end{cases} \quad (21)$$

$\Gamma$ -converges to the functional

$$J(u) = \begin{cases} 4\text{Per}_Q(\{u = 0\}) & \text{if } u \in BV(Q, \{0, 1\}), \\ +\infty & \text{otherwise,} \end{cases} \quad (22)$$

where  $\text{Per}_Q(\{u = 0\})$  denotes the perimeter of the set  $\{u = 0\}$  relative to the set  $Q$ .

**Remark 8** We have stated the result for the special two-well energy  $\text{dist}^2(u, \{0, 1\})$  to emphasize the similarities and differences with the choice (3). The result in [3] is proved for a general local energy  $W$  with  $W(0) = W(1) = 0$ ,  $W > 0$  otherwise and  $W(u) \geq c|u|^2 - C$ . Interestingly, and in contrast to the situation with regularizing energy of Dirichlet type [14, 15, 13], the  $\Gamma$ -limit does not depend on the shape of  $W$ , but just on the position of its zeros. This is an effect of the logarithmic rescaling. In fact in the proof of the upper bound in the verification of the  $\Gamma$ -limit one has a large amount of freedom. The precise choice of the transition profile by which one approximates a jump does not matter, as long as the length scale of the transition is chosen correctly. We will also exploit this fact in Section 6.

Comparing the above result with Theorem 6 it is natural to conjecture that the critical scaling regime for the number of obstacles  $N_\varepsilon$  which leads to an interaction of the line energy in the result above and the pinning energy discussed earlier is given by  $N_\varepsilon \approx \varepsilon^{-1} |\log \varepsilon|$ . This is what we will establish in the next section. As a byproduct we also obtain the limiting energy functionals in the subcritical regime  $N_\varepsilon \ll \varepsilon^{-1} |\log \varepsilon|$  and the supercritical regime  $N_\varepsilon \gg \varepsilon^{-1} |\log \varepsilon|$ .

## 3 Main results

We establish compactness and  $\Gamma$ -convergence results for the energy functionals  $E_\varepsilon(u)$ , both with and without the pinning condition. We first describe the behaviour of the functionals  $I_\varepsilon(u) = E_\varepsilon(u)/|\log \varepsilon|$ , thus extending the results of [3] to energies with infinitely many wells (and to an anisotropic kernel).

**Theorem 9** *The functional*

$$I_\varepsilon(u) = \begin{cases} \frac{1}{|\log \varepsilon| \varepsilon} \int_{T^2} \text{dist}^2(u, \mathbf{Z}) dx + \frac{1}{|\log \varepsilon|} \iint_{T^2 \times T^2} K_\nu(x-y) |u(x) - u(y)|^2 dx dy & \text{if } u \in H^{\frac{1}{2}}(T^2), \\ +\infty & \text{otherwise,} \end{cases} \quad (23)$$

$\Gamma$ -converges to the functional

$$I(u) = \begin{cases} \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1 & \text{if } u \in BV(T^2, \mathbf{Z}), \\ +\infty & \text{otherwise,} \end{cases} \quad (24)$$

where  $n$  denotes the normal on the jump set  $S_u$  of  $u$  and the anisotropic line energy density  $\gamma(n)$  is defined, for any  $n \in S^1$ , by

$$\gamma(n) := 2 \int_{x \cdot n = 1} \Gamma_\nu(x) d\mathcal{H}^1 = 2 \lim_{\delta \rightarrow 0} \delta^2 \int_{x \cdot n = \delta} K_\nu(x) d\mathcal{H}^1, \quad (25)$$

where  $\Gamma_\nu$  is given by (11). More precisely

- i) Every sequence  $\{u_\varepsilon\}$  such that  $\sup_\varepsilon I_\varepsilon(u_\varepsilon) < \infty$  is bounded in  $L^2(T^2)$ , up to a translation, and is pre-compact in  $L^q(T^2)$  for every  $q < 2$ . Every cluster point  $u$  of the translated sequence belongs to  $BV(T^2, \mathbf{Z})$  and satisfies

$$\int_{T^2} |Du| \leq C \liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon).$$

- ii) Every sequence  $\{u_\varepsilon\}$  strongly converging in  $L^1(T^2)$  to some function  $u$  satisfies

$$\liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) \geq I(u).$$

- iii) For every  $u \in BV(T^2, \mathbf{Z})$  there exists a sequence  $\{u_\varepsilon\}$  strongly converging in  $L^q(T^2)$ , for all  $q < 2$ , to  $u$  such that

$$\lim_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) = I(u).$$

The proof of the result above will be obtained as a consequence of a compactness theorem (Theorem 12 in Section 4) a lower bound given in Section 5 (Theorem 17), and an upper bound proved in Section 6 (Theorem 24).

The asymptotic analysis for the functional with the pinning condition in the critical scaling  $N_\varepsilon \approx \varepsilon^{-1} |\log \varepsilon|$  is summarized in the following theorem.

**Theorem 10** *Assume  $N_\varepsilon \varepsilon / |\log \varepsilon| \rightarrow \Lambda$ ,  $0 < \Lambda < \infty$ , and that the discs  $B_{R\varepsilon}^i$  are uniformly distributed and well separated. Then the functional*

$$F_\varepsilon(u) = \begin{cases} \frac{1}{|\log \varepsilon| \varepsilon} \int_{T^2} \text{dist}^2(u, \mathbf{Z}) dx + \frac{1}{|\log \varepsilon|} \iint_{T^2 \times T^2} K_\nu(x-y) |u(x) - u(y)|^2 dx dy & \text{if } u \in H^{\frac{1}{2}}(T^2), \text{ and } u = 0 \text{ on } \bigcup_i B_{R\varepsilon}^i \\ +\infty & \text{otherwise,} \end{cases} \quad (26)$$

$\Gamma$ -converges to the functional

$$F(u) = \begin{cases} \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1 + \Lambda \int_{T^2} D_\nu(u, B_R) dx & \text{if } u \in BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z}), \\ +\infty & \text{otherwise.} \end{cases} \quad (27)$$

More precisely

i) Every sequence  $\{u_\varepsilon\}$  such that  $\sup_\varepsilon F_\varepsilon(u_\varepsilon) < \infty$  is bounded in  $L^2(T^2)$  and is pre-compact in  $L^q(T^2)$ , for every  $q < 2$ . Every cluster point  $u$  belongs to  $BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$  and satisfies

$$\int_{T^2} |Du| \leq C \liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon).$$

ii) Every sequence  $\{u_\varepsilon\}$  strongly converging in  $L^1(T^2)$  to some function  $u$  satisfies

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) \geq F(u).$$

iii) For every  $u \in BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$  there exists a sequence  $\{u_\varepsilon\}$  strongly converging in  $L^q(T^2)$ , for any  $q < 2$ , to  $u$  such that

$$\lim_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) = F(u).$$

Theorem 10 follows from the compactness result obtained in Section 4 (Proposition 16) and the lower and the upper bounds obtained in Section 5 (Theorem 21) and Section 6 (Theorem 27), respectively.

Finally we extend the result to the subcritical and supercritical scaling, which correspond to  $\Lambda = 0$  and  $\Lambda = \infty$ , respectively. If we consider the rescaling  $E_\varepsilon/(N_\varepsilon\varepsilon)$ , then formally the limit is given by dividing (27) by  $\Lambda$ . Thus for  $\Lambda = 0$  we expect the line energy to dominate, leading to the constraint  $u = \text{const}$ , while for  $\Lambda = \infty$  the line energy becomes negligible so we can no longer expect compactness in  $L^1$  and the limiting energy density functional has to be replaced by its relaxation, given by the convex hull  $D_\nu^{**}(u, B_R)$ . More precisely

$$\begin{aligned} u \mapsto D_\nu^{**}(u, B_R) \quad &\text{is the convex hull of the function } \tilde{D}, \text{ where} \\ \tilde{D}(u) = D_\nu(u, B_R), \quad &\text{if } u \in \mathbf{Z}, \quad \tilde{D}(u) = \infty \quad \text{otherwise.} \end{aligned} \quad (28)$$

**Corollary 11** *Assume that the discs  $B_{R\varepsilon}^i$  are uniformly distributed and well separated. Consider the functional*

$$\mathcal{F}_\varepsilon(u) = \begin{cases} \frac{1}{\varepsilon^2 N_\varepsilon} \int_{T^2} \text{dist}^2(u, \mathbf{Z}) \, dx + \frac{1}{\varepsilon N_\varepsilon} \iint_{T^2 \times T^2} K_\nu(x-y) |u(x) - u(y)|^2 \, dx \, dy \\ \quad \text{if } u \in H^{\frac{1}{2}}(T^2), \text{ and } u = 0 \text{ on } \bigcup_i B_{R\varepsilon}^i, \\ +\infty \quad \text{otherwise.} \end{cases} \quad (29)$$

(i) *If  $N_\varepsilon \rightarrow \infty$  and  $\varepsilon N_\varepsilon / |\log \varepsilon| \rightarrow 0$  then  $\mathcal{F}_\varepsilon$   $\Gamma$ -converges (with respect to the strong  $L^2$  topology) to the functional*

$$\mathcal{F}(u) = \begin{cases} D_\nu(a, B_R) \, dx & \text{if } u \equiv a, \, a \in \mathbf{Z}, \\ +\infty & \text{otherwise.} \end{cases} \quad (30)$$

(ii) *If  $\varepsilon N_\varepsilon / |\log \varepsilon| \rightarrow \infty$  then  $\mathcal{F}_\varepsilon$   $\Gamma$ -converges (with respect to the weak  $L^2$  topology) to the functional*

$$\mathcal{F}(u) = \int_{T^2} D_\nu^{**}(u, B_R) \, dx. \quad (31)$$

The proof of Corollary 11 follows closely that of Theorems 9 and 10. In Section 7 we sketch the necessary modifications in the argument. The main difficulty is that in the supercritical case one no longer has  $L^1$  compactness. Nonetheless we will show that on a scale which is large compared to the typical particle separation  $1/\sqrt{N_\varepsilon}$  the  $L^2$  oscillation of a finite energy sequence  $u_\varepsilon$  can still be controlled. This will allow us to locally estimate the energy from below essentially in the same way as in the presence of strong convergence.

## 4 Compactness

In this section we establish the following compactness result.

**Theorem 12** *Let  $\{u_\varepsilon\}$  be such that*

$$\liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) < +\infty,$$

*then there exists a sequence  $\{a_\varepsilon\}$  such that the sequence  $\{u_\varepsilon - a_\varepsilon\}$  is bounded in  $L^2$  and relatively compact in  $L^q(Q)$ , for every  $q < 2$ . Every cluster point  $u$  of  $\{u_\varepsilon - a_\varepsilon\}$  belongs to  $BV(T^2, \mathbf{Z})$  and satisfies*

$$\int_Q |Du| \leq C \liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon). \quad (32)$$

The main difference with the result of Alberti, Bouchitté and Seppecher [3] is that the local part of the energy  $\text{dist}^2(u, \mathbf{Z})$  is not coercive. Hence the crucial point of the proof consists in the derivation of a uniform  $L^2$  bound on  $u_\varepsilon$  (up to a translation). Pointwise convergence can then be deduced for [3] by a truncation argument.

### 4.1 The $L^2$ -bound

**Theorem 13** *Let  $\{u_\varepsilon\}$  such that*

$$I_\varepsilon(u_\varepsilon) \leq M \quad \forall \varepsilon > 0 \quad (33)$$

*and assume that  $|\{u_\varepsilon > 0\}| > \frac{1}{2}|Q|$  and  $|\{u_\varepsilon < 1\}| > \frac{1}{2}|Q|$ . Let  $Q' \subset\subset Q$  be a square centered in zero. Then there exists a constant  $C(Q')$  such that*

$$\int_{Q'} |u_\varepsilon|^2 dx \leq C(Q').$$

#### Remarks

1. The bound is optimal, since the embedding of  $BV$  into  $L^2$  is optimal. Indeed if  $I_\varepsilon(u_\varepsilon) \leq M$  would imply a uniform  $L^p_{\text{loc}}$  bound for  $p > 2$  then, by Theorem 9, every function in  $BV(Q, \mathbf{Z})$  would be  $L^p_{\text{loc}}(Q)$ , which is false.
2. If one considers the analogue of the functional  $I_\varepsilon$  in  $n$  dimensions ( $n \geq 2$ ), with kernel  $K(z) \approx |z|^{-(n+1)}$  the argument below gives an  $L^{\frac{n}{n-1}}_{\text{loc}}$  bound, which is again optimal. For  $n = 1$  Kurzke [11] showed that the optimal bound corresponds to the Orlicz space  $e^L$ .

The idea of the proof of Theorem 13 is simple. From Theorem 9 we expect that  $I_\varepsilon(u_\varepsilon)$  behaves asymptotically similar to the  $BV$  seminorm. Comparing the super-level sets  $\{u_\varepsilon > k\}$  and  $\{u_\varepsilon > k-1\}$  we thus expect that the perimeter of  $\{u_\varepsilon > k\}$  is controlled by  $I_\varepsilon(T_{k-1}u_\varepsilon)$ , where  $T_{k-1}u_\varepsilon$  is given by  $u_\varepsilon \vee (k-1) \wedge k$ . In combination with the isoperimetric inequality we would get  $|\{u_\varepsilon > k\}|^{\frac{1}{2}} \leq CI_\varepsilon(T_{k-1}u_\varepsilon)$ , and from this the assertion would already follow (see (49) and (50) below).

Unfortunately  $I_\varepsilon$  does not really control the perimeter of the level sets uniformly in  $\varepsilon$ . Instead we will establish directly a bound of the form

$$|A_k^\varepsilon|^{\frac{1}{2}} = |Q' \cap \{u_\varepsilon > k + 1 - \sigma\}|^{\frac{1}{2}} \leq CI_\varepsilon(T_k u_\varepsilon), \quad (34)$$

for a fixed  $\sigma \in (0, \frac{1}{4})$ . To prove this we first replace the singular kernel by a regular kernel and drop the logarithmic rescaling. Then we bound the singular kernel  $\frac{1}{|t|^\alpha}$  from below by a dyadic sum of scaled regular kernels and with a covering argument we get an estimate like (34) at least if  $|A_k^\varepsilon|$  is not too small, see (46) below.

If  $|A_k^\varepsilon|$  is very small then a scaling argument shows that the term  $\text{dist}^2(u_\varepsilon, \mathbf{Z})$  becomes negligible and we directly use the embedding of  $H^{\frac{1}{2}}$  in  $L^4$  and conclude after a short calculation.

We begin with the relevant estimates for the regular kernels.

**Proposition 14** Fix  $\varphi \in C_c^\infty(B_1(0))$ , with  $\int \varphi(x) dx = 1$  and  $\varphi > 0$  in  $B_{\frac{1}{2}}(0)$ . For every  $\lambda \in (0, 1)$  there exists a constant  $c(\lambda)$  such that for every subset  $A$  of  $Q$  with  $\lambda \leq |A| \leq \frac{1+\lambda}{2}$  we have

$$\gamma_\delta(A, Q) := \frac{1}{\delta} \int_A \int_{Q \setminus A} \varphi_\delta(x - x') dx dx' \geq c(\lambda) \quad \forall \delta \in (0, 1), \quad (35)$$

where  $\varphi_\delta(x) = \frac{1}{\delta^2} \varphi\left(\frac{x}{\delta}\right)$ .

**Proof.** Fix  $\lambda$ . Assume by contradiction that there exists a sequence of sets  $\{A_k\}$  such that

$$\lambda \leq |A_k| \leq \frac{1+\lambda}{2} \quad (36)$$

and a sequence  $\{\delta_k\}$ , such that

$$\lim_{k \rightarrow \infty} \gamma_{\delta_k}(A_k, Q) = 0. \quad (37)$$

We may assume that  $\delta_k$  converges to some  $\delta$  and the characteristic function  $\chi_{A_k}(x)$  of  $A_k$  converges to  $\theta(x)$  weak\* in  $L^\infty$  as  $k \rightarrow \infty$ . If  $\delta \neq 0$ , then from (37) we get

$$\iint_{Q \times Q} \theta(x)(1 - \theta(x')) \varphi_\delta(x - x') dx dx' = 0.$$

This implies either  $\theta = 0$  a.e. in  $Q$  or  $\theta = 1$  a.e. in  $Q$ , which is impossible, since (36) implies  $\lambda \leq \int_Q \theta(x) dx \leq \frac{1+\lambda}{2}$ . If  $\delta = 0$ , we can rewrite  $\gamma_{\delta_k}(A_k)$  as follows

$$\begin{aligned} & \frac{1}{\delta_k} \int_{A_k} \int_{Q \setminus A_k} \varphi_{\delta_k}(x - x') dx dx' = \frac{1}{2\delta_k} \int_Q \int_Q \varphi_{\delta_k}(x - x') |\chi_{A_k}(x) - \chi_{A_k}(x')|^2 dx dx' \\ & = \frac{1}{2\delta_k} \int_Q \int_Q \varphi_{\delta_k}(x - x') |\chi_{A_k}(x) - \chi_{A_k}(x')|^2 dx dx' + \frac{1}{2\delta_k} \int_Q \text{dist}^2(\chi_{A_k}, \{0, 1\}) dx =: \frac{1}{2} \Phi_{\delta_k}(\chi_{A_k}) \end{aligned}$$

The asymptotics of the functional  $\Phi_\delta$  has been studied by Alberti and Bellettini in [1]. By their compactness result we get that there exists a set  $A$  with finite perimeter such that, up to a subsequence, the sequence  $\chi_{A_k}$  strongly converges to  $\chi_A$  in  $L^1$ . Moreover applying [1], Theorem 1.4, we obtain that there exists a positive constant  $C$  such that

$$\liminf_{k \rightarrow \infty} \gamma_{\delta_k}(A_k, Q) \geq C \text{Per}_Q(A).$$

By (37) we get  $\text{Per}_Q(A) = 0$ , but this again contradicts (36).  $\square$

**Lemma 15** Let  $Q' \subset \subset Q$ . There exist two constants  $C_1$  and  $C_2$ , depending on  $Q'$ , such that for every  $A \subseteq Q$  and  $B \subseteq Q$  with  $A \cap B = \emptyset$  disjoint the following inequality holds (with  $E = Q \setminus (A \cup B)$ )

$$\int_{A \cap Q'} \int_{B \cap Q'} \frac{1}{|x - y|^3} dx dy \geq C_1 |A \cap Q'|^{\frac{1}{2}} \log \frac{C_2 |A \cap Q'|}{|E|}. \quad (38)$$

**Proof.** We first prove the equivalent of (35) for the singular kernel  $\frac{1}{|t|^3}$ , i.e. we prove that if  $A \subseteq Q$  satisfies

$$\lambda \leq |A| \leq \frac{\lambda + 1}{2},$$

then there exists a constant  $C(\lambda)$  such that

$$\int_A \int_B \frac{1}{|x - y|^3} dx dy \geq C(\lambda) \log \frac{c(\lambda)}{2|E|}, \quad (39)$$

$c(\lambda)$  being the constant given by Lemma 14. Clearly we may assume that  $|E| = |Q \setminus (A \cup B)| < \frac{c(\lambda)}{2}$ . Since  $A \cup B \cup E = Q$ , by Lemma 14 it is easy to see that

$$\frac{1}{\delta} \int_A \int_B \varphi_\delta(x-y) dx dy \geq c(\lambda) - \frac{|E|}{\delta} \geq \frac{c(\lambda)}{2} \quad \text{as long as } \frac{2|E|}{c(\lambda)} < \delta < 1. \quad (40)$$

We will get (39) by estimating the singular kernel with a dyadic sum of regular kernels. Take  $m \in \mathbb{N}$  such that  $2^{-m} \leq \frac{2|E|}{c(\lambda)} \leq 2^{-m+1}$ , hence  $m \geq \log(c(\lambda)/2|E|)/\log 2$ . Then take  $\delta_i = 2^{-i}$  for  $i = 0, \dots, m$ . By the definition of  $\varphi_\delta$  and taking into account that  $\varphi_\delta(x-y) = 0$  if  $|x-y| > \delta$ , we have

$$\frac{1}{|x-y|^3} \geq \frac{1}{\delta \sup \varphi} \varphi_\delta(x-y) \quad \forall x, y \in Q.$$

More generally we have that there exists a constant  $C$ , depending on  $\varphi$ , such that

$$\frac{1}{|x-y|^3} \geq C \sum_{i=0}^m \frac{1}{\delta_i} \varphi_{\delta_i}(x-y) \quad (41)$$

for all  $x, y \in Q$ . Indeed

$$\sum_{i=0}^m \frac{1}{\delta_i} \varphi_{\delta_i}(x-y) \leq \sup \varphi \sum_{i=0}^m \frac{1}{\delta_i^3} \leq 2^4 \sup \varphi \frac{1}{\delta_{m-1}^3},$$

and this, by the definition of  $\delta_{m-1}$ , implies immediately that (41) holds if  $|x-y| < 2|E|/c(\lambda)$ . On the other hand since  $\text{supp } \varphi \subseteq B_1(0)$ , for any  $i_0 \in \{0, 1, \dots, m-1\}$  and  $\delta_{i_0+1} \leq |x-y| \leq \delta_{i_0}$ , we get

$$\sum_{i=0}^m \frac{1}{\delta_i} \varphi_{\delta_i}(x-y) \leq \sup \varphi \sum_{i=0}^{i_0+1} \frac{1}{\delta_i^3} \leq 2^4 (\sup \varphi) \frac{1}{\delta_{i_0}^3}$$

which gives (41) for every  $x, y \in Q$ . Thus (39) follows by (41) and (40).

The proof of the result can be now obtained with a covering argument. Indeed by scaling (39) we get that for every  $r > 0$  and for every  $A \subseteq Q$  such that

$$\lambda \leq \frac{|A \cap Q_r|}{|Q_r|} \leq \frac{\lambda+1}{2},$$

we have

$$\int_{A \cap Q_r} \int_{B \cap Q_r} \frac{1}{|x-y|^3} dx dy \geq Cr \log \frac{c(\lambda)r^2}{2|E \cap Q_r|}, \quad (42)$$

where  $Q_r$  denotes a square of size  $r$ . Fix  $Q' \subset\subset Q$ , let  $d = \text{dist}(\partial Q, Q')$ . We claim that there exists a constant  $\lambda = \lambda(d)$  such that for every  $x \in A \cap Q'$  there exists a square  $Q_{r_x}(x)$  centered in  $x$  and of size  $r_x$  such that

$$\lambda \leq \frac{|A \cap Q' \cap Q_{r_x}(x)|}{|Q_{r_x}(x)|} \leq \frac{1+\lambda}{2}. \quad (43)$$

Indeed on the one hand a.e.  $x \in A \cap Q'$  has density 1 and on the other hand for every  $r$

$$\min_{r: Q_r(x) \subset\subset Q} \frac{|A \cap Q' \cap Q_r(x)|}{|Q_r(x)|} \leq \max_{x \in Q'} \min_{r: Q_r(x) \subset\subset Q} \frac{|Q' \cap Q_r(x)|}{|Q_r(x)|} =: \lambda(d).$$

By the continuity of  $\frac{|A \cap Q' \cap Q_r(x)|}{|Q_r(x)|}$  with respect to  $r$  we get (43). By the Besicovitch Covering Theorem we obtain a family of disjoint squares  $Q_{r_i}(x_i)$ ,  $i \in I$ , satisfying (43) such that

$$\sum_{i \in I} r_i^2 = \sum_{i \in I} |Q_{r_i}(x_i)| \geq \tilde{c} |A \cap Q'|,$$

where  $\tilde{c}$  is a universal constant. From this family of disjoint squares we can extract a family of indices  $J$  such that

$$\frac{|E \cap Q_{r_i}(x_i)|}{r_i^2} < \frac{2}{\tilde{c}} \frac{|E|}{|A \cap Q'|} \quad \forall i \in J. \quad (44)$$

We have

$$\sum_{i \in J} r_i^2 = \sum_{i \in I} r_i^2 - \sum_{i \notin J} r_i^2 > \tilde{c}|A \cap Q'| - \frac{\tilde{c}}{2} \sum_{i \notin J} \frac{|E \cap Q_{r_i}(x_i)|}{|E|} |A \cap Q'| > \frac{\tilde{c}}{2} |A \cap Q'|. \quad (45)$$

Finally by (42) and (44) we get

$$\begin{aligned} \int_{A \cap Q'} \int_{B \cap Q'} \frac{1}{|x-y|^3} dx dy &\geq \sum_{i \in J} \int_{A \cap Q_{r_i}(x_i)} \int_{B \cap Q_{r_i}(x_i)} \frac{1}{|x-y|^3} dx dy \\ &\geq \sum_{i \in J} C r_i \log \frac{c(\lambda) r_i^2}{2|E \cap Q_{r_i}(x_i)|} \geq C \left( \sum_i r_i^2 \right)^{\frac{1}{2}} \log \frac{c(\lambda) \tilde{c} |A \cap Q'|}{4|E|}. \end{aligned}$$

which concludes the proof together with (45).  $\square$

**Proof of Theorem 13.** Fix  $\sigma \in (0, \frac{1}{4})$  and  $k \in \mathbf{Z}$ . Denote by  $A_k^\varepsilon$  and  $B_k^\varepsilon$  the following two super and sub-level sets

$$A_k^\varepsilon = \{x \in Q' : u_\varepsilon > k + 1 - \sigma\} \quad \text{and} \quad B_k^\varepsilon = \{x \in Q' : u_\varepsilon < k + \sigma\}.$$

We will give the proof in several steps. Assume that  $u_\varepsilon > 0$ . The idea is to get an estimate for the super-level sets of the following type

$$\sum_{k=0}^{\infty} |A_k^\varepsilon|^{\frac{1}{2}} \leq C,$$

in order to deduce the  $L^2$  estimate.

*Step 1.* Fix  $\alpha < 1$ , then there exists a constant  $C$  depending on  $\alpha$  and  $Q'$  such that

$$\int_{A_k^\varepsilon} \int_{B_k^\varepsilon} K_\nu(x-y) dx dy \geq C |\log \varepsilon| |A_k^\varepsilon|^{\frac{1}{2}} \quad (46)$$

whenever  $|A_k^\varepsilon|^{\frac{1}{2}} > \varepsilon^\alpha$ .

Denote by  $E_k^\varepsilon$  the set  $\{x \in Q' : k + \sigma \leq u_\varepsilon \leq k + 1 - \sigma\}$ . By (33) we deduce that  $|E_k^\varepsilon| \leq \frac{M}{\sigma^2} \varepsilon |\log \varepsilon|$ . Then we may assume  $|E_k^\varepsilon| \leq \varepsilon^\alpha$ . Thus, by Lemma 15 and the fact that  $K_\nu(t) \leq C1/|t|^3$ , we get

$$\int_{A_k^\varepsilon} \int_{B_k^\varepsilon} K_\nu(x-y) dx dy \geq C |A_k^\varepsilon|^{\frac{1}{2}} \log \frac{C_2 \varepsilon^{2\alpha}}{\varepsilon^\alpha}$$

which gives (46).

*Step 2.* Let  $\alpha$  be as in Step 1 and assume  $|A_k^\varepsilon|^{\frac{1}{2}} \leq \varepsilon^\alpha$ . Fix  $\gamma \in (1, 2)$ . Then there exists a positive constant  $C$  such that

$$|A_k^\varepsilon|^{\frac{1}{2}} \leq \frac{C}{(k - \sigma)^\gamma}. \quad (47)$$

By the Sobolev inequality and the fact that  $|\{u_\varepsilon < 1\}| \geq \frac{1}{2}|Q|$ , we have

$$\|(u_\varepsilon - 1)_+\|_{L^4(Q)} \leq C [(u_\varepsilon - 1)_+]_{H^{\frac{1}{2}}} \leq C (|\log \varepsilon| I_\varepsilon(u_\varepsilon))^{\frac{1}{2}}.$$

Thus by Hölder's inequality we get

$$\begin{aligned} |A_k^\varepsilon|(k - \sigma) &\leq \int_{A_k^\varepsilon} |(u_\varepsilon - 1)_+| dx \\ &\leq |A_k^\varepsilon|^{\frac{3}{4}} \|(u_\varepsilon - 1)_+\|_{L^4(Q)} \leq C |A_k^\varepsilon|^{\frac{3}{4}} (|\log \varepsilon| I_\varepsilon(u_\varepsilon))^{\frac{1}{2}}. \end{aligned}$$

Hence

$$|A_k^\varepsilon|^{\frac{1}{4}} \leq C \frac{1}{(k - \sigma)} (|\log \varepsilon| I_\varepsilon(u_\varepsilon))^{\frac{1}{2}}.$$

Since  $|A_k^\varepsilon|^{\frac{1}{2}} \leq \varepsilon^\alpha$ , this implies that

$$|A_k^\varepsilon|^{\frac{1}{2\gamma}} \leq C \frac{1}{(k - \sigma)} (I_\varepsilon(u_\varepsilon))^{\frac{1}{2}}$$

and (47) follows by raising the last inequality to the power  $\gamma$ .

*Step 3.* There exists a positive constant  $C$  such that

$$\sum_{k=0}^{\infty} |A_k^\varepsilon|^{\frac{1}{2}} \leq C. \quad (48)$$

This is a consequence of Step 1 and Step 2. Indeed if  $|A_k^\varepsilon|^{\frac{1}{2}} > \varepsilon^\alpha$ , then we apply Step 1 and we get

$$|A_k^\varepsilon|^{\frac{1}{2}} \leq C I_\varepsilon(T_k u_\varepsilon),$$

where  $T_k u_\varepsilon = (u_\varepsilon \vee k) \wedge (k + 1)$ . If  $|A_k^\varepsilon|^{\frac{1}{2}} \leq \varepsilon^\alpha$ , then we apply Step 2. Thus we get

$$\sum_{k=1}^{\infty} |A_k^\varepsilon|^{\frac{1}{2}} \leq C \sum_{k=1}^{\infty} \frac{1}{(k - \sigma)^\gamma} + I_\varepsilon(T_k u_\varepsilon) \leq C(1 + I_\varepsilon(u_\varepsilon))$$

and this gives (48).

*Step 4.* We conclude the proof by noting that for any decreasing sequence  $a_k$  of positive numbers we have

$$\sum_{k=1}^{\infty} k a_k \leq \sup_{k \geq 1} (k a_k^{\frac{1}{2}}) \sum_{k=1}^{\infty} a_k^{\frac{1}{2}} \leq \left( \sum_{k=1}^{\infty} a_k^{\frac{1}{2}} \right)^2. \quad (49)$$

We apply this inequality with  $a_k = |A_{k-2}^\varepsilon|$  and using Step 3, we get

$$\int_0^\infty t |\{x \in Q'; u_\varepsilon > t\}| dt \leq \sum_{k=1}^{\infty} k |\{x \in Q'; u_\varepsilon > k - 1\}| \leq \sum_{k=1}^{\infty} k |A_{k-2}^\varepsilon| \leq C. \quad (50)$$

Hence

$$\int_{Q' \cap \{u_\varepsilon > 0\}} |u_\varepsilon|^2 dx = \int_0^\infty t |\{x \in Q'; u_\varepsilon > t\}| dt \leq C.$$

The conclusion follows by arguing in a similar way for the negative part of  $u_\varepsilon$ . ○

## 4.2 Compactness

**Proof of Theorem 12.** We may assume that  $|\{u_\varepsilon \geq 0\}| \geq \frac{1}{2}|Q|$  and  $|\{u_\varepsilon \leq 1\}| \geq \frac{1}{2}|Q|$ , since otherwise we may replace  $\{u_\varepsilon\}$  with  $\{u_\varepsilon - a_\varepsilon\}$ , where

$$a_\varepsilon = \max\{a \in \mathbf{Z} : |\{u_\varepsilon - a > 0\}| \geq \frac{1}{2}|Q|\}. \quad (51)$$

Furthermore we may suppose that  $I_\varepsilon(u_\varepsilon) \leq C_1$  and by passage to a subsequence  $\liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) = \lim_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon)$ .

Theorem 13 shows that  $u_\varepsilon$  is uniformly bounded in  $L^2(Q')$  for every compactly contained concentric subsquare  $Q' \subset\subset Q$ . Since  $u_\varepsilon$  is periodic and since we can apply Theorem 13 also to a translation of  $u_\varepsilon$  we easily see that  $u_\varepsilon$  is bounded in  $L^2(T^2)$ . Hence there exists  $u \in L^2(T^2)$  such that, up a subsequence,

$$u_\varepsilon \rightharpoonup u \quad \text{in } L^2(T^2). \quad (52)$$

To obtain strong convergence in  $L^1$  we consider again the truncation operator  $T_a u = (u \vee a) \wedge (a + 1)$ , for all  $a \in \mathbf{Z}$ . Clearly by (8)

$$\frac{1}{|\log \varepsilon|} \left( [T_a u_\varepsilon]_{H^{\frac{1}{2}}(Q)}^2 + \frac{1}{\varepsilon} \int_Q \text{dist}^2(T_a u_\varepsilon, \{a, a + 1\}) dx \right) \leq 2I_\varepsilon(T_a u_\varepsilon) \leq 2I_\varepsilon(u_\varepsilon) \leq C.$$

By [3], Theorem 4.7, (see also [2]) we have that, up to a subsequence, for every  $a \in \mathbf{Z}$ , there exists an  $L^1$  function  $u_a \in BV(Q, \{a, a + 1\})$  such that

$$T_a u_\varepsilon \rightarrow u_a \quad \text{in } L^1(T^2) \quad (53)$$

and

$$\int_{T^2} |Du_a| \leq C \liminf_{\varepsilon \rightarrow 0} I_\varepsilon(T_a u_\varepsilon). \quad (54)$$

Now consider  $M \in \mathbf{N}$  and the truncation operator  $T^M u = (u \vee -M) \wedge M$ . By (53) there exists  $u^M \in BV(T^2, \mathbf{Z})$  such that

$$T^M u_\varepsilon \rightarrow u^M \quad \text{in } L^1(T^2).$$

Since  $|\{|u_\varepsilon| > M\}|^{\frac{1}{2}} \leq \|u_\varepsilon\|_{L^2}/M$ , weak lower semi-continuity of the  $L^1$  norm and the  $L^2$  bound yield

$$\|u^M - u\|_{L^1(T^2)} \leq \liminf_{\varepsilon \rightarrow 0} \|T^M u_\varepsilon - u_\varepsilon\|_{L^1(T^2)} \leq \liminf_{\varepsilon \rightarrow 0} 2 \int_{\{|u_\varepsilon| > M\}} |u_\varepsilon| dx \leq \frac{C}{M}.$$

Now  $u_\varepsilon - u = u_\varepsilon - T^M u_\varepsilon + T^M u_\varepsilon - u^M + u^M - u$  and thus

$$\limsup_{\varepsilon \rightarrow 0} \|u_\varepsilon - u\|_{L^1(T^2)} \leq \frac{2C}{M}.$$

Since  $M$  was arbitrary this shows that  $u_\varepsilon \rightarrow u$  in  $L^1(T^2)$  (and hence in all  $L^q(T^2)$ , for  $q < 2$ ) and thus  $u_a = T_a u$ . It is easy to verify that

$$\sum_{a \in \mathbf{Z}} \iint_{Q \times Q} K_\nu(x - y) |T_a v_\varepsilon(x) - T_a v_\varepsilon(y)|^2 dx dy \leq \iint_{Q \times Q} K_\nu(x - y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy.$$

Hence (54) yields (32), after summation over  $a$ . ○

### 4.3 The effect of the pinning

Theorem 12 gives compactness up to translation by integers of any sequences  $\{u_\varepsilon\}$  which satisfies  $\liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) < \infty$ . If in addition  $u_\varepsilon$  is also subject to the pinning condition, this eliminates the translation invariance of the problem and yields, via a Poincaré's inequality, an  $L^2$  bound and compactness of the sequence  $u_\varepsilon$ .

**Proposition 16** *Let  $u_\varepsilon$  such that*

$$\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) < +\infty,$$

*then  $u_\varepsilon$  is bounded in  $L^2(T^2)$  and relatively compact in  $L^q(T^2)$ , for every  $q < 2$ . Every cluster point  $u$  belongs to  $BV(T^2, \mathbf{Z})$  and satisfies*

$$\int_{T^2} |Du| \leq C \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon). \quad (55)$$

**Proof.** Without loss of generality, by (8) we may assume that

$$[u_\varepsilon]_{H^{\frac{1}{2}}(T^2)}^2 \leq C |\log \varepsilon|. \quad (56)$$

By Theorem 12 we know that for every  $\varepsilon$  there exists an integer number  $a_\varepsilon$  such that  $\{u_\varepsilon - a_\varepsilon\}$  is bounded in  $L^2(T^2)$ , relatively compact in  $L^q(T^2)$ , for  $q < 2$ , and every cluster point satisfies (55). In order to conclude it is enough to have an  $L^2$  bound for  $\{u_\varepsilon\}$  which yields that the sequence  $\{a_\varepsilon\}$  is bounded. This can be obtained through the Poincaré inequality. Fix  $\rho_\varepsilon = \sqrt{\frac{(L+1)\varepsilon}{|\log \varepsilon|}}$  ( $L$  is the constant given by (14)). With a little abuse of notation we denote by  $Q_{\rho_\varepsilon}^j$  the squares of a lattice on  $Q$  of size approximately  $\rho_\varepsilon$ . Applying the Poincaré inequality (13), scaled to the square  $Q_{\rho_\varepsilon}^j$ , we get

$$\int_{Q_{\rho_\varepsilon}^j} |u_\varepsilon|^2 dx \leq C_1 \rho_\varepsilon \left( 1 + \frac{\rho_\varepsilon}{\text{Cap}(\{u_\varepsilon = 0\} \cap Q_{\rho_\varepsilon}^j \times \{0\})} \right) [u_\varepsilon]_{H^{\frac{1}{2}}(Q_{\rho_\varepsilon}^j)}^2. \quad (57)$$

By our choice of  $\rho_\varepsilon$  and assumption (14) we have

$$1 \leq \#(\mathcal{I}_\varepsilon(Q_{\rho_\varepsilon}^j)) \leq 2L + 1$$

and then  $\text{Cap}(\{u_\varepsilon = 0\} \cap Q_{\rho_\varepsilon}^j \times \{0\}) > C'R\varepsilon$ . Taking the sum over all  $j$  in (57), by (56), we get

$$\int_Q |u_\varepsilon|^2 dx \leq \sum_j C_1 \rho_\varepsilon \left( 1 + \frac{\rho_\varepsilon}{C'R\varepsilon} \right) [u_\varepsilon]_{H^{\frac{1}{2}}(Q_{\rho_\varepsilon}^j)}^2 \leq C \rho_\varepsilon \left( 1 + \frac{\rho_\varepsilon}{C'R\varepsilon} \right) |\log \varepsilon| \leq C$$

which concludes the proof.  $\square$

## 5 Lower bound

**Theorem 17** *Let  $\{u_\varepsilon\}$  be a sequence such that  $\liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) < +\infty$ . Assume that  $\{u_\varepsilon\}$  converges strongly in  $L^1(T^2)$  to some function  $u$ , then  $u \in BV(T^2, \mathbf{Z})$  and*

$$\int_{T^2} \gamma \left( \frac{Du}{|Du|} \right) |Du| = \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1 \leq \liminf_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon), \quad (58)$$

where the anisotropic line energy density  $\gamma(n)$  is defined by (25)

The proof of Theorem 17 is based on a blow-up argument. Let  $n \in S^1$  and denote by  $Q^n$  a square centered at 0, with size 1 and parallel to  $n$ . We now estimate from below the energy of a function on  $Q^n$  which is close to the characteristic function of the half plane  $\{n \cdot x > 0\}$ . We denote the latter function by  $u_0^n = \chi_{\{n \cdot x > 0\}}$ .

**Lemma 18** *Fix  $0 < \delta < \frac{1}{2}$  and  $\frac{1}{2} < \alpha < 1$ . Then there exist  $\delta_0 > 0$  and  $\varepsilon_0 > 0$  such that for every  $n \in S^1$ , for every  $\varepsilon \in (0, \varepsilon_0)$  and for every  $u \in L^1(Q)$  satisfying*

$$\int_{Q^n} |u - u_0^n| dx \leq \delta_0 \quad (59)$$

and

$$\int_{Q^n} \text{dist}^2(u, \mathbf{Z}) dx \leq \varepsilon^\alpha \quad \text{for some } , \quad (60)$$

we have

$$\iint_{Q^n \times Q^n} \Gamma_\nu(x-y) |u(x) - u(y)|^2 dx dy \geq \gamma(n) \alpha (1-\delta) |\log \varepsilon|, \quad (61)$$

where  $\gamma(n)$  is defined by (25).

**Proof.** After rotation we can easily restrict our analysis to the case  $n = e_1$ , in which the square  $Q^n$  reduces to  $Q = (-\frac{1}{2}, \frac{1}{2})^2$ . It then suffices to prove the statement with  $\gamma(n) = \gamma(e_1)$  and we write from now  $u_0$  instead of  $u_0^{\varepsilon_1}$ . By scaling we may also assume that  $\gamma(e_1) = 2$ . Since  $\Gamma_\nu$  is homogeneous of degree  $-3$  we have then

$$\Gamma_\nu^1(x_1) := \int_{\mathbf{R}} \Gamma_\nu(x_1, x_2) dx_2 = \frac{1}{x_1^2}. \quad (62)$$

We also assume that  $0 \leq u \leq 1$  (otherwise we truncate the function by 0 and 1). Let us consider the following sub and super-level sets

$$A = \left\{ u < \frac{\delta}{8} \right\} \quad \text{and} \quad B = \left\{ u > 1 - \frac{\delta}{8} \right\}.$$

Then

$$\iint_{Q \times Q} \Gamma_\nu(x-y) |u(x) - u(y)|^2 dx dy \geq 2 \left(1 - \frac{\delta}{4}\right)^2 \int_A \int_B \Gamma_\nu(x-y) dx dy.$$

Hence it is sufficient to show that

$$\int_A \int_B \Gamma_\nu(x-y) dx dy \geq \alpha \left(1 - \frac{\delta}{2}\right) |\log \varepsilon|. \quad (63)$$

Using the change of variables,  $y = x + z$ , we get

$$\int_A \int_B \Gamma_\nu(x-y) dx dy = \int_{\mathbf{R}^2} \Gamma_\nu(-z) \left( \int_{A \cap (B-z)} dx \right) dz = \int_{\mathbf{R}^2} \Gamma_\nu(z) |A \cap (B-z)| dz. \quad (64)$$

Let  $\frac{1}{2} < \sigma < \alpha$ . By (60) we know that for  $\varepsilon_0$  small enough the set  $E = Q \setminus (A \cup B)$  satisfies

$$|E| \leq \varepsilon^\sigma. \quad (65)$$

If the function  $u$  is independent of  $x_2$  and  $E$  is a simple strip, then  $|A \cap (B-z)| \geq z_1 - \varepsilon^\sigma$  and the result can be obtained from (64) by an explicit computation. In general these conditions are not satisfied, but the idea is that they are approximatively satisfied. In order to deal with the general case let us estimate the difference between  $\chi_B$  and the characteristic function of  $\{x_1 > 0\}$ , i.e.  $u_0$ . If  $x_1 > 0$  and  $x \in A$ , then  $|u - u_0| \geq \frac{1}{2}$ . Thus

$$|\chi_B - u_0| = |\chi_B - 1| \leq \chi_E + \chi_A \leq \chi_E + 2|u - u_0|.$$

Similarly, if  $x_1 < 0$  and  $x \in B$ , then  $|u - u_0| \geq \frac{1}{2}$ . Hence

$$|\chi_B - u_0| \leq \chi_E + 2|u - u_0| \quad \text{in } Q. \quad (66)$$

Now consider a concentric subsquare  $Q'$  of  $Q$ , i.e.  $Q' = \rho Q$  with  $\rho < 1$ . By (65) we have

$$|A \cap Q'| + |B \cap Q'| \geq |Q'| - \varepsilon^\sigma \quad (67)$$

Suppose that  $z$  satisfies  $|z|_\infty = \max(|z_1|, |z_2|) < (1-\rho)/2$  and  $z_1 > 0$ . Since  $|(B-z) \cap Q'| = |B \cap (Q'+z)|$ , we have

$$\begin{aligned} |(B-z) \cap Q'| - |B \cap Q'| &\geq \int_{Q'+z} u_0 dx - \int_{Q'} u_0 dx - \int_{(Q'+z) \Delta Q'} |\chi_B - u_0| dx \\ &\geq \rho z_1 - \int_U |\chi_B - u_0| dx, \end{aligned}$$

where  $U$  is the annulus  $U = (\rho + |z|_\infty)Q \setminus (\rho - |z|_\infty)Q$  and  $(Q'+z) \Delta Q' \subset U$ . Together with (67) this yields

$$\begin{aligned} m(z) := |A \cap (B-z)| &\geq |A \cap (B-z) \cap Q'| \geq |A \cap Q'| + |(B-z) \cap Q'| - |Q'| \quad (68) \\ &\geq z_1 \rho - \varepsilon^\sigma - \int_U |\chi_B - u_0| dx. \end{aligned}$$

We now must choose  $\rho$  and  $U$  properly in order to get in (68) the right hand side large enough. We fix  $\delta_2 > 0$  and  $\rho_0 = 1 - \delta_2$  and we cover  $Q \setminus \rho_0 Q$  with annuli of thickness  $2|z|_\infty$ , i.e. we take  $\rho_i = \rho_0 + 2i|z|_\infty$ ,  $U_i = \rho_i Q \setminus \rho_{i-1} Q$ , with  $i = 1, \dots, k$  and  $k < \delta_2/4|z|_\infty \leq k+1 \leq 2k$ . We apply (68) with  $\rho = \rho_i$  and  $U = U_i$ , we sum all the inequalities for  $i$  from 1 to  $k$ , we divide by  $k$  and we get

$$m(z) \geq z_1 \rho_0 - \varepsilon^\sigma - \frac{1}{k} \int_Q |\chi_B - u_0| dx.$$

This, together with (65), (66) and (59) yields

$$m(z) \geq (1 - \delta_2)z_1 - \varepsilon^\sigma - \frac{8}{\delta_2} |z|_\infty (2\delta_0 + \varepsilon^\sigma),$$

whenever  $|z|_\infty \leq \delta_2/2$ . We now assume  $\varepsilon_0^\sigma \leq 2\delta_0$  and choose  $\delta_2 = 2\sqrt{\delta_0}$ . This gives

$$m(z) \geq (1 - 2\sqrt{\delta_0})z_1 - \varepsilon^\sigma - 16\sqrt{\delta_0}|z|_\infty, \quad \text{if } |z|_\infty \leq \sqrt{\delta_0}. \quad (69)$$

Then, in order to conclude from here together with (64) and to obtain (63), we must estimate the following integrals:

$$I_1 = \int_{\{\varepsilon^\sigma \leq z_1 \leq 1\}} \varepsilon^\sigma \Gamma_\nu(z) dz_1 dz_2 \leq \varepsilon^\sigma \int_{\varepsilon^\sigma}^1 \Gamma_\nu^1(z_1) dz_1 \leq 1, \quad (70)$$

where we used (62). Since  $|z|_\infty \leq 2|z|$  and  $\int_{\mathbf{R}} (a^2 + z^2)^{-1} dz_2 = \pi/|a|$

$$I_2 = \int_{\{\varepsilon^\sigma \leq z_1 \leq 1\}} |z|_\infty \Gamma_\nu(z) dz_1 dz_2 \leq \int_{\varepsilon^\sigma}^1 \frac{C}{|z_1|} dz_1 \leq C\sigma |\log \varepsilon|. \quad (71)$$

Finally, by the definition of  $\Gamma_\nu^1$  and the fact that  $\Gamma_\nu(z) \leq C/z_2^2$ , we get

$$\begin{aligned} I_3 = \int_{\varepsilon^\sigma}^{\delta_0} \int_{\{|z_2| \leq \delta_0\}} z_1 \Gamma_\nu(z) dz_2 dz_1 &\geq \int_{\varepsilon^\sigma}^{\delta_0} z_1 (\Gamma_\nu^1(z_1) - \frac{C}{\delta_0^2}) dz_1 \quad (72) \\ &\geq \log \frac{\delta_0}{\varepsilon^\sigma} - C = \sigma |\log \varepsilon| + \log \delta_0 - C. \end{aligned}$$

Thus by (69) - (72) we have

$$\begin{aligned} \frac{1}{|\log \varepsilon|} \int_{\mathbf{R}^2} m(z) \Gamma_\nu(z) dz &\geq \frac{1}{|\log \varepsilon|} \left[ (1 - 2\sqrt{\delta_0})I_3 - I_1 - 16\sqrt{\delta_0}I_2 \right] \\ &\geq (1 - 2\sqrt{\delta_0})\sigma - C(\sqrt{\delta_0} + \frac{1}{|\log \varepsilon_0|}) \end{aligned}$$

and this proves (63) if  $\alpha - \sigma$ ,  $\delta_0$  and  $\varepsilon_0$  are chosen sufficiently small.

○

With the following lemma we prove that the measure defined on the product  $T^2 \times T^2$  by

$$\mu_\varepsilon(A \times B) = \frac{1}{|\log \varepsilon|} \int_A \int_B K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \quad (73)$$

converges weakly to a measure whose support is on the diagonal and that it can be estimated from below by a measure concentrated on the jump set of the limit function  $u$ .

**Lemma 19** *Let  $u_\varepsilon$  be a sequence which converges strongly in  $L^1(T^2)$  to some function  $u \in BV(T^2, \mathbf{Z})$  and assume  $0 \leq u_\varepsilon \leq 1$ . Let  $\mu_\varepsilon$  be defined by (73) and let  $\mu$  be its weak\*-limit (for a subsequence). Then  $\mu$  is concentrated on the diagonal  $D = \{(x, x) : x \in T^2\}$ , i.e.  $\mu(E) = 0$  if  $E \cap D = \emptyset$ . Moreover the measure  $\lambda$  on  $T^2$  which is defined by*

$$\lambda(A) = \mu(\{(x, x) : x \in A\}), \quad (74)$$

satisfies

$$\lambda \geq \gamma(n) d\mathcal{H}^1 \llcorner S_u, \quad (75)$$

where  $S_u$  is the jump set of  $u$ .

**Proof.** To prove that  $\text{supp} \mu \subseteq D$  it is enough to show that for any continuous nonnegative function  $\varphi : T^2 \times T^2 \rightarrow \mathbf{R}$  with  $\text{supp} \varphi \cap D = \emptyset$  we have  $\int \varphi d\mu = 0$ . Let  $\delta = \text{dist}(\text{supp} \varphi, D) > 0$ . Since  $u_\varepsilon$  is bounded by 1 we have

$$\begin{aligned} \int_{T^2} \varphi(x, y) d\mu &= \lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \iint_{T^2 \times T^2} \varphi(x, y) K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \\ &\leq \lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \frac{C}{\delta^3} \iint_{T^2 \times T^2} \varphi(x, y) dx dy = 0. \end{aligned}$$

Thus  $\mu$  is concentrated on  $D$ . Hence we can define the measure  $\lambda$  on  $T^2$  by

$$\lambda(A) = \mu(\{(x, x) : x \in A\}) = \mu(A \times A).$$

In order to conclude it is enough to show that for  $\mathcal{H}^1$ -a.e.  $x_0 \in S_u$  we have

$$\liminf_{r \rightarrow 0} \frac{\lambda(Q_r^n(x_0))}{r} = \liminf_{r \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \frac{\mu_\varepsilon(Q_r^n(x_0) \times Q_r^n(x_0))}{r} \geq \gamma(n), \quad (76)$$

where, with a little abuse of notation,  $Q_r^n$  denotes the square centered at  $x_0$  with side  $r$  parallel to the normal  $n$  on  $S_u$  at  $x_0$ .

For  $\mathcal{H}^1$ -a.e.  $x_0 \in S_u$  we have

$$\lim_{r \rightarrow 0} \int_{Q_1^n} |u(rx + x_0) - \chi_{\{x \cdot n > 0\}}| dx = 0.$$

Fix such a  $x_0 \in S_u$ . We will proceed with a blow-up argument. Consider the sequence  $v_\varepsilon$  obtained from  $u_\varepsilon$  by a rescaling, i.e.  $v_\varepsilon(x) = u_\varepsilon(rx + x_0)$ . By a change of variables we have

$$\frac{1}{r} \mu_\varepsilon(Q_r^n(x_0) \times Q_r^n(x_0)) = \iint_{Q_1^n \times Q_1^n} K_\nu^r(x-y) |v_\varepsilon(x) - v_\varepsilon(y)|^2 dx dy,$$

where  $K_\nu^r(t) = r^3 K_\nu(rt)$ . From Proposition 1 and the homogeneity of  $\Gamma_\nu$  we have

$$\frac{1}{r} \mu_\varepsilon(Q_r^n(x_0) \times Q_r^n(x_0)) = \iint_{Q_1^n \times Q_1^n} \Gamma_\nu(x-y) |v_\varepsilon(x) - v_\varepsilon(y)|^2 dx dy + o(1), \quad (77)$$

as  $r$  goes to zero. The idea is to use Lemma 18 with the function  $v_\varepsilon$ . Fix  $0 < \delta < \frac{1}{2}$  and  $\frac{1}{2} < \alpha < 1$  and let  $\delta_0 > 0$  and  $\varepsilon_0 > 0$  be the constants given by Lemma 18. For  $r > 0$  small enough we have

$$\int_{Q_1^n} |u(rx + x_0) - \chi_{\{x \cdot n > 0\}}| dx \leq \frac{\delta_0}{2}$$

and thus, for such  $r$ , we can choose  $\varepsilon_0$  such that for every  $\varepsilon < \varepsilon_0$

$$\frac{1}{r^2} \int_{Q_r^n} |u_\varepsilon(x) - u(x)| dx \leq \frac{\delta_0}{2}.$$

We then get

$$\int_{Q_1^n} |v_\varepsilon(x) - \chi_{\{x \cdot n > 0\}}| dx \leq \int_{Q_1^n} |v_\varepsilon(x) - u(rx + x_0)| dx + \int_{Q_1^n} |u(rx + x_0) - \chi_{\{x \cdot n > 0\}}| \leq \delta_0.$$

Since  $v_\varepsilon$  also satisfies (60), applying Lemma 18 we obtain

$$\frac{1}{|\log \varepsilon|} \iint_{Q_1^n \times Q_1^n} \Gamma_\nu(x - y) |v_\varepsilon(x) - v_\varepsilon(y)|^2 dx dy \geq \gamma(n) \alpha (1 - \delta). \quad (78)$$

By (77), taking the limit as  $\varepsilon \rightarrow 0$  and then  $r \rightarrow 0$ , we get

$$\liminf_{r \rightarrow 0} \frac{\lambda(Q_r^n(x_0))}{r} \geq \gamma(n) \alpha (1 - \delta)$$

which gives (76) since  $\alpha \in (\frac{1}{2}, 1)$  and  $\delta > 0$  where arbitrary.  $\circ$

**Proof of Theorem 17.** By the compactness result (Theorem 12) we can deduce that  $u \in BV(T^2, \mathbf{Z})$ . We obtain the result by Lemma 19 using a truncation argument. Let  $j \in \mathbf{Z}$  and let us consider the truncations  $T_j u_\varepsilon = (u_\varepsilon \vee j) \wedge (j + 1)$ . Clearly each truncation  $T_j u_\varepsilon$  converges to  $T_j u$  and, up to a translation, it satisfies the assumptions of Lemma 19. Thus we have that for any  $j \in \mathbf{Z}$

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \iint_{T^2 \times T^2} K_\nu(x - y) |T_j u_\varepsilon(x) - T_j u_\varepsilon(y)|^2 dx dy \geq \int_{S_{T_j u}} \gamma(n) d\mathcal{H}^1. \quad (79)$$

Note that  $|u_\varepsilon(x) - u_\varepsilon(y)|^2 \geq \sum_{j \in \mathbf{Z}} |T_j u_\varepsilon(x) - T_j u_\varepsilon(y)|^2$  for every  $x, y \in T^2$ , that  $\cup_{j \in \mathbf{Z}} S_{T_j u} = S_u$   $\mathcal{H}^1$ -a.e., and that  $||u|(x)| = \sum_{j \in \mathbf{Z}} |[T_j u](x)|$   $\mathcal{H}^1$ -a.e.  $x \in S_u$ . Hence, by (79), we have

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} & \iint_{T^2 \times T^2} K_\nu(x - y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \\ & \geq \sum_{j \in \mathbf{Z}} \liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \iint_{T^2 \times T^2} K_\nu(x - y) |T_j u_\varepsilon(x) - T_j u_\varepsilon(y)|^2 dx dy \\ & \geq \sum_{j \in \mathbf{Z}} \int_{S_{T_j u}} \gamma(n) d\mathcal{H}^1 = \int_{S_u} \gamma(n) |[u](x)| d\mathcal{H}^1(x), \end{aligned} \quad (80)$$

which concludes the proof.  $\circ$

**Remark 20** Reasoning as in the proof above we conclude that for any sequence  $u_\varepsilon$  converging to  $u$  in  $L^1$  and satisfying  $I_\varepsilon(u_\varepsilon) \leq C$ , and for every  $\varphi \geq 0$  continuous function on  $T^2 \times T^2$  we get

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \iint_{T^2 \times T^2} \varphi(x, y) K_\nu(x - y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \geq \int_{S_u} \varphi(x, x) \gamma(n) |[u](x)| d\mathcal{H}^1(x)$$

and thus  $\lambda \geq \gamma(n) |[u]| \mathcal{H}^1 \llcorner S_u$ , where  $\lambda(A) = \mu(\{(x, x) : x \in A\})$  and  $\mu$  is the weak\*-limit of the measures  $\mu_\varepsilon$  defined in (73). Moreover it is easy to see that if  $u_\varepsilon$  is also bounded in  $L^2$ ,  $\mu$  is concentrated on the diagonal  $D$ , i.e.  $\lambda(A) = \mu(A \times A)$ .

Combining Theorem 17 and the results proved in [8] we now establish the following lower bound for the  $\Gamma$ -limit of the functional  $F_\varepsilon$ .

**Theorem 21** *Assume that  $N_\varepsilon\varepsilon/|\log\varepsilon| \rightarrow \Lambda$  and that the discs  $B_{R\varepsilon}^i$  are uniformly distributed and well separated. Let  $u_\varepsilon$  be a sequence such that  $\liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon) < +\infty$ . Assume that  $u_\varepsilon$  converges strongly in  $L^1(T^2)$  to some function  $u$ , then  $u \in BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$  and*

$$\int_{S_u} \gamma(n)|[u]|d\mathcal{H}^1 + \Lambda \int_{T^2} D_\nu(u, B_R) dx \leq \liminf_{\varepsilon \rightarrow 0} F_\varepsilon(u_\varepsilon), \quad (81)$$

where  $\gamma(n)$  is defined by (25) and  $D_\nu(\cdot, B_R)$  is defined by (17).

The proof is also based on the following two lemmas proved in [8].

**Lemma 22** *Given  $\mathcal{R} : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ , with  $\mathcal{R}(\varepsilon) \rightarrow \infty$  as  $\varepsilon \rightarrow 0$ , there exists a function  $\omega : \mathbf{R}_+ \times \mathbf{R}_+ \rightarrow \mathbf{R}_+$ , with  $\omega(\varepsilon, \delta) \rightarrow 0$  as  $(\varepsilon, \delta) \rightarrow (0, 0)$ , such that the following statement holds. Let  $a \in \mathbf{Z}$ . If  $\zeta \in H^{\frac{1}{2}}(B_{\mathcal{R}(\varepsilon)})$  satisfies*

$$\int_{B_{\mathcal{R}(\varepsilon)}} |\zeta - a| dx \leq \delta \quad (82)$$

and  $\zeta = 0$  on  $B_R$ , then

$$\int_{B_{\mathcal{R}(\varepsilon)}} \text{dist}^2(\zeta, \mathbf{Z}) dx + \iint_{B_{\mathcal{R}(\varepsilon)} \times B_{\mathcal{R}(\varepsilon)}} K^\varepsilon(x-y)|\zeta(x) - \zeta(y)|^2 dx dy \geq D_\nu(a, B_R) - \omega(\varepsilon, \delta), \quad (83)$$

where  $K^\varepsilon(t) = \varepsilon^3 K_\nu(\varepsilon t)$ .

**Lemma 23** *There exists a positive constant  $C$  such that for every  $0 < \rho < \hat{\rho}$  the following inequality holds*

$$\int_{B_\rho} |u| dx \leq \int_{B_{\hat{\rho}}} |u| dx + \frac{C}{\sqrt{\rho}} [u]_{H^{\frac{1}{2}}(B_{\hat{\rho}})} \quad (84)$$

for all  $u \in H^{\frac{1}{2}}(B_{\hat{\rho}})$ .

**Proof of Theorem 21.** The proof is based on a blow-up argument like the proof of Theorem 17. Let  $\mu_\varepsilon$  be the measure defined in (73), let  $\mu$  be its weak\*-limit,  $\lambda$  defined by (74) and define  $\eta_\varepsilon$  be the following measure

$$\eta_\varepsilon(A) = \frac{1}{\varepsilon|\log\varepsilon|} \int_A \text{dist}^2(u_\varepsilon, \mathbf{Z}) dx. \quad (85)$$

Since  $\eta_\varepsilon$  is bounded, it converges weakly\* up to a subsequence. Let  $\eta$  be its weak\* limit. The main step is now to prove that

$$\lambda + \eta \geq \Lambda D_\nu(u(x), B_R) dx, \quad (86)$$

i.e. that for a.e.  $x \in T^2$  we have

$$\lim_{r \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} \frac{\mu_\varepsilon(Q_r(x) \times Q_r(x)) + \eta_\varepsilon(Q_r(x))}{|Q_r|} \geq \Lambda D_\nu(u(x), B_R). \quad (87)$$

Let us fix  $a \in \mathbf{Z}$  and  $x_0$  be a Lebesgue point of  $u$  with  $u(x_0) = a$ . In order to simplify the notation assume that  $x_0 = 0$ . The proof of (87) follows using Lemmas 22 and 23 and the same strategy of the proof of the lower bound in [8] taking into account that here the domain is  $Q_r$  and the regime for the obstacles is different, i.e.  $N_\varepsilon \approx |\log\varepsilon|/\varepsilon$ . We repeat it here for the convenience of the reader.

Consider on  $Q_r$  a lattice of squares, denoted by  $Q_j^\varepsilon$ , of size approximately  $1/\sqrt{N_\varepsilon}$ . Let  $\widehat{Q}_j^\varepsilon$  be concentric squares of three times the size. Since each point is contained at most in 9 of the squares  $\widehat{Q}_j^\varepsilon$  we have

$$\sum_j \iint_{\widehat{Q}_j^\varepsilon \times \widehat{Q}_j^\varepsilon} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^2}{|x - y|^3} dx dy \leq C |\log \varepsilon| \approx N_\varepsilon \varepsilon$$

and

$$\sum_j \int_{\widehat{Q}_j^\varepsilon} |u_\varepsilon - a| dx \leq \omega_r,$$

where  $\omega_r \rightarrow 0$  as  $r \rightarrow 0$ . Let  $\theta > 0$ . Recall that  $\mathcal{I}_\varepsilon(Q_r)$  is the set of indices  $i$  such that  $x_\varepsilon^i \in Q_r$  where  $x_\varepsilon^i$  are the center of the discs  $B_{R\varepsilon}^i$ . Then there exist a set of indices  $\mathcal{J}^\varepsilon(Q_r)$  such that  $\#(\mathcal{J}^\varepsilon(Q_r)) \geq (1 - \theta)\#(\mathcal{I}_\varepsilon(Q_r))$  and a constant  $C_\theta$  such that

$$\iint_{\widehat{Q}_j^\varepsilon \times \widehat{Q}_j^\varepsilon} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^2}{|x - y|^3} dx dy \leq C_\theta \varepsilon$$

and

$$\int_{\widehat{Q}_j^\varepsilon} |u_\varepsilon - a| dx \leq C_\theta \omega_r$$

for all  $j \in \mathcal{J}^\varepsilon(Q_r)$ . Let  $0 < \delta < 1$ . By applying Lemma 23 with  $\rho = \varepsilon^\beta$ , with  $\frac{1}{2} < \beta < 1$ , and  $\widehat{\rho} = \frac{1}{\sqrt{N_\varepsilon}}$ , for each  $x_\varepsilon^i \in Q_j^\varepsilon$  we also have

$$\int_{B_{\varepsilon^\beta}^i} |u_\varepsilon - a| dx \leq \delta \quad \text{if } \varepsilon \leq \varepsilon_0(\delta, \theta) \text{ and } r \leq r_0(\theta). \quad (88)$$

Then by Lemma 22 applied with  $\mathcal{R}(\varepsilon) = \varepsilon^{\beta-1}$  we get (after the scaling  $X \rightarrow x/\varepsilon$ )

$$\mu_\varepsilon(B_{\varepsilon^\beta}^i \times B_{\varepsilon^\beta}^i) + \eta_\varepsilon(B_{\varepsilon^\beta}^i) \geq \frac{\varepsilon}{|\log \varepsilon|} (D_\nu(a, B_R) - \omega(\varepsilon, \delta))$$

for any  $i$  such that  $x_\varepsilon^i \in Q_j^\varepsilon$ . Since the points  $x_\varepsilon^i$  are assumed to be well separated (see (15)), summation over all  $i$  yields

$$\mu_\varepsilon(Q_r \times Q_r) + \eta_\varepsilon(Q_r) \geq \frac{\varepsilon}{|\log \varepsilon|} \left[ \sum_{j \in \mathcal{J}^\varepsilon(Q_r)} \#(\mathcal{I}_\varepsilon(Q_j^\varepsilon)) \right] (D_\nu(a, B_R) - \omega(\varepsilon, \delta))$$

The uniform distribution of the obstacles (see condition (14)) implies that

$$\begin{aligned} \sum_{j \in \mathcal{J}^\varepsilon(Q_r)} \#(\mathcal{I}_\varepsilon(Q_j^\varepsilon)) &= \#(\mathcal{I}_\varepsilon(Q_r)) - \sum_{j \notin \mathcal{J}^\varepsilon(Q_r)} \#(\mathcal{I}_\varepsilon(Q_j^\varepsilon)) \geq \#(\mathcal{I}_\varepsilon(Q_r)) - \sum_{j \notin \mathcal{J}^\varepsilon(Q_r)} (N_\varepsilon |Q_j^\varepsilon| + L) \\ &= \#(\mathcal{I}_\varepsilon(Q_r)) - (L + 1)\#(\{j : j \notin \mathcal{J}^\varepsilon(Q_r)\}). \end{aligned}$$

Since  $\#(\{j : j \notin \mathcal{J}^\varepsilon(Q_r)\}) \leq \#(\mathcal{I}_\varepsilon(Q_r))\theta$ , by (14) we get

$$\begin{aligned} \mu_\varepsilon(Q_r \times Q_r) + \eta_\varepsilon(Q_r) &\geq \frac{\varepsilon \#(\mathcal{I}_\varepsilon(Q_r)) (1 - \theta(L + 1))}{|\log \varepsilon|} (D_\nu(a, B_R) - \omega(\varepsilon, \delta)) \\ &\geq \Lambda |Q_r| (1 - \theta(L + 1)) (D_\nu(a, B_R) - \omega(\varepsilon, \delta)) + o(1) \end{aligned}$$

as  $\varepsilon$  goes to zero. Taking the limits  $\varepsilon \rightarrow 0$ , then  $r \rightarrow 0$ , then  $\delta \rightarrow 0$  and finally  $\theta \rightarrow 0$ , we obtain (87). Now we conclude easily by (87) and Lemma 19 together with Remark 20 that

$$\lambda + \eta \geq \Lambda D_\nu(u(x), B_R) dx \quad \text{and} \quad \lambda \geq \gamma(n) |[u]| \mathcal{H}^1 \llcorner S_u.$$

Since the two measures  $D_\nu(u(x), B_R) dx$  and  $\gamma(n) |[u]| \mathcal{H}^1 \llcorner S_u$  are mutually singular, we get

$$\lambda + \eta \geq \Lambda D_\nu(u(x), B_R) dx + \gamma(n) |[u]| \mathcal{H}^1 \llcorner S_u$$

and this concludes the proof. ○

## 6 Upper bound

In this section we establish an upper bound for the  $\Gamma$ -limit of  $I_\varepsilon$  and  $F_\varepsilon$ . This concludes the proof of the  $\Gamma$ -convergence result in the critical scaling (see Theorems 9 and 10).

**Theorem 24** *For every  $u \in BV(T^2, \mathbf{Z})$  there exists a sequence  $\{u_\varepsilon\}$  converging to  $u$  in  $L^q(T^2)$ , for any  $q < 2$ , such that*

$$\limsup_{\varepsilon \rightarrow 0} I_\varepsilon(u_\varepsilon) \leq \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1.$$

**Proof.** By a standard density argument, we can restrict our analysis to the case of  $u \in BV(T^2, \mathbf{Z})$  such that  $S_u$  is polygonal, with a finite number of sides, and  $|[u]| = 1$   $\mathcal{H}^1$ -a.e. on  $S_u$ . We also choose a fundamental domain  $Q$  of the torus such that  $|Du|(\partial Q) = 0$ . We construct  $u_\varepsilon$  simply mollifying  $u$  at the scale  $\varepsilon$ .

Fix  $\varphi \in C_c^\infty(B(0, 1))$ ,  $\varphi \geq 0$  and  $\int \varphi dx = 1$ , and set

$$u_\varepsilon = \varphi_\varepsilon * u \quad \text{where } \varphi_\varepsilon(x) = \varepsilon^{-2} \varphi\left(\frac{x}{\varepsilon}\right).$$

Since  $u_\varepsilon = u$  outside an  $\varepsilon$ -neighbourhood of  $S_u$  we clearly have that

$$\frac{1}{\varepsilon} \int_Q \text{dist}^2(u_\varepsilon, \mathbf{Z}) dx \leq C. \quad (89)$$

The conclusion follows if we prove that

$$\lim_{\varepsilon \rightarrow 0} \mu_\varepsilon(Q \times Q) = \lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \iint_{Q \times Q} K_\nu(x - y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy = \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1, \quad (90)$$

where the measure  $\mu_\varepsilon$  is defined as in (73). We first prove that the total variation of  $\mu_\varepsilon$  is bounded. Let  $A$  be an open subset of  $Q$  and denote by  $d(A)$  the diameter of  $A$ . Using a change of variables  $z = y - x$  we get

$$\mu_\varepsilon(A \times A) \leq \frac{C}{|\log \varepsilon|} \int_{|z| \leq d(A)} \frac{1}{|z|^3} \int_A |u_\varepsilon(x + z) - u_\varepsilon(x)|^2 dx dz. \quad (91)$$

Let us denote now by  $\mathcal{N}_\delta$  the  $\delta$ -neighbourhood of  $S_u$ . Since  $u \in BV(Q, \mathbf{Z})$  and  $S_u$  is polygonal (with a finite number of sides) we have

$$|\mathcal{N}_\delta \cap A| \leq Cd(A)\delta \quad (92)$$

and, by the definition of  $u_\varepsilon$ , we easily deduce that  $u_\varepsilon$  is constant on  $B_\delta(x)$  if  $x \notin \mathcal{N}_{\delta+\varepsilon}$ . Thus, using that  $u$  is bounded by a constant  $M$ , we get

$$\begin{aligned} \int_{2\varepsilon \leq |z| \leq d(A)} \frac{1}{|z|^3} \int_A |u_\varepsilon(x + z) - u_\varepsilon(x)|^2 dx dz &\leq 4M^2 \int_{2\varepsilon \leq |z| \leq d(A)} \frac{1}{|z|^3} |\mathcal{N}_{2|z|} \cap A| \\ &\leq Cd(A) \int_{2\varepsilon \leq |z| \leq d(A)} \frac{1}{|z|^2} dz \leq Cd(A) |\log \varepsilon|. \end{aligned}$$

On the other hand we have that  $|Du_\varepsilon| \leq C/\varepsilon$  and hence

$$\begin{aligned} \int_{|z| \leq 2\varepsilon} \frac{1}{|z|^3} \int_A |u_\varepsilon(x + z) - u_\varepsilon(x)|^2 dx dz &\leq C \int_{|z| \leq 2\varepsilon} \frac{1}{|z|^3} |\mathcal{N}_{2\varepsilon} \cap A| \frac{|z|^2}{\varepsilon^2} dz \\ &\leq Cd(A) \int_{|z| \leq 2\varepsilon} \frac{1}{|z| \varepsilon} dz \leq C. \end{aligned}$$

By (91) we get

$$\mu_\varepsilon(A \times A) \leq Cd(A) + o(1) \quad (93)$$

as  $\varepsilon$  tends to zero. In particular  $\mu_\varepsilon$  is bounded and thus, up to a subsequence, converges weakly\* to a measure  $\mu$ . Since  $u_\varepsilon$  is bounded in  $L^\infty$ , by Remark 20,  $\mu$  is concentrated on the diagonal and we can define as above the measure  $\lambda(A) = \mu(A \times A)$ . By the definition of  $u_\varepsilon$  it is easy to check that  $\text{supp}\lambda \subseteq S_u$ . Moreover, taking the limit  $\varepsilon \rightarrow 0$  in (93) we get a similar estimate for  $\lambda$ , i.e.  $\lambda(A) \leq Cd(A)$ . Taking the Radon-Nikodym derivative of  $\lambda$  with respect to  $\mathcal{H}^1$ , we deduce that

$$\lambda \leq C\mathcal{H}^1 \llcorner S_u. \quad (94)$$

We conclude the proof if we prove that for  $\mathcal{H}^1$ -a.e.  $x_0 \in S_u$

$$\lim_{r \rightarrow 0} \frac{\lambda(B_r(x_0))}{2r} \leq \gamma(n). \quad (95)$$

Indeed, this together with the lower bound implies (90). The latter inequality can be obtained by an explicit calculation. Let us assume, for the sake of simplicity, that  $x_0 = 0$  and that the normal  $n$  of  $S_u$  at  $x_0$  is  $e_1$ , i.e.  $S_u$  is locally contained in  $\{x_1 = 0\}$ . Then

$$u_\varepsilon(x) = \psi\left(\frac{x_1}{\varepsilon}\right), \quad \text{with} \quad \psi(t) = \int_{\{z_1 \geq -t\}} \varphi(z) dz.$$

Hence  $\psi$  is smooth and decreasing and  $\psi(t) = 0$  if  $t \leq -1$ ,  $\psi(t) = 1$  if  $t \geq 1$ . Set  $R = r/\varepsilon$ . It follows by a change of variables, from Proposition 1, that

$$\frac{1}{2r} \iint_{B_r \times B_r} K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy = \frac{1}{2R} \iint_{B_R \times B_R} \Gamma_\nu(x-y) |\psi(x_1) - \psi(y_1)|^2 dx dy + o(1). \quad (96)$$

Now note that since  $\Gamma_\nu(t) \geq 0$  we have

$$\begin{aligned} \frac{1}{2R} \int_{-R}^R \int_{-R}^R \Gamma_\nu(x-y) dx_2 dy_2 &= \frac{1}{2R} \int_{-2R}^{2R} \Gamma_\nu(x_1 - y_1, z_2) (2R - |z_2|) dz_2 \\ &\leq \int_{-\infty}^{\infty} \Gamma_\nu(x_1 - y_1, z_2) dz_2 = \Gamma_\nu^1(x_1 - y_1) = \frac{\gamma(e_1)}{2|x_1 - y_1|^2}. \end{aligned}$$

Thus using the fact that, for  $t > 0$ ,  $|\psi(x_1) - \psi(x_1 + t)|$  is zero if  $x_1 \leq -t - 1$  or if  $x_1 \geq 1$  and is bounded by  $\text{Lip}\psi t$  if  $t$  is small and by 1 if  $t$  is big, we deduce

$$\begin{aligned} \frac{1}{2R} \iint_{B_R \times B_R} \Gamma_\nu(x-y) |\psi(x_1) - \psi(y_1)|^2 dx dy &\leq \frac{\gamma(e_1)}{2} \int_{-R}^R \int_{-R}^R \frac{1}{|x_1 - y_1|^2} |\psi(x_1) - \psi(y_1)|^2 dx_1 dy_1 \\ &= \gamma(e_1) \int_0^{2R} \frac{1}{t^2} \int_{-R}^{R-t} |\psi(x_1) - \psi(x_1 + t)|^2 dx_1 dt \\ &\leq \gamma(e_1) \int_0^{2R} \frac{1}{t^2} \int_{-t-1}^1 |\psi(x_1) - \psi(x_1 + t)|^2 dx_1 dt \\ &\leq \gamma(e_1) \left( \int_0^1 \frac{(t+2)}{t^2} (\text{Lip}\psi t)^2 dt + \int_1^{2R} \frac{(t+2)}{t^2} dt \right) \\ &\leq \gamma(e_1)(C + \log(2R)). \end{aligned}$$

Since  $\log(2R) = \log(2r) + \log \frac{1}{\varepsilon}$ , from (96) and (89) we finally get

$$\frac{\lambda(B_r(x_0))}{2r} \leq \liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \frac{1}{2r} \iint_{B_r \times B_r} K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \leq \gamma(e_1),$$

which concludes the proof. ○

**Remark 25** Note that in the construction of the optimal sequence for the  $\Gamma$ -limit (Theorem 24) the precise shape of the profile  $\varphi$  is irrelevant. It does not influence the logarithmic contribution of the  $H^{\frac{1}{2}}$  norm of  $u_\varepsilon$  but only the terms which are of order one.

**Remark 26** Together with the lower bound (Theorem 17) and (89), we see that a sequence constructed by convolution as in the above proof is also optimal on any open subset  $A$  of  $Q$  and, up to a subsequence, satisfies

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_A \int_A K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy = \int_{S_u \cap A} \gamma(n) |[u]| d\mathcal{H}^1.$$

In particular we proved that the measure  $\lambda$  defined in (74) satisfies

$$\lambda = \gamma(n) d\mathcal{H}^1 \llcorner S_u.$$

With the theorem below we will give the upper bound for the  $\Gamma$ -convergence result stated in Theorem 10.

**Theorem 27** *Assume that  $N_\varepsilon \varepsilon / |\log \varepsilon| \rightarrow \Lambda$  and that the discs  $B_{R\varepsilon}^i$  are uniformly distributed and well separated. For every  $u \in BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$  there exists a sequence  $\{v_\varepsilon\}$  converging to  $u$  in  $L^q(T^2)$ , for any  $q < 2$ , such that  $v_\varepsilon \in H^{\frac{1}{2}}(T^2)$ ,  $v_\varepsilon = 0$  a.e. on  $\cup_i B_{R\varepsilon}^i$  and*

$$\limsup_{\varepsilon \rightarrow 0} F_\varepsilon(v_\varepsilon) \leq \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1 + \int_Q \Lambda D_\nu(u, B_R) dx. \quad (97)$$

**Proof.** The general idea of the proof is to consider the optimal sequence obtained by convolution in Theorem 24 and to modify it in order to let it satisfy the pinning condition using the  $H^{\frac{1}{2}}$ -dislocation capacitary potentials as cut-off functions, as we did for the dilute case in [8]. However the presence of the non local term in the energy makes this argument more involved. In the proof we will always assume that the fundamental domain  $Q$  for  $T^2$  is chosen such that  $|Du|(\partial Q) = 0$ .

*Step 1. The statement of the theorem holds for any  $u$  constant,  $u = a \in \mathbf{Z}$ .*

In this case the sequence  $v_\varepsilon$  can be constructed exactly as in [8] using the  $H^{\frac{1}{2}}$ -dislocation capacitary potentials (see Section 2.2 for the relevant definitions). We give here some details in order to fix some notation for the following steps. Denote by  $\zeta_\varepsilon^a$  the  $H^{\frac{1}{2}}$ -dislocation capacitary potential of  $B_R$  with respect to  $B_{\varepsilon^{\alpha-1}}$  at the level  $a$ , where  $\alpha \in (1, \frac{\beta+1}{2})$  and  $\beta$  is given by (15). Then the sequence  $v_\varepsilon = v_\varepsilon^a$  is given by

$$v_\varepsilon^a(x) = \begin{cases} a - \zeta_\varepsilon^a \left( \frac{x - x_\varepsilon^i}{\varepsilon} \right) & \text{if } x \in \cup_i B_{\varepsilon^\alpha}^i \\ a & \text{otherwise} \end{cases}.$$

By  $\mu_\varepsilon^a$  and  $\eta_\varepsilon^a$  we denote the following measures

$$\mu_\varepsilon^a(A \times B) = \frac{1}{|\log \varepsilon|} \int_A \int_B K_\nu(x-y) |v_\varepsilon^a(x) - v_\varepsilon^a(y)|^2 dx dy \quad (98)$$

and

$$\eta_\varepsilon^a(A) = \frac{1}{\varepsilon |\log \varepsilon|} \int_A \text{dist}(v_\varepsilon^a, \mathbf{Z}) dx, \quad (99)$$

by  $\mu^a$  and  $\eta^a$  we denote their weak\*-limit (which exist up to a subsequence) and by  $\lambda^a$  we denote the measure such that  $\lambda^a(A) = \mu^a(A \times A)$ . With this notation what we have to prove is  $\lambda^a + \eta^a \leq \Lambda D_\nu(a, B_R) dx$ .

It is easy to check, by using the properties of the kernel  $K_\nu(t)$  (see Proposition 1) and a rescaling argument that

$$\mu_\varepsilon^a(B_{\varepsilon^\alpha} \times B_{\varepsilon^\alpha}) + \eta_\varepsilon^a(B_{\varepsilon^\alpha}) \leq \frac{\varepsilon}{|\log \varepsilon|} D_\nu(a, B_R, B_{\varepsilon^{\alpha-1}}). \quad (100)$$

The choice of the exponent  $\alpha$  is motivated by the following argument that will permit us to control the non local term in the energy and to show that long range interactions are negligible, i.e.

$$\mu_\varepsilon^\alpha((Q \times Q) \cap \{|x - y| > \varepsilon^\beta\}) \leq C a^2 \varepsilon^{(2\alpha - \beta - 1)} |Q|. \quad (101)$$

As a consequence, in view of (100) and (20), we get

$$\mu_\varepsilon^\alpha(Q \times Q) + \eta_\varepsilon^\alpha(Q) \leq \sum_{I_\varepsilon(Q)} \mu_\varepsilon^\alpha(B_{\varepsilon^\alpha}^i \times B_{\varepsilon^\alpha}^i) + \eta_\varepsilon^\alpha(B_{\varepsilon^\alpha}^i) + o(1) \leq \Lambda D_\nu(a, B_R) + o(1)$$

as  $\varepsilon$  goes to zero. Once (101) is shown, Step 1 is finished (see [8] for more details).

It only remains to prove (101). Indeed we will prove the following more general statement which will be useful later.

*There exists a constant  $C > 0$  with the following property. If  $\{w_\varepsilon\}$  and  $\{z_\varepsilon\}$  are two sequences in  $H^{\frac{1}{2}}(Q)$  which are bounded in  $L^\infty$  by a constant  $M$  and satisfy  $w_\varepsilon(x) = z_\varepsilon(x) = \text{const.}$  in  $Q \setminus \cup_i B_{\varepsilon^\alpha}^i$ , then for every  $r > 0$*

$$\frac{1}{|\log \varepsilon|} \iint_{\substack{Q_r \times Q_r \\ |x-y| > \varepsilon^\beta}} K_\nu(x-y) |w_\varepsilon(x) - z_\varepsilon(y)|^2 dx dy \leq C M^2 \varepsilon^{(2\alpha - \beta - 1)} |Q_r|. \quad (102)$$

To show (102) it is enough to use the properties of the kernel  $K_\nu$  and the uniform distribution of the obstacles (cfr. (14)). We have

$$\begin{aligned} \frac{1}{|\log \varepsilon|} \iint_{\substack{Q_r \times Q_r \\ |x-y| > \varepsilon^\beta}} K_\nu(x-y) |w_\varepsilon(x) - z_\varepsilon(y)|^2 dx dy \\ \leq 2 \frac{1}{|\log \varepsilon|} \sum_i \int_{B_{\varepsilon^\alpha}^i} \int_{Q_r} \chi_{|x-y| > \varepsilon^\beta} K_\nu(x-y) |w_\varepsilon(x) - z_\varepsilon(y)|^2 dx dy \\ \leq \frac{1}{|\log \varepsilon|} C 8 M^2 \# \mathcal{I}_\varepsilon(Q_r) \varepsilon^{2\alpha} \int_{B_{4r} \setminus B_{\varepsilon^\beta}} \frac{1}{|y|^3} dy \\ \leq C M^2 \frac{N_\varepsilon |Q_r| \varepsilon^{2\alpha}}{|\log \varepsilon| \varepsilon^\beta} \leq C M^2 \varepsilon^{(2\alpha - \beta - 1)} |Q_r|. \end{aligned}$$

*Step 2. Let  $u \in BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$ , with  $S_u$  polygonal with a finite number of sides. We will prove that for every  $\delta > 0$  there exists a sequence  $v_\varepsilon^\delta$  converging to  $u$  strongly in  $L^q$ , for  $q < 2$ , such that*

$$\limsup_{\varepsilon \rightarrow 0} F_\varepsilon(v_\varepsilon^\delta) \leq \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1 + \int_{T^2} (1 - \chi_{\mathcal{N}_\delta}) D_\nu(u(x), B_R) + \chi_{\mathcal{N}_\delta} |u(x)|^2 D_\nu(1, B_R) dx, \quad (103)$$

where  $\chi_{\mathcal{N}_\delta}$  denotes the characteristic function of the  $\delta$ -neighbourhood of  $S_u$ .

By our choice of  $u$  there exist  $N$  integer number  $a_i$ ,  $i = 1, \dots, N$ , and  $N$  polygons  $P_i$  such that

$$u = \sum_{i=1}^N a_i \chi_{P_i}.$$

From Theorem 24 we find a sequence  $u_\varepsilon$ , obtained by convolution, such that  $u_\varepsilon = u$  in  $Q \setminus \mathcal{N}_\varepsilon$ , where  $\mathcal{N}_\varepsilon$  denotes an  $\varepsilon$ -neighbourhood of  $S_u$ , and

$$\lim_{\varepsilon \rightarrow 0} \mu_\varepsilon(Q \times Q) = \lim_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \iint_{Q \times Q} K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy = \int_{S_u} \gamma(n) |[u]| d\mathcal{H}^1, .$$

We must modify  $u_\varepsilon$  in order to let it satisfy the pinning condition. Thus for any point  $x_\varepsilon^j$  which is not in  $\mathcal{N}_\delta$  we modify  $u_\varepsilon$  exactly as in the previous step using the  $H^{\frac{1}{2}}$ -dislocation capacity potential. If

$x_\varepsilon^j \in \mathcal{N}_\delta$  we modify  $u_\varepsilon$  by multiplying it by an appropriate cut-off function, namely the scaled  $H^{\frac{1}{2}}$ -dislocation capacity potential at level 1. This permits in particular to achieve the pinning condition in  $\mathcal{N}_\varepsilon$  where  $u_\varepsilon$  is not constant. Precisely we define the function  $w_\varepsilon^\delta$  as follows

$$w_\varepsilon^\delta(x) = \begin{cases} a_i - \zeta_\varepsilon^{a_i} \left( \frac{x - x_\varepsilon^j}{\varepsilon} \right) & \text{if } x \in B_{\varepsilon^\alpha}(x_\varepsilon^j) \text{ with } x_\varepsilon^j \in P_i \setminus \mathcal{N}_\delta \\ u_\varepsilon(x) \left( 1 - \zeta_\varepsilon^1 \left( \frac{x - x_\varepsilon^j}{\varepsilon} \right) \right) & \text{if } x \in B_{\varepsilon^\alpha}(x_\varepsilon^j) \text{ with } x_\varepsilon^j \in \mathcal{N}_\delta \\ u_\varepsilon(x) & \text{otherwise.} \end{cases} \quad (104)$$

Let us define again, as above, by  $\tilde{\mu}_\varepsilon$  and  $\tilde{\eta}_\varepsilon$  the following measures

$$\tilde{\mu}_\varepsilon(A \times B) = \frac{1}{|\log \varepsilon|} \int_A \int_B K_\nu(x-y) |w_\varepsilon^\delta(x) - w_\varepsilon^\delta(y)|^2 dx dy \quad (105)$$

and

$$\tilde{\eta}_\varepsilon(A) = \frac{1}{\varepsilon |\log \varepsilon|} \int_A \text{dist}(w_\varepsilon^\delta, \mathbf{Z}) dx, \quad (106)$$

and let us denote by  $\tilde{\mu}$  and  $\tilde{\eta}$  their weak\*-limits and by  $\tilde{\lambda}$  the measure such that  $\tilde{\lambda}(A) = \tilde{\mu}(A \times A)$ . For the sake of simplicity we drop the dependence on  $\delta$  in the notation for the measures introduced above. We shall prove that

$$\tilde{\lambda} + \tilde{\mu} \leq \Lambda[(1 - \chi_{\mathcal{N}_\delta}(x))D_\nu(u(x), B_R) + \chi_{\mathcal{N}_\delta}(x)|u(x)|^2 D_\nu(1, B_R)]dx + \gamma(n)|[u](x)d\mathcal{H}^1 \llcorner S_u. \quad (107)$$

This clearly implies the assertion of Step 2.

First we prove that  $\tilde{\lambda} + \tilde{\mu}$  is absolutely continuous with respect to the Lebesgue measure outside  $S_u$ . Fix  $x_0 \in Q \setminus S_u$ . Since for any  $a \in \mathbf{Z}$  and  $v \in \mathbf{R}$  we have that  $\text{dist}^2(av, \mathbf{Z}) \leq a^2 \text{dist}^2(v, \mathbf{Z})$ , by the minimality of the  $H^{\frac{1}{2}}$ -dislocation capacity potentials it is easy to check that

$$D_\nu(a, B_R) \leq a^2 D_\nu(1, B_R)$$

and for all  $x_\varepsilon^i \in Q_r(x_0)$  we have

$$\tilde{\mu}_\varepsilon(B_{\varepsilon^\alpha}^i \times B_{\varepsilon^\alpha}^i) + \tilde{\eta}_\varepsilon(B_{\varepsilon^\alpha}^i) \leq |u(x_0)|^2 (\mu_\varepsilon^1(B_{\varepsilon^\alpha}^i \times B_{\varepsilon^\alpha}^i) + \eta_\varepsilon^1(B_{\varepsilon^\alpha}^i))$$

where  $\mu_\varepsilon^1$  and  $\eta_\varepsilon^1$  have been defined in Step 1 (take  $a = 1$ ). Thus in view of (102), applied also for  $w_\varepsilon = v_\varepsilon^a$  and  $z_\varepsilon = av_\varepsilon^1$ , we have

$$\tilde{\mu}_\varepsilon(Q_r(x_0) \times Q_r(x_0)) + \tilde{\eta}_\varepsilon(Q_r(x_0)) \leq |u(x_0)|^2 (\mu_\varepsilon^1(Q_r(x_0) \times Q_r(x_0)) + \eta_\varepsilon^1(Q_r(x_0))) + o(1).$$

Then taking the limit as  $\varepsilon$  goes to zero and using Step 1 we see that  $\tilde{\lambda} + \tilde{\eta}$  is absolutely continuous with respect to the Lebesgue measure outside  $S_u$  and satisfies (107) in  $\mathcal{N}_\delta \setminus S_u$ . In the case  $x_0 \in Q \setminus \mathcal{N}_\delta$  we shall prove that

$$\tilde{\lambda}(Q_r(x_0)) + \tilde{\eta}(Q_r(x_0)) \leq \Lambda D_\nu(u(x_0), B_R) |Q_r| + o(1) \quad (108)$$

as  $r$  goes to zero. This clearly follows as above by Step 1 and the fact that  $v_\varepsilon$  coincide with  $v_\varepsilon^a$  in a neighbourhood of  $x_0$ , where  $a = u(x_0)$ . For  $r$  small enough

$$\tilde{\mu}_\varepsilon(Q_r(x_0) \times Q_r(x_0)) + \tilde{\eta}_\varepsilon(Q_r(x_0)) = \mu_\varepsilon^a(Q_r(x_0) \times Q_r(x_0)) + \eta_\varepsilon^a(Q_r(x_0)).$$

Thus outside  $S_u$  inequality (107) follows from Step 1.

Let now consider  $x_0 \in S_u$ . We will prove that

$$\limsup_{r \rightarrow 0} \frac{\tilde{\lambda}(Q_r(x_0)) + \tilde{\eta}(Q_r(x_0))}{r} \leq \gamma(n)|[u](x_0)|. \quad (109)$$

It is easy to see that

$$\lim_{r \rightarrow 0} \frac{\tilde{\eta}(Q_r(x_0))}{r} = 0.$$

Indeed, using a change of variable and the definition of  $v_\varepsilon$ , in particular the fact that  $u_\varepsilon$  is constant in  $Q \setminus \mathcal{N}_\varepsilon$ , for any  $r < \delta$ , we get

$$\begin{aligned} \tilde{\eta}_\varepsilon(Q_r(x_0)) &\leq \tilde{\eta}_\varepsilon(Q_r(x_0) \cap \mathcal{N}_\varepsilon) + \tilde{\eta}_\varepsilon(Q_r(x_0) \setminus \mathcal{N}_\varepsilon) \\ &\leq \frac{C\varepsilon 2r}{\varepsilon |\log \varepsilon|} + \frac{1}{\varepsilon |\log \varepsilon|} \#\mathcal{I}_\varepsilon(Q_r) \varepsilon^2 \int_{B_{\varepsilon^{\alpha-1}}} \text{dist}^2(\zeta_\varepsilon^a, \mathbf{Z}) dx \leq \Lambda D_\nu(a, B_R) |Q_r| + o(1), \end{aligned}$$

as  $\varepsilon$  goes to zero, where we denoted by  $a$  the maximum of  $|u|$  in  $Q_r$ . Thus in order to prove (109) it is enough to show that

$$\limsup_{r \rightarrow 0} \frac{\tilde{\lambda}(Q_r(x_0))}{r} \leq \gamma(n) |[u](x_0)|. \quad (110)$$

For  $r < \delta$  we have  $Q_r \subseteq \mathcal{N}_\delta$ . Since by definition  $w_\varepsilon^\delta(x) = u_\varepsilon(x) v_\varepsilon^1(x)$  in  $\mathcal{N}_\delta$  we have, for every  $\sigma > 0$ ,

$$\begin{aligned} \tilde{\mu}_\varepsilon(Q_r \times Q_r) &= \frac{1}{|\log \varepsilon|} \iint_{Q_r \times Q_r} K_\nu(x-y) |v_\varepsilon^1(x) u_\varepsilon(x) - v_\varepsilon^1(x) u_\varepsilon(y) + v_\varepsilon^1(x) u_\varepsilon(y) - v_\varepsilon^1(y) u_\varepsilon(y)|^2 dx dy \\ &\leq \frac{1}{|\log \varepsilon|} (1 + \sigma) \iint_{Q_r \times Q_r} K_\nu(x-y) |v_\varepsilon^1(x)|^2 |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \\ &\quad + \frac{1}{|\log \varepsilon|} \left(1 + \frac{1}{\sigma}\right) \iint_{Q_r \times Q_r} K_\nu(x-y) |u_\varepsilon(y)|^2 |v_\varepsilon^1(x) - v_\varepsilon^1(y)|^2 dx dy \\ &\leq (1 + \sigma) \mu_\varepsilon(Q_r \times Q_r) + \sup_Q |u|^2 \left(1 + \frac{1}{\sigma}\right) \mu_\varepsilon^1(Q_r \times Q_r), \end{aligned}$$

where in the last inequality we used the fact that  $v_\varepsilon^1$  is bounded by 1 and  $\sup |u_\varepsilon| \leq \sup |u|$ . Then (110) follows by the above estimate, taking the limit as  $\varepsilon$  goes to zero. Indeed, in view of Remark 26 and Step 1 we have

$$\frac{1}{r} \lim_{\varepsilon \rightarrow 0} \tilde{\mu}_\varepsilon(Q_r \times Q_r) \leq (1 + \sigma) \frac{1}{r} \int_{Q_r \cap S_u} \gamma(n) |[u]| d\mathcal{H}^1 + \left(1 + \frac{1}{\sigma}\right) \Lambda \sup_Q |u|^2 D_\nu(1, B_R) \frac{|Q_r|}{r}.$$

After taking the limit as  $r$  goes to zero the estimate (110), and hence (107) and (109), follows by the arbitrariness of  $\sigma$ .

*Step 3.* In this step we conclude the proof. First we can obtain the thesis of the theorem for functions  $u \in BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$ , with  $S_u$  polygonal, with a finite number of sides by Step 2 and a diagonalization argument.

By estimate (19) the map  $u \mapsto \int_Q D_\nu(u, B_R) dx$  is continuous in  $L^2(Q, \mathbf{Z})$  and satisfies

$$\int_Q D_\nu(u, B_R) dx \geq C_1 \|u\|_{L^2}^2.$$

This implies that the limit functional is continuous in  $BV(T^2, \mathbf{Z}) \cap L^2(T^2, \mathbf{Z})$ . Hence the general case can be recovered by a density argument.  $\circ$

## 7 The sub-critical and super-critical regimes

In this section we will briefly discuss the asymptotic behaviour of  $E_\varepsilon(u)/(N_\varepsilon \varepsilon)$  in the sub-critical and in the super-critical regime. In the latter case we also assume  $N_\varepsilon \ll \frac{1}{\varepsilon^{2\beta}}$ , with  $\beta < 1$ , in order to keep the obstacles well separated (see condition (15)).

## 7.1 The sub-critical regime

In the case  $N_\varepsilon \ll |\log \varepsilon|/\varepsilon$  the result can be deduced from the Theorem 10. Indeed the compactness result given in Theorem 12 implies compactness in  $L^q(T^2)$ , for  $q < 2$ , of any sequence  $\{u_\varepsilon\}$  such that  $\frac{E_\varepsilon(u_\varepsilon)}{\varepsilon N_\varepsilon} \leq C$ . Moreover every cluster point  $u \in BV(T^2, \mathbf{Z})$  of such a sequence must satisfy  $\int_{T^2} |Du| = 0$  and hence is a constant  $a \in \mathbf{Z}$ . Thus also in this case, as for the dilute case discussed in Theorem 6, the effect of the pinning condition is weaker than the line tension and in a similar way one can deduce that the  $\Gamma$ -limit is given by

$$E(u) = \begin{cases} D_\nu(a, B_R) & \text{if } u = \text{const.} \in \mathbf{Z}, \\ +\infty & \text{otherwise.} \end{cases}$$

## 7.2 The super-critical regime

We now consider the case  $\frac{|\log \varepsilon|}{\varepsilon} \ll N_\varepsilon \ll \frac{1}{\varepsilon^2}$ . It is easy to check that also in this case the Poincaré inequality, as in Proposition 16, yields an  $L^{\frac{2}{3}}(T^2)$  bound for all sequences with equibounded energy, but in general we cannot expect more than weak convergence up to a subsequence.

The upper bound in the proof of Corollary 11 can be obtained for the class of piecewise constant functions with integer values taking into account that in the super-critical scale the line tension effect becomes negligible. This class is weakly dense in  $L^2(T^2, \mathbf{R})$  and then the general case follows by an energy density argument.

The proof of the lower bound is more delicate. By a blow-up argument it is enough to understand the case of a sequence  $\{u_\varepsilon\}$  converging to a real constant  $c$  weakly in  $L^2(T^2)$ . The idea in this case is that even if the sequence converges only weakly, it oscillates at a larger scale than the average distance between the obstacles. Then at this scale, we still can find an integer value to which we can apply Lemma 22 and get a local lower bound for the energy.

**Proposition 28** *If  $u_\varepsilon \rightharpoonup c = \text{const.}$ , with  $c \in \mathbf{R}$ , then*

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon N_\varepsilon} E_\varepsilon(u_\varepsilon) \geq D_\nu^{**}(c, B_R), \quad (111)$$

where  $D_\nu^{**}(\cdot, B_R)$  is the convex envelope of  $D_\nu(\cdot, B_R)$  as defined in (17).

**Proof.** *Step 1 (Selection of good pinning sites  $x_\varepsilon^i$ ).* We may assume that  $\frac{E_\varepsilon(u_\varepsilon)}{\varepsilon N_\varepsilon} \leq C$ . Thus for every  $\theta \in (0, 1)$  we can find a set of indices  $\mathcal{I}_\varepsilon^\theta$  and a constant  $C(\theta)$  such that  $\#\mathcal{I}_\varepsilon^\theta \geq (1 - \theta)N_\varepsilon$  and for all  $i \in \mathcal{I}_\varepsilon^\theta$  the points  $x_\varepsilon^i$  satisfy

$$\frac{1}{\varepsilon} \int_{B_{\rho_\varepsilon}^i} \text{dist}^2(u_\varepsilon, \mathbf{Z}) dx \leq C(\theta)\varepsilon, \quad (112)$$

$$\iint_{B_{\rho_\varepsilon}^i \times B_{\rho_\varepsilon}^i} K_\nu(x - y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \leq C(\theta)\varepsilon \quad (113)$$

and

$$\int_{B_{\rho_\varepsilon}^i} |u_\varepsilon|^2 dx \leq C(\theta), \quad (114)$$

where  $B_{\rho_\varepsilon}^i$  denotes the disc of radius  $\rho_\varepsilon = 1/\sqrt{N_\varepsilon}$  and center  $x_\varepsilon^i$ .

*Step 2 (Assignment of a value  $c_\varepsilon^i$  to each good pinning site).* By the (scaling invariant) embedding of  $H^{\frac{1}{2}}$  into  $L^4$ , there exist constants  $c_\varepsilon^i$  such that

$$\int_{B_{\rho_\varepsilon}^i} |u_\varepsilon - c_\varepsilon^i|^4 dx \leq C[u_\varepsilon]_{H^{\frac{1}{2}}(B_{\rho_\varepsilon}^i)}^4 \leq C\varepsilon^2.$$

Thus

$$\int_{B_{\rho_\varepsilon}^i} |u_\varepsilon - c_\varepsilon^i|^4 dx \leq CN_\varepsilon \varepsilon^2 \rightarrow 0 \quad (115)$$

and

$$\int_{B_{\rho_\varepsilon}^i} |u_\varepsilon - c_\varepsilon^i| dx \leq C(N_\varepsilon \varepsilon^2)^{\frac{1}{4}} \rightarrow 0.$$

By the interpolation inequality, given by Lemma 23, applied to  $u_\varepsilon - c_\varepsilon^i$  we have

$$\int_{B_{\varepsilon^\beta}^i} |u_\varepsilon - c_\varepsilon^i| dx \leq \int_{B_{\rho_\varepsilon}^i} |u_\varepsilon - c_\varepsilon^i| dx + C \frac{\varepsilon^{\frac{1}{2}}}{\varepsilon^{\frac{\beta}{2}}} \rightarrow 0 \quad (116)$$

where  $\beta < 1$  is chosen as in (15). From (115) we also deduce

$$\int_{B_{\rho_\varepsilon}^i} |u_\varepsilon - c_\varepsilon^i|^2 dx \leq C(N_\varepsilon \varepsilon^2)^{\frac{1}{2}} \rightarrow 0.$$

Together with (114) this yields

$$|c_\varepsilon^i| \leq C. \quad (117)$$

*Step 3 (Lower bound for each good pinning site).* We claim that there exist integers  $a_\varepsilon^i$  such that

$$\sup_{i \in \mathcal{I}_\varepsilon^\theta} |c_\varepsilon^i - a_\varepsilon^i| \rightarrow 0 \quad (118)$$

and

$$\frac{1}{\varepsilon^2} \int_{B_{\varepsilon^\beta}^i} \text{dist}^2(u_\varepsilon, \mathbf{Z}) dx + \frac{1}{\varepsilon} \iint_{B_{\varepsilon^\beta}^i \times B_{\varepsilon^\beta}^i} K_\nu(x-y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \geq D_\nu(a_\varepsilon^i, B_R) - o(1). \quad (119)$$

To see this we consider the scaled function  $\zeta_\varepsilon^i(x) = u_\varepsilon(x_\varepsilon^i + \varepsilon x)$ . Then (116) gives

$$\omega_1(\varepsilon) = \int_{B_{\varepsilon^{\beta-1}}} |\zeta_\varepsilon^i - c_\varepsilon^i| dx \rightarrow 0 \quad (120)$$

uniformly in  $i \in \mathcal{I}_\varepsilon^\theta$ , and the left hand side of (119) can be written as

$$T_\varepsilon^i := \int_{B_{\varepsilon^{\beta-1}}} \text{dist}^2(\zeta_\varepsilon^i, \mathbf{Z}) dx + \iint_{B_{\varepsilon^{\beta-1}} \times B_{\varepsilon^{\beta-1}}} K_\nu(x-y) |\zeta_\varepsilon^i(x) - \zeta_\varepsilon^i(y)|^2 dx dy.$$

By (112) and (113) we know that  $T_\varepsilon^i$  is bounded. Moreover by (120) the set

$$\{z \in B_{\varepsilon^{\beta-1}} : |\zeta_\varepsilon^i(z) - c_\varepsilon^i| < 2\omega_1(\varepsilon)\}$$

has measure at least  $|B_{\varepsilon^{\beta-1}}|/2$ . Hence  $T_\varepsilon^i$  can be bounded only if

$$\text{dist}(c_\varepsilon^i, \mathbf{Z}) \leq 2\omega_1(\varepsilon) + \frac{C}{|B_{\varepsilon^{\beta-1}}|^{\frac{1}{2}}} \rightarrow 0$$

This proves (118). In view of (120) we also get

$$\int_{B_{\varepsilon^{\beta-1}}} |\zeta_\varepsilon^i - a_\varepsilon^i| dx \rightarrow 0.$$

Hence Lemma 22 and (117) yield

$$T_\varepsilon^i \geq D_\nu(a_\varepsilon^i, B_R) - o(1).$$

Summation of (119) over the good centers yields

$$\frac{1}{\varepsilon N_\varepsilon} E_\varepsilon(u_\varepsilon) \geq (1 - \theta) \frac{1}{\#(\mathcal{I}_\varepsilon^\theta)} \sum_{i \in \mathcal{I}_\varepsilon^\theta} D_\nu(a_\varepsilon^i, B_R). \quad (121)$$

*Step 4 (Relation between the values  $a_\varepsilon^i$  assigned to the pinning sites and the weak limit  $c$  of  $u_\varepsilon$ ).* By (115) and (118) we see that  $u_\varepsilon$  is close to an (integer) constant on balls of radius  $\frac{1}{\sqrt{N_\varepsilon}}$  centered on ‘good’ pinning sites. We now show that  $u_\varepsilon$  is actually nearly constant on a slightly larger scale  $r_\varepsilon$ . This will allow us to exploit the uniform distribution of the pinning sites to conclude that  $\int_Q u_\varepsilon dx \approx \frac{1}{\#(\mathcal{I}_\varepsilon^\theta)} \sum_{i \in \mathcal{I}_\varepsilon^\theta} a_\varepsilon^i$ . Since  $\int_Q u_\varepsilon dx \rightarrow c$

Since  $N_\varepsilon \ll \varepsilon^{-2}$  there exists  $r_\varepsilon$  such that

$$\frac{1}{\sqrt{N_\varepsilon}} \ll r_\varepsilon \ll \frac{1}{\varepsilon^{\frac{1}{2}} N_\varepsilon^{\frac{3}{4}}}. \quad (122)$$

We assume that  $\frac{1}{r_\varepsilon}$  is an integer and we cover  $Q$  by a lattice of squares  $\tilde{Q}_{r_\varepsilon}^j$  of size  $r_\varepsilon$ . By  $\hat{Q}_{r_\varepsilon}^j$  we denote the concentric squares of three times the size. Given  $\theta$  we can find a set of indices  $\mathcal{J}_\varepsilon^\theta$  such that  $\#(\mathcal{J}_\varepsilon^\theta) \geq (1 - \theta)r_\varepsilon^{-2}$  and for all  $j \in \mathcal{J}_\varepsilon^\theta$  the squares  $\tilde{Q}_{r_\varepsilon}^j$  satisfy

$$\int \int_{\hat{Q}_{r_\varepsilon}^j \times \hat{Q}_{r_\varepsilon}^j} K_\nu(x - y) |u_\varepsilon(x) - u_\varepsilon(y)|^2 dx dy \leq C(\theta) \varepsilon N_\varepsilon r_\varepsilon^2, \quad (123)$$

$$\int_{\hat{Q}_{r_\varepsilon}^j} |u_\varepsilon|^2 dx \leq C(\theta) r_\varepsilon^2. \quad (124)$$

By the embedding of  $H^{\frac{1}{2}}$  into  $L^4$ , there exist constants  $A_\varepsilon^j$  such that

$$\int_{\tilde{Q}_{r_\varepsilon}^j} |u_\varepsilon - A_\varepsilon^j|^4 dx \leq C \varepsilon^2 N_\varepsilon^2 r_\varepsilon^2 \rightarrow 0.$$

Now consider a good pinning site in a good square, i.e.  $x_\varepsilon^i \in \tilde{Q}_{r_\varepsilon}^j$ ,  $i \in \mathcal{I}_\varepsilon^\theta$  and  $j \in \mathcal{J}_\varepsilon^\theta$ . Then by the interpolation inequality (applied to  $u_\varepsilon - A_\varepsilon^j$ ) and (122)

$$\int_{B_{\rho_\varepsilon}^i} |u_\varepsilon - A_\varepsilon^j| dx \leq \int_{\tilde{Q}_{r_\varepsilon}^j} |u_\varepsilon - A_\varepsilon^j| dx + C \frac{\varepsilon^{\frac{1}{2}} N_\varepsilon^{\frac{1}{2}} r_\varepsilon}{N_\varepsilon^{-\frac{1}{4}}} \leq C o(1).$$

Thus in view of (115) and (118)

$$|a_\varepsilon^i - A_\varepsilon^j| \leq C o(1).$$

Since  $a_\varepsilon^i \in \mathbf{Z}$  this shows that for good pinning sites in good squares  $a_\varepsilon^i$  depends only on the square  $\tilde{Q}_{r_\varepsilon}^j$ .

Using the uniform distribution of  $x_\varepsilon^i$  and the fact that  $r_\varepsilon \gg N_\varepsilon^{-\frac{1}{2}}$  as well as (114) we deduce that

$$\left| \frac{1}{N_\varepsilon} \sum_{i \in \mathcal{I}_\varepsilon^\theta, x_\varepsilon^i \in \tilde{Q}_{r_\varepsilon}^j} a_\varepsilon^i - \int_{\tilde{Q}_{r_\varepsilon}^j} u_\varepsilon dx \right| \leq C |\tilde{Q}_{r_\varepsilon}^j| o(1) + \frac{\#(\mathcal{I}_\varepsilon(\tilde{Q}_{r_\varepsilon}^j) \setminus \mathcal{I}_\varepsilon^\theta)}{N_\varepsilon}. \quad (125)$$

Now we sum (125) over all  $j \in \mathcal{J}_\varepsilon^\theta$ . Let  $E_\varepsilon$  be the union of the squares  $\tilde{Q}_{r_\varepsilon}^j$ , with  $j \notin \mathcal{J}_\varepsilon^\theta$ . Then  $|E_\varepsilon| \leq \theta$  and by the  $L^2$  bound on  $u_\varepsilon$  we have  $\int_{E_\varepsilon} |u_\varepsilon| dx \leq C \theta^{\frac{1}{2}}$ . Combining this with (117) and (118) we deduce after a short calculation that

$$\lim_{\theta \rightarrow 0} \lim_{\varepsilon \rightarrow 0} \left| \frac{1}{\#(\mathcal{I}_\varepsilon^\theta)} \sum_{i \in \mathcal{I}_\varepsilon^\theta} a_\varepsilon^i - \int_Q u_\varepsilon dx \right| = 0$$

Together with (121) this finishes the proof. ○

## References

- [1] ALBERTI, G., AND BELLETTINI, G. A non-local anisotropic model for phase transitions: asymptotic behaviour of rescaled energies. *European J. Appl. Math.* 9, 3 (1998), 261–284.
- [2] ALBERTI, G., AND BELLETTINI, G. A nonlocal anisotropic model for phase transitions. I. the optimal profile problem. *Math. Ann.* 310, 3 (1998), 527–560.
- [3] ALBERTI, G., BOUCHITTÉ, G., AND SEPPECHÉ, P. Phase transition with the line-tension effect. *Arch. Rational Mech. Anal.* 144, 1 (1998), 1–46.
- [4] CIORANESCU, D., AND MURAT, F. Un terme trange venu d’ailleurs. (french) [a strange term brought from somewhere else]. In *Nonlinear partial differential equations and their applications. Collge de France Seminar Vol. II (Paris, 1979/1980)*, Res. Notes in Math., 60. Pitman, Boston, Mass.-London, 1982, pp. 98–138, 389–390.
- [5] CONTI, S., FONSECA, I., AND LEONI, G. A  $\Gamma$ -convergence result for the two-gradient theory of phase transitions. *Comm. Pure Appl. Math.* 55 (2002), 857–936.
- [6] DAL MASO, G., AND GARRONI, A. New results on the asymptotic behaviour of dirichlet problems in perforated domains. *Math. Models Methods Appl. Sci.* 3 (1994), 373–407.
- [7] FONSECA, I., AND TARTAR, L. The gradient theory of phase transitions. *Proc. Roy. Soc. Edinburgh Sect. A* 111 (1989), 89–102.
- [8] GARRONI, A., AND MÜLLER, S.  $\Gamma$ -limit of a phase-field model of dislocations. Preprint 92/2003 Max Planck Institute, Leipzig, to appear in *SIAM J. Math. Anal.*
- [9] KOHN, R. V., AND STERNBERG, P. Local minimizers and singular perturbations. *Proc. Roy. Soc. Edinburgh Sect. A* 111 (1989), 69–84.
- [10] KOSŁOWSKI, M., CUITINO, A. M., AND ORTIZ, M. A phase-field theory of dislocation dynamics, strain hardening and hysteresis in ductile single crystal. *J. Mech. Phys. Solids* 50 (2002), 2597–2635.
- [11] KURZKE, M. A nonlocal singular perturbation problem with periodic well potential. Preprint 106/2003 MPI-MIS, Leipzig.
- [12] MARCHENKO, A., AND KRUSLOV, E. Y. New results of boundary value problems for regions with closed-grained boundaries. *Uspekhi Mat. Nauk* 33, 127 (1978).
- [13] MODICA, L. The gradient theory of phase transitions and the minimal interface criterion. *Arch. Rational Mech. Anal.* 98 (1987), 123–142.
- [14] MODICA, L., AND MORTOLA, S. Il limite nella  $\Gamma$ -convergenza di una famiglia di funzionali ellittici. (italian). *Boll. Un. Mat. Ital. A* 14, 3 (1977), 526–529.
- [15] MODICA, L., AND MORTOLA, S. Un esempio di  $\Gamma$ -convergenza. (italian). *Boll. Un. Mat. Ital. B* 14, 1 (1977), 285–299.