

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

On the gradient set of Lipschitz maps

by

Bernd Kirchheim, and László Székelyhidi

Preprint no.: 16

2007



ON THE GRADIENT SET OF LIPSCHITZ MAPS

BERND KIRCHHEIM AND LÁSZLÓ SZÉKELYHIDI JR.

ABSTRACT. We prove that the essential range of the gradient of planar Lipschitz maps has a connected rank-one convex hull. As a corollary, in combination with the results in [7] we obtain a complete characterization of incompatible sets of gradients for planar maps in terms of rank-one convexity.

1. INTRODUCTION

This paper is concerned with the range of gradients of Lipschitz maps. More precisely, let $\Omega \subset \mathbb{R}^n$ be a bounded domain, and let $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a Lipschitz map. We denote by $[Du]$ the essential range of the gradient of u , i.e. the smallest closed subset of $\mathbb{R}^{m \times n}$ such that $Du(x) \in [Du]$ for almost every $x \in \Omega$. Our aim is to find geometric restrictions on, or characterizations of the essential range of gradients of Lipschitz maps.

This issue plays a central role in the study of material microstructure [2, 3, 5, 9], and is linked to the question of existence and regularity of solutions to partial differential inclusions of the type

$$Du(x) \in K \text{ a.e. } x \in \Omega,$$

where $K \subset \mathbb{R}^{m \times n}$ is a prescribed (compact) set of matrices.

The following construction is well known: let $A, B \in \mathbb{R}^{m \times n}$ be two matrices such that $\text{rank}(A - B) = 1$, so that $A - B = a \otimes \nu$ for some vectors $a \in \mathbb{R}^m$ and $\nu \in \mathbb{R}^n$. For any Lipschitz “profile” $h : \mathbb{R} \rightarrow \mathbb{R}$ with $h'(t) \in \{0, 1\}$ a.e., the map

$$u(x) = Bx + ah(x \cdot \nu)$$

is a Lipschitz map whose gradient takes the values A or B almost everywhere. This type of example is called a *simple laminate*, and whenever two matrices A, B satisfy $\text{rank}(A - B) = 1$, one speaks of a *rank-one connection* (or, more classically, A and B are said to satisfy the Hadamard jump condition). On the other hand, it is also well known that if $A, B \in \mathbb{R}^{m \times n}$ with $\text{rank}(A - B) > 1$, then the only Lipschitz maps with gradient $Du(x) \in \{A, B\}$ a.e. are affine maps. Moreover, in [2] J. M. Ball and R. D. James established the much stronger statement that whenever $\{u_j\}$ is a sequence of maps bounded in $W^{1,1}$ such that $\text{dist}(Du_j, \{A, B\}) \rightarrow 0$ in L^1 strongly, then - up to a subsequence - $Du_j \rightarrow A$ or $Du_j \rightarrow B$ strongly in L^1 .

A general question, that has received considerable attention recently, is to understand to what extent the above construction is universal. In other

words to understand to what extent the presence of rank-one connections is necessary in the essential range of gradients of Lipschitz maps. To put this question into proper perspective, we need to recall the example given by the first author together with D. Preiss of a Lipschitz map $u : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$, where $[Du]$ consists of 5 matrices, none of which are rank-one connected to each other ([12, Chapter 4], see also [11] for similar examples). Such examples show that it may happen that the set $[Du]$ itself contains no rank-one connections. On the other hand, the construction of such maps relies very much on the presence of rank-one segments in the sense that it proceeds via an iteration scheme known as convex integration (see [13] for a survey of the theory). In technical terms one key ingredient for this scheme to work is that the rank-one convex hull $[Du]^{rc}$ is a connected set, which contains many rank-one segments in the sense that for any matrix $A \in [Du]^{rc} \setminus [Du]$ there exists a rank-one segment through A contained in $[Du]^{rc}$. The upshot is that although the iterative process of convex integration can eliminate rank-one connections in the essential range $[Du]$ of the limit, the “trail” it leaves behind is a large rank-one convex hull (see Section 4 for a precise definition of the rank-one convex hull).

Our main result shows that for planar maps this is in some sense optimal:

THEOREM 1. *Let $\Omega \subset \mathbb{R}^2$ be a bounded domain, and $u : \Omega \rightarrow \mathbb{R}^2$ a Lipschitz map. Then the rank-one convex hull $[Du]^{rc}$ of the essential range of the gradient is connected.*

It is important to note that connectedness itself does not imply that $[Du]^{rc}$ contains rank-one segments. The standard example is simply a planar conformal map. However, in some sense this is the only example. Indeed, if $[Du]^{rc}$ is connected and contains no rank-one connections, then in fact the differential inclusion

$$Du(x) \in [Du]^{rc} \text{ a.e. } x \in \Omega$$

can be viewed as a (possibly degenerate) elliptic system (see [23, 25]). In particular, we have the following statement.

COROLLARY 1. *If the essential range of the gradient of a Lipschitz map $u : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^2$ contains an isolated matrix $A \in [Du]$, then there exists another matrix $B \in [Du]^{rc} \setminus \{A\}$ such that $\text{rank}(A - B) = 1$.*

We emphasize that Theorem 1 and Corollary 1 are very specific for *planar* mappings, and the analogue statements are false in higher dimensions in general (see for example [9]).

Our Theorem 1 has interesting implications concerning the study of *incompatible sets* of gradients. In combination with the results in [6] and [7] we obtain the following theorem.

THEOREM 2. *Let $K_1, K_2 \subset \mathbb{R}^{2 \times 2}$ be disjoint compact sets which are rank-one incompatible in the sense that*

$$K_1^{rc} \cap K_2^{rc} = \emptyset \text{ and } K_1^{rc} \cup K_2^{rc} = (K_1 \cup K_2)^{rc}.$$

Then for any bounded Lipschitz domain $\Omega \subset \mathbb{R}^2$ and any $p \in [1, \infty)$ there exists a constant $C = C(p, \Omega)$ such that

$$\min\left(\int_{\Omega} \text{dist}^p(Du, K_1), \int_{\Omega} \text{dist}^p(Du, K_2)\right) \leq C \int_{\Omega} \text{dist}^p(Du, K_1 \cup K_2)$$

for all $u \in W^{1,p}(\Omega, \mathbb{R}^2)$.

In the proof of Theorem 1 we follow the approach of [7], which is based on the geometric characterization of rank-one incompatibility via a separating curve, introduced by the second author in [24]. As in the proof of [7, Theorem 4], the existence of a separating curve $\Gamma : \mathcal{S}^1 \rightarrow \mathbb{R}^{2 \times 2}$ allows us to consider the associated family of quasiregular mappings

$$u^t(x) = u(x) - \Gamma(t)x.$$

However, since in our case we have no control over the boundary values of u^t , we cannot conclude that these maps are homeomorphisms. In general they may have branch points. Our strategy is to prove that the set of branch points is in fact independent of $t \in \mathcal{S}^1$, and therefore cannot disconnect Ω . In this way we will be able to conclude the incompatibility just as in [7, Theorem 4].

Our paper is organized as follows. In Section 2 we prove a general result about stability of the branch set of quasiregular mappings in \mathbb{R}^n , and show in Proposition 2 how it can be used to prove separation results for gradients of Lipschitz maps in \mathbb{R}^n . Then in Section 3 we utilize the stability result together with the existence of a separating curve in case $[Du]^{rc}$ is disconnected (as in [24]) to prove Theorem 1 and Corollary 1.

Finally, in Section 4 we discuss the implications of Theorem 1 to the study of incompatible sets of gradients and in particular the proof of Theorem 2. As the explanation of these implications requires introducing the language of gradient Young measures which does not otherwise play a central role in our paper, we defer the statements and proofs until that section.

2. STABILITY OF THE BRANCH SET

In the following, given a domain $\Omega \subset \mathbb{R}^n$ and an open and discrete mapping $u : \Omega \rightarrow \mathbb{R}^n$, we shall write $\mu(y, u, G)$ for the *local degree* of the mapping at $y \in \mathbb{R}^n$ with respect to G (provided $y \notin u(\partial G)$), $N(y, u, G) = \text{card } u^{-1}(y) \cap G$, $N(u, G) = \sup_y N(y, u, G)$ and $i(x, u)$ for the *local index* of u at $x \in \Omega$. We recall that a domain $D \subset \Omega$ is called a *normal domain* for the mapping u if $u(\partial D) = \partial u(D)$ (note that $\partial u(D) \subset u(\partial D)$ follows automatically from openness of the map). A *normal neighbourhood* D of $x \in \Omega$ is a normal domain such that $D \cap u^{-1}(u(x)) = \{x\}$.

A map $u : \Omega \rightarrow \mathbb{R}^n$ is said to be *quasiregular*, if for some constant $\mathcal{K} \geq 1$

$$\|Du(x)\|^n \leq \mathcal{K} \det Du(x) \quad \text{a.e. } x \in \Omega,$$

where $\|Du(x)\|$ denotes the operator norm of the matrix $Du(x)$. It is well known since the pioneering work of Y. G. Reshetnyak that non-constant

quasiregular mappings are open and discrete. The branch set $B(u)$ is defined as the set of points $x \in \Omega$ where u is not locally homeomorphic, that is,

$$B(u) := \{x \in \Omega : i(x, u) > 1\}.$$

In particular for quasiregular maps $B(u)$ is a closed set of topological dimension $(n - 2)$ and Lebesgue measure zero [20]. For the basic theory of quasiregular mappings, and their topological properties, we refer the reader to [21].

PROPOSITION 1. *Let $u : \Omega \rightarrow \mathbb{R}^n$ be a \mathcal{K} -quasiregular mapping such that $\|Du(x)\| \geq \varepsilon$ a.e. $x \in \Omega$, let $G \subset \Omega$ be a subdomain with $\bar{G} \subset \Omega$, and assume that $M := N(u, G) < \infty$.*

There exists a constant $\delta = \delta(\varepsilon, \mathcal{K}, M, n) > 0$ and for each $x_0 \in G$ a radius $r(x_0) > 0$ so that for any Lipschitz mapping $\phi : \Omega \rightarrow \mathbb{R}^n$ with $\|D\phi(x)\| < \delta$ a.e. $x \in \Omega$,

$$(1) \quad \min_{|x-x_0|=r} |u^t(x) - u^t(x_0)| \geq \delta r \quad \text{for all } r < r(x_0), t \in [0, 1],$$

where $u^t = u + t\phi$. In particular

$$i(x_0, u) = i(x_0, u + \phi) \text{ for all } x_0 \in G,$$

and $B(u) \cap G = B(u + \phi) \cap G$.

PROOF. From [19] for every $x_0 \in G$ there exists a radius $r(x_0) > 0$ so that $B_{r(x_0)}(x_0) \subset \Omega$, and for $r < r(x_0)$

$$\max_{|x-x_0|=r} |u(x) - u(x_0)| \leq L \min_{|x-x_0|=r} |u(x) - u(x_0)|,$$

where $L = L(\mathcal{K}, M, n)$. Moreover,

$$\int_{B_r(x_0)} \det Du(x) dx = \int_{\mathbb{R}^n} N(y, u, B_r(x_0)) dy \leq M |f(B_r(x_0))|,$$

and on the other hand $|\det Du(x)| \geq \mathcal{K}^{-1} \varepsilon^n$. Hence

$$\frac{1}{M\mathcal{K}} \varepsilon^n \leq \frac{|f(B_r(x_0))|}{|B_r(x_0)|}.$$

In particular we deduce that for all $r < r(x_0)$

$$\frac{1}{L(M\mathcal{K})^{1/n}} \varepsilon r \leq \min_{|x-x_0|=r} |u(x) - u(x_0)|.$$

For $t \in [0, 1]$ define $u^t = u + t\phi$. Then

$$u^t(x) - u^t(x_0) = u(x) - u(x_0) + t(\phi(x) - \phi(x_0)),$$

so that

$$\min_{|x-x_0|=r} |u^t(x) - u^t(x_0)| \geq \left(\frac{1}{L(M\mathcal{K})^{1/n}} \varepsilon - t\delta \right) r.$$

Choosing $\delta = \frac{1}{2} \frac{1}{L(M\mathcal{K})^{1/n}} \varepsilon$ we find

$$\min_{|x-x_0|=r} |u^t(x) - u^t(x_0)| \geq \delta r \quad \text{for all } r < r(x_0), t \in [0, 1],$$

and in particular $u^t(x) \neq u^t(x_0)$ for all $x \in \partial B_r(x_0)$. Hence

$$(2) \quad \mu(u^0(x_0), u^0, B_r(x_0)) = \mu(u^1(x_0), u^1, B_r(x_0)) \text{ for all } 0 < r < r(x_0).$$

For $x_0 \in \Omega$ the local topological index of the mapping u at x_0 is defined to be

$$i(x_0, u) = \mu(u(x_0), u, B_r(x_0)),$$

where $r > 0$ is chosen sufficiently small so that $\overline{B_r(x_0)} \cap u^{-1}\{u(x_0)\} = \{x_0\}$. Therefore (2) implies that

$$i(x_0, u) = i(x_0, u + \phi).$$

Since the branch set is defined as $B(u) = \{x \in \Omega : i(x, u) > 1\}$, we deduce that $B(u) \cap G = B(u + \phi) \cap G$. Q.E.D.

PROPOSITION 2. *Let $\Gamma \subset \mathbb{R}^{n \times n}$ be a compact set of $n \times n$ matrices and $\Omega \subset \mathbb{R}^n$ an open domain. Let $u \in W^{1,n}(\Omega, \mathbb{R}^n)$, and suppose that there exists $\mathcal{K} \geq 1$ and $\varepsilon > 0$ such that for all $A \in \Gamma$*

$$\varepsilon \leq \|Du(x) - A\|^n \leq \mathcal{K} \det(Du(x) - A) \quad \text{a.e. } x \in \Omega.$$

Then there exists an open and connected subset $\Omega_0 \subset \Omega$ with $|\Omega \setminus \Omega_0| = 0$ such that for all $x_0 \in \Omega_0$ there exists a radius $\tilde{r}(x_0) > 0$ such that

$$u(x) - u(y) \neq A(x - y) \quad \text{for all } x, y \in B_{\tilde{r}(x_0)}(x_0) \text{ and all } A \in \Gamma.$$

PROOF. For simplicity of notation let us treat $\Gamma \subset \mathbb{R}^{n \times n}$ as the image of a continuous map $\Gamma : \mathcal{S} \rightarrow \mathbb{R}^{n \times n}$, where \mathcal{S} is a compact metric space which we think of as an index set. Consider for any $t \in \mathcal{S}$ the mapping

$$u^t(x) := u(x) - \Gamma(t)x.$$

By assumption $u^t \in W^{1,n}(\Omega, \mathbb{R}^n)$ and

$$\varepsilon \leq \|Du^t(x)\|^n \leq \mathcal{K} \det Du^t(x) \quad \text{a.e. } x \in \Omega,$$

in particular $\{u^t\}_{t \in \mathcal{S}}$ is an equicontinuous family of quasiregular mappings.

Let $G \subset \Omega$ be a subdomain with $\overline{G} \subset \Omega$ and $|\partial G| = 0$. From [18] we know that $N(u^t, G) < \infty$ for each $t \in \mathcal{S}$. We aim to show that in fact $\sup_{t \in \mathcal{S}} N(u^t, G) < \infty$. To this end note that since each u^t is a discrete mapping, for each $x \in G$ and each $t \in \mathcal{S}$ there exists $r = r(x, t) > 0$ so that

$$\overline{B_r(x)} \cap (u^t)^{-1}\{u^t(x)\} = \{x\}.$$

More precisely from Proposition 1 we deduce that there exists $r = r(x, t) > 0$ and $\delta = \delta(t) > 0$ so that

$$\overline{B_r(x)} \cap (u^s)^{-1}\{u^s(x)\} = \{x\} \text{ for all } |s - t| < \delta,$$

and

$$\text{dist}(u^s(x), u^s(\partial B_r(x))) \geq \delta r \text{ for all } |s - t| < \delta$$

holds for all $t \in \mathcal{S}$. Hence by compactness of \mathcal{S} there exists $r = r(x) > 0$ and $\delta > 0$ (now independent of t) so that

$$(3) \quad \overline{B_r(x)} \cap (u^t)^{-1}\{u^t(x)\} = \{x\} \text{ for all } t \in \mathcal{S},$$

and

$$(4) \quad \text{dist} (u^t(x), u^t(\partial B_r(x))) \geq \delta r \text{ for all } t \in \mathcal{S}.$$

Indeed, the sets $V(t) := \{s \in \mathcal{S} : |t - s| < \delta(t)\}$ form an open cover for \mathcal{S} , so it suffices to take a finite subcover $V(t_1), \dots, V(t_N)$ and then define

$$r(x) = \min_{i=1, \dots, N} r(x, t_i) \text{ and } \delta = \min_{i=1, \dots, N} \delta(t_i)$$

in (3) and (4). Let

$$s(x, t) := \text{dist} (u^t(x), u^t(\partial B_r(x))),$$

and let $U(x, t)$ be the connected component of

$$(u^t)^{-1} (B_{s(x,t)}(u^t(x)))$$

containing x . Then $U(x, t) \subset B_{r(x)}(x)$ is a normal neighbourhood of x for the mapping u^t . Since the family $\{u^t\}$ is equicontinuous, from (4) we deduce that there exists $\tilde{r} = \tilde{r}(x) > 0$ so that $B_{\tilde{r}}(x) \subset U(x, t)$. Since $\tilde{r}(x)$ is independent of t , there exists a number $J \in \mathbb{N}$ so that for each fixed $t \in \mathcal{S}$ the compact set \overline{G} can be covered by at most J normal neighbourhoods $U(x_1, t), \dots, U(x_J, t)$. Then

$$N(u^t, G) \leq \sum_{j=1}^J N(u^t, U(x_j, t)) = \sum_{j=1}^J i(x_j, u^t).$$

On the other hand Proposition 1 implies that for each fixed $x_0 \in G$ the function $t \mapsto i(x_0, u^t)$ is continuous, hence bounded on \mathcal{S} . Therefore we deduce that $N(u^t, G)$ is bounded independently of t .

Proposition 1 now implies that there exists $\delta > 0$ (not depending on t) so that

$$B(u^t) \cap G = B(u^s) \cap G \text{ for all } s, t \in \mathcal{S} \text{ with } |s - t| < \delta.$$

In particular the set

$$B = \bigcup_{t \in \mathcal{S}} B(u^t) \cap G$$

is a finite union of closed sets of topological dimension $(n - 2)$ and Lebesgue measure zero [20], hence B is a closed set of dimension $(n - 2)$ and Lebesgue measure zero. This implies that the set $G_0 := G \setminus B$ is open and connected (see [8, Theorem IV.4]), and $|G \setminus G_0| = 0$.

In G_0 each mapping u^t is a local homeomorphism. More precisely, let $x_0 \in G_0$. Since $U(x_0, t)$ is a normal neighbourhood of x_0 for the mapping u^t , we have

$$N(u^t, U(x_0, t)) = i(x_0, u^t) = 1 \quad \text{for all } t \in \mathcal{S}.$$

Since $B_{\tilde{r}(x_0)}(x_0) \subset U(x_0, t)$ for all $t \in \mathcal{S}$, we deduce that each mapping u^t is injective on $B_{\tilde{r}(x_0)}(x_0)$. Therefore

$$u^t(x) \neq u^t(y) \text{ for all } t \in \mathcal{S}, x, y \in B_{\tilde{r}(x_0)}(x_0),$$

in other words

$$u(x) - u(y) - A(x - y) \neq 0 \text{ for all } x, y \in B_{\tilde{r}(x_0)}(x_0) \text{ and all } A \in \Gamma.$$

The proposition now follows by exhausting Ω with a nested sequence of subdomains $G \subset \Omega$ with $|\partial G| = 0$ and $\overline{G} \subset \Omega$. Q.E.D.

3. PROOF OF THE MAIN RESULT

Proof of Theorem 1. Let $K = [Du]$. Recall, that by definition $[Du]$ is the smallest closed subset of $\mathbb{R}^{2 \times 2}$ such that $Du(x) \in [Du]$ for almost every $x \in \Omega$.

We argue by contradiction, assuming that K^{rc} is not connected. According to [7, Theorem 5] there exists a continuous closed curve $\Gamma : \mathcal{S}^1 \rightarrow \mathbb{R}^{2 \times 2}$ such that

$$K \subset \mathcal{U}_\Gamma := \{A \in \mathbb{R}^{2 \times 2} : \det(A - \Gamma(t)) > 0 \text{ for all } t \in \mathcal{S}^1\},$$

where K is contained in more than one component of \mathcal{U}_Γ , and the curve Γ satisfies $\Gamma(t) \neq \Gamma(s)$ for $t \neq s$ and for some $\mathcal{K}' \geq 1$

$$(5) \quad \|\Gamma(t) - \Gamma(s)\|^2 \leq \mathcal{K}' \det(\Gamma(t) - \Gamma(s)) \quad \text{for all } t, s \in \mathcal{S}^1.$$

Since K and Γ are compact, there exist $\mathcal{K} > \mathcal{K}'$ and $\varepsilon > 0$ such that for all $t \in \mathcal{S}^1$

$$\varepsilon \leq \|Du(x) - \Gamma(t)\|^2 \leq \mathcal{K} \det(Du(x) - \Gamma(t)) \quad \text{a.e. } x \in \Omega.$$

But then Proposition 2 implies that there exists a connected and open subset $\Omega_0 \subset \Omega$ with $|\Omega \setminus \Omega_0| = 0$ and for each $x_0 \in \Omega_0$ there exists a radius $\tilde{r}(x_0) > 0$ such that

$$u(x) - u(y) \neq \Gamma(t)(x - y) \quad \text{for all } x, y \in B_{\tilde{r}(x_0)}(x_0) \text{ and } t \in \mathcal{S}^1.$$

Setting $y = x + \varepsilon e_1$ for $0 < \varepsilon < \tilde{r}(x)$ we obtain

$$\frac{u(x + \varepsilon e_1) - u(x)}{\varepsilon} \neq \Gamma(t)e_1 \quad \text{for all } t \in \mathcal{S}^1, (x, \varepsilon) \in \Delta,$$

where $\Delta = \{(x, \varepsilon) : x \in \Omega_0, 0 < \varepsilon < \tilde{r}(x)\}$. Since Γ satisfies (5),

$$\Gamma(\cdot)e_1 : \mathcal{S}^1 \rightarrow \mathbb{R}^2$$

is a continuous imbedding, hence by the Jordan separation theorem the image $\{\Gamma(t)e_1 : t \in \mathcal{S}^1\}$ separates \mathbb{R}^2 into two disjoint regions ω and $\mathbb{R}^2 \setminus \overline{\omega}$. Since Δ is a connected set, we deduce that

$$\frac{u(x + \varepsilon e_1) - u(x)}{\varepsilon} \in \omega \text{ for all } (x, \varepsilon) \in \Delta,$$

or

$$\frac{u(x + \varepsilon e_1) - u(x)}{\varepsilon} \in \mathbb{R}^2 \setminus \overline{\omega} \text{ for all } (x, \varepsilon) \in \Delta.$$

Since u is quasiregular, it is differentiable almost everywhere in Ω . Therefore, recalling that $|\Omega \setminus \Omega_0| = 0$, we obtain

$$(6) \quad \begin{aligned} & \partial_{x_1} u(x) \in \omega \text{ for a.e. } x \in \Omega, \\ & \text{or} \\ & \partial_{x_1} u(x) \in \mathbb{R}^2 \setminus \bar{\omega} \text{ for a.e. } x \in \Omega. \end{aligned}$$

In light of [7, Lemma 2] this implies that K has to be contained in a single component of \mathcal{U}_Γ , giving us the required contradiction.

Q.E.D.

Proof of Corollary 1. Suppose that $A \in [Du]$ is an isolated point, and assume for a contradiction that for all $B \in [Du]^{rc} \setminus \{A\}$ we have $\text{rank}(A - B) > 1$.

If $\det(A - B) > 0$ for all $B \in [Du] \setminus \{A\}$, then - since A is isolated and hence $[Du] \setminus \{A\}$ is compact - there exists a constant $\gamma \geq 1$ so that

$$\|Du(x) - A\|^2 \leq \gamma \det(Du(x) - A) \quad \text{a.e. } x \in \Omega.$$

This means that the map $x \mapsto u(x) - Ax$ is quasiregular. By the unique continuation property of quasiregular mappings we deduce that $Du(x) = Ax$ a.e., a contradiction. Similarly, we obtain the same contradiction if $\det(A - B) < 0$ for all $B \in [Du] \setminus \{A\}$ (by just considering a linear change of variables).

Therefore, we may assume that there exists at least two matrices

$$A_1, A_2 \in [Du] \setminus \{A\}$$

such that $\det(A - A_1) < 0$ and $\det(A - A_2) > 0$. If $[Du]^{rc} \setminus \{A\}$ is connected, we obtain by continuity the existence of $B \in [Du]^{rc} \setminus \{A\}$ with $\det(A - B) = 0$.

Otherwise let K_1, K_2 be disjoint connected components of $[Du]^{rc} \setminus \{A\}$ containing A_1 and A_2 , respectively. We claim first of all that

$$(7) \quad A \in \overline{K_1} \cap \overline{K_2}.$$

Indeed, assume the contrary, so that, without loss of generality, $A \notin \overline{K_1}$. Then there exists $\eta > 0$ with

$$(8) \quad B_{2\eta}(A) \cap K_1 = \emptyset.$$

As $[Du]^{rc}$ and hence $\tilde{K} := [Du]^{rc} \setminus B_\eta(A)$ is compact, and since K_1 is clearly the connected component of \tilde{K} containing A_1 , we see from [14, §42 II 2] or [4, §4.4 II 6] that K_1 is equal to the intersection of the family \mathcal{F} of all open and closed subsets of \tilde{K} which contain A_1 . In particular, since \mathcal{F} is closed under finite intersections, we conclude that there is some $V \in \mathcal{F}$ with

$$V \subset B_\eta(K_1) \text{ and } A_2 \notin V.$$

Here $B_\eta(K_1)$ denotes the open η -neighbourhood of K_1 . But then $V \cap B_\eta(A) = \emptyset$ because of (8), and hence V is closed and open in $[Du]^{rc}$. We

conclude that $[Du]^{rc}$ would be disconnected, in contradiction with Theorem 1. This proves the claim (7).

Now suppose without loss of generality that $\det(A_1 - A_2) > 0$, and consider the function $f(X) = \det(X - A_1)$ restricted to K_2 . Since $\det(A - A_1) < 0$ and $A \in \overline{K_2}$, there exists $A' \in K_2$ such that $f(A') < 0$, by continuity. On the other hand $f(A_2) > 0$, therefore there exists, again by continuity, $A_3 \in K_2$ with $f(A_3) = 0$. In particular $[A_1, A_3]$ is a rank-one segment, which therefore is contained in $[Du]^{rc}$. If $A \notin [A_1, A_3]$ then we obtain a contradiction with the assumption that $A_1 \in K_1$ and $A_3 \in K_2$ are contained in different connected components of $[Du]^{rc} \setminus \{A\}$. On the other hand, if $A \in [A_1, A_3]$, then in particular $\det(A - A_1) = 0$, contradicting the assumption that $\det(A - A_1) < 0$. This finishes the proof. Q.E.D.

4. INCOMPATIBLE SETS OF GRADIENTS

Following [1] two disjoint compact sets of matrices $K_1, K_2 \subset \mathbb{R}^{m \times n}$ are said to be incompatible if whenever Ω is a bounded domain and $\{u_j\}$ is a sequence of maps bounded in $W^{1,1}(\Omega)$ such that

$$\text{dist}(Du_j, K_1 \cup K_2) \rightarrow 0 \text{ in } L^1(\Omega) \text{ strongly,}$$

then - up to a subsequence -

$$\text{dist}(Du_j, K_1) \rightarrow 0 \text{ or } \text{dist}(Du_j, K_2) \rightarrow 0 \text{ strongly in } L^1(\Omega).$$

In the language of Young measures this is equivalent to saying that whenever $\{\nu_x\}_{x \in \Omega}$ is a gradient Young measure supported in $K_1 \cup K_2$, that is,

$$\text{supp } \nu_x \subset K_1 \cup K_2 \text{ a.e. } x \in \Omega,$$

then

$$\text{either } \text{supp } \nu_x \subset K_1 \text{ a.e. or } \text{supp } \nu_x \subset K_2 \text{ a.e.}$$

In short, the sets K_1 and K_2 are *incompatible for gradient Young measures*. From the point of view of material microstructure it is of interest to be able to characterize such incompatible sets. Indeed, in this situation the inclusion problem $Du(x) \in K_1 \cup K_2$ would correspond to energy-minimizing deformations of an elastic material, and roughly speaking incompatibility prevents large scale oscillations (oscillations *between* K_1 and K_2), whilst still allowing for local oscillations *within* each energy-well K_1 or K_2 .

Pairs of incompatible sets have several nice features. First of all, if K_1 and K_2 are incompatible for gradient Young measures, then sufficiently small ε -neighbourhoods $(K_1)_\varepsilon$ and $(K_2)_\varepsilon$ are still incompatible. This was established by Ball and James in the early 90s in their study of metastability [1]. Moreover, one gets precise control of the gradient for approximating sequences in the form of a rigidity estimate:

$$\min \left(\int_{\Omega} \text{dist}^p(Du, K_1), \int_{\Omega} \text{dist}^p(Du, K_2) \right) \leq C(p, \Omega) \int_{\Omega} \text{dist}^p(Du, K_1 \cup K_2),$$

valid for all $u \in W^{1,p}(\Omega, \mathbb{R}^m)$ and all $p \in [1, \infty)$. This was proved in [6] using the method of Ball and James [1].

The simplest example of incompatible sets, as already pointed out in the introduction, is given by the singleton sets $\{A\}, \{B\}$ whenever $A, B \in \mathbb{R}^{m \times n}$ with $\text{rank}(A - B) > 1$. In [27] K. Zhang showed that in this case there exists $\varepsilon > 0$, so that the sets

$$K_1 = \{X \in \mathbb{R}^{m \times n} : |X - A| \leq \varepsilon\} \text{ and } K_2 = \{X \in \mathbb{R}^{m \times n} : |X - B| \leq \varepsilon\}$$

are still incompatible (in fact Zhang's result applies to the neighbourhood of any finite collection of matrices contained in a subspace without rank-one connections). More precisely, Zhang obtains explicit estimates for $\varepsilon > 0$ in terms of Schauder $L^\infty - BMO$ estimates (see also [26] for a similar technique applied to incompatible wells in 2D). In contrast, in the aforementioned stability result of Ball and James $\varepsilon > 0$ is obtained in a contradiction argument. Other types of explicit examples of incompatible sets were obtained by V. Šverák [22] in connection with the Monge-Ampère equation and by J. P. Matos in [15] concerning the two-well problem in 3D.

Our Theorem 1, combined with results in [7] allows us to completely characterize incompatible sets in $\mathbb{R}^{2 \times 2}$ in terms of the underlying rank-one geometry.

COROLLARY 2. *Two disjoint compact sets $K_1, K_2 \subset \mathbb{R}^{2 \times 2}$ are incompatible for gradient Young measures if and only if $K_1^{rc} \cap K_2^{rc} = \emptyset$ and $K_1^{rc} \cup K_2^{rc} = (K_1 \cup K_2)^{rc}$.*

In order to explain the meaning of this result, we briefly recall a few more notions from the nonconvex calculus of variations. First of all, a gradient Young measure $\{\nu_x\}_{x \in \Omega}$ is said to be *homogeneous* if ν_x is independent of $x \in \Omega$. Homogeneous gradient Young measures appear in the study of compactness of sequences of gradients $\{Du_j\}$. A further subclass of homogeneous gradient Young measures is formed by *laminates*. Roughly speaking laminates are probability measures that can be constructed through rank-one connections. More precisely, laminates are the smallest class of probability measures on the space of matrices that are

- (i) closed under splitting,
- (ii) closed under weak* convergence,
- (iii) and contain all measures of the form $\lambda\delta_A + (1 - \lambda)\delta_B$ whenever $\text{rank}(A - B) \leq 1$ and $\lambda \in [0, 1]$.

Being closed under splitting means that if ν is a laminate of the form

$$\nu = \lambda\delta_A + (1 - \lambda)\tilde{\nu}$$

for some probability measure $\tilde{\nu}$, and μ is a laminate with barycenter $\bar{\mu} = A$, then the measure

$$\lambda\mu + (1 - \lambda)\tilde{\nu}$$

is also a laminate. For basic properties of these classes of measures we refer the reader to [16, 17].

We recall in particular that the rank-one convex hull K^{rc} of a compact set of matrices can be defined as the set of barycenters of laminates supported in K :

$$K^{rc} = \{\bar{\mu} : \mu \text{ is a laminate with } \text{supp } \mu \subset K\}.$$

To each class of measures one can associate a notion of *incompatibility* for pairs of compact sets. Thus for example K_1, K_2 are said to be *incompatible for laminates* if whenever μ is a laminate with support

$$\text{supp } \mu \subset K_1 \cup K_2,$$

then

$$\text{supp } \mu \subset K_1 \text{ or } \text{supp } \mu \subset K_2.$$

Similarly, K_1, K_2 are said to be *homogeneously incompatible* if they are incompatible for homogeneous gradient Young measures. Equivalently, K_1, K_2 are homogeneously incompatible if whenever $\{u_j\}$ is a sequence of maps bounded in $W_0^{1,1}(\Omega)$ such that

$$\text{dist}(A + Du_j, K_1 \cup K_2) \rightarrow 0 \text{ in } L^1(\Omega) \text{ strongly}$$

for some matrix A , then - up to a subsequence -

$$\text{dist}(A + Du_j, K_1) \rightarrow 0 \text{ or } \text{dist}(A + Du_j, K_2) \rightarrow 0 \text{ strongly in } L^1(\Omega).$$

The meaning of Corollary 2 is that in the space of 2×2 matrices the three notions of incompatibility are equivalent:

COROLLARY 3. *Let $K_1, K_2 \subset \mathbb{R}^{2 \times 2}$ be disjoint compact sets. The following are equivalent:*

- (i) K_1, K_2 are incompatible for gradient Young measures,
- (ii) K_1, K_2 are incompatible for homogeneous gradient Young measures,
- (iii) K_1, K_2 are incompatible for laminates.

The equivalence between (ii) and (iii) was already proved in [7, Corollary 1]. Here we establish the equivalence of (i) and (ii). Proving this equivalence amounts to a passage from approximating sequences of the form $\{A + Du_j\}$ with $Du_j \in W_0^{1,1}(\Omega)$ to general sequences $\{Du_j\} \subset W^{1,1}(\Omega)$. Indeed, a crucial aspect of Theorem 1 is that there is no assumption made on the boundary values of the map $u : \Omega \rightarrow \mathbb{R}^2$.

Proof of Corollary 2. One direction is easy: if K_1, K_2 are incompatible for gradient Young measures, then in particular they are incompatible for laminates. Thus any laminate μ with support $\text{supp } \mu \subset K_1 \cup K_2$ has to be supported in K_1 or K_2 . Therefore the definition of rank-one convex hull implies that $(K_1 \cup K_2)^{rc} = K_1^{rc} \cup K_2^{rc}$. It remains to show that $K_1^{rc} \cap K_2^{rc} = \emptyset$. Assume for a contradiction that $K_1^{rc} \cap K_2^{rc} \neq \emptyset$, so that there exist laminates μ_1, μ_2 with support $\text{supp } \mu_i \subset K_i$ with common barycenter $\bar{\mu}_1 = \bar{\mu}_2 \in K_1^{rc} \cap K_2^{rc}$. But then the laminate defined as $\mu := \frac{1}{2}\mu_1 + \frac{1}{2}\mu_2$ has support $\text{supp } \mu \subset K_1 \cup K_2$, but doesn't satisfy $\text{supp } \mu \subset K_1$ or $\text{supp } \mu \subset K_2$. This gives a contradiction, and therefore necessarily $K_1^{rc} \cap K_2^{rc} = \emptyset$.

For the other direction suppose now that $K_1^{rc} \cap K_2^{rc} = \emptyset$ and $(K_1 \cup K_2)^{rc} = K_1^{rc} \cup K_2^{rc}$. We claim that in this case K_1 and K_2 are incompatible for laminates. Indeed, suppose μ is a laminate with support $\text{supp } \mu \subset K_1 \cup K_2$. Then

$$\text{supp } \mu \subset (K_1 \cup K_2)^{rc} = K_1^{rc} \cup K_2^{rc},$$

and on the other hand it is well known that $(\text{supp } \mu)^{rc}$ is a connected set (see [12, Theorem 4.9]). Therefore necessarily

$$(\text{supp } \mu)^{rc} \subset K_1^{rc} \text{ or } (\text{supp } \mu)^{rc} \subset K_2^{rc}.$$

To conclude that $\text{supp } \mu \subset K_1$ or $\text{supp } \mu \subset K_2$ just note that $\text{supp } \mu \subset (\text{supp } \mu)^{rc}$ and that $K_i \cap K_j^{rc} = \emptyset$ for $i \neq j$.

Having just shown that K_1 and K_2 are incompatible for laminates, we can now invoke [7, Corollary 1], which implies that K_1 and K_2 are incompatible for homogeneous gradient Young measures. In particular $K_1^{qc} \cap K_2^{qc} = \emptyset$ and $(K_1 \cup K_2)^{qc} = K_1^{qc} \cup K_2^{qc}$ (see [7, Corollary 3]).

Now suppose that $\{\nu_x\}_{x \in \Omega}$ is a gradient Young measure such that

$$\text{supp } \nu_x \subset K_1 \cup K_2 \text{ for a.e. } x \in \Omega.$$

Since ν_x coincides with a homogeneous gradient Young measure for a.e. x and K_1, K_2 are incompatible for homogeneous gradient Young measures, we deduce that for almost every $x \in \Omega$ there exists $i = i_x \in \{1, 2\}$ such that

$$\text{supp } \nu_x \subset K_{i_x}.$$

It remains to show that $i_x = 1$ a.e. or $i_x = 2$ a.e. To this end recall (see [10]) that because $\{\nu_x\}_{x \in \Omega}$ is a gradient Young measure, there exists a Lipschitz mapping $u : \Omega \rightarrow \mathbb{R}^2$ such that $Du(x) = \bar{\nu}_x$ a.e. $x \in \Omega$. In particular

$$[Du] \subset (K_1 \cup K_2)^{qc}.$$

By Theorem 1 we know that $[Du]^{rc}$ is connected, and on the other hand

$$[Du]^{rc} \subset [Du]^{qc} \subset (K_1 \cup K_2)^{qc} = K_1^{qc} \cup K_2^{qc}.$$

Since $K_1^{qc} \cap K_2^{qc} = \emptyset$, we deduce that

$$[Du] \subset K_1^{qc} \text{ or } [Du] \subset K_2^{qc}.$$

Finally, note that $\bar{\nu}_x \in K_i^{qc}$ if and only if $\text{supp } \nu_x \subset K_i$ (for $i = 1, 2$) since $K_1^{qc} \cap K_2^{qc} = \emptyset$. Hence we conclude that $\text{supp } \nu_x \subset K_1$ a.e. $x \in \Omega$ or $\text{supp } \nu_x \subset K_2$ a.e. $x \in \Omega$.

Q.E.D.

Proof of Corollary 3. Since the implications (i) \Rightarrow (ii) \Rightarrow (iii) follow from the definitions, it suffices to prove that (iii) \Rightarrow (i). Suppose that K_1, K_2 are incompatible for laminates. Then, precisely as in the proof of Corollary 2 above, we have that

$$(K_1 \cup K_2)^{rc} = K_1^{rc} \cup K_2^{rc} \text{ and } K_1^{rc} \cap K_2^{rc} = \emptyset.$$

But then Corollary 2 implies that K_1, K_2 are incompatible for gradient Young measures.

Q.E.D.

Proof of Theorem 2. The statement of the theorem is a direct consequence of Corollary 2 together with [6, Theorem 1.2]. Q.E.D.

REFERENCES

- [1] BALL, J. M., AND JAMES, R. D. Incompatible sets of gradients and metastability. *in preparation*.
- [2] BALL, J. M., AND JAMES, R. D. Fine phase mixtures as minimizers of energy. *Arch. Rational Mech. Anal.* 100 (1987), 13–52.
- [3] BHATTACHARYA, K., FIROOZY, N. B., JAMES, R. D., AND KOHN, R. V. Restrictions on microstructure. *Proc. Roy. Soc. Edinburgh Sect. A* 124, 5 (1994), 843–878.
- [4] BOURBAKI, N. *General topology. Chapters 1–4*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1998. Translated from the French, Reprint of the 1989 English translation.
- [5] CHIPOT, M., AND KINDERLEHRER, D. Equilibrium configurations of crystals. *Arch. Rational Mech. Anal.* 103 (1988), 237–277.
- [6] DE LELLIS, C., AND SZÉKELYHIDI, JR., L. Simple proof of two-well rigidity. *C. R. Math. Acad. Sci. Paris* 343, 5 (2006), 367–370.
- [7] FARACO, D., AND SZÉKELYHIDI, JR., L. Tartar’s conjecture and localization of the quasiconvex hull in $\mathbb{R}^{2 \times 2}$. Preprint, MPI-MIS, 2006.
- [8] HUREWICZ, W., AND WALLMAN, H. *Dimension Theory*. Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941.
- [9] IWANIEC, T., VERCHOTA, G. C., AND VOGEL, A. L. The failure of rank-one connections. *Arch. Ration. Mech. Anal.* 163, 2 (2002), 125–169.
- [10] KINDERLEHRER, D., AND PEDREGAL, P. Characterizations of Young measures generated by gradients. *Arch. Rational Mech. Anal.* 115 (1991), 329–365.
- [11] KIRCHHEIM, B. Deformations with finitely many gradients and stability of quasiconvex hulls. *C. R. Acad. Sci. Paris Sér. I Math.* 332, 3 (2001), 289–294.
- [12] KIRCHHEIM, B. Rigidity and Geometry of microstructures. Habilitation thesis, University of Leipzig, 2003.
- [13] KIRCHHEIM, B., MÜLLER, S., AND ŠVERÁK, V. Studying nonlinear PDE by geometry in matrix space. In *Geometric analysis and Nonlinear partial differential equations*, S. Hildebrandt and H. Karcher, Eds. Springer-Verlag, 2003, pp. 347–395.
- [14] KURATOWSKI, C. *Topologie. Vol. II*. Troisième édition, corrigée et complétée de deux appendices. Monografie Matematyczne, Tom 21. Państwowe Wydawnictwo Naukowe, Warsaw, 1961.
- [15] MATOS, J. P. Young measures and the absence of fine microstructures in a class of phase transitions. *European J. Appl. Math.* 3, 1 (1992), 31–54.
- [16] MÜLLER, S. Variational models for microstructure and phase transitions. In *Calculus of variations and geometric evolution problems (Cetraro, 1996)*, vol. 1713 of *Lecture Notes in Math*. Springer, Berlin, 1999, pp. 85–210.
- [17] PEDREGAL, P. Laminates and microstructure. *European J. Appl. Math.* 4 (1993), 121–149.
- [18] RESHETNYAK, Y. G. A condition for boundedness of the index for mappings with bounded distortion. *Sibirsk. Mat. Ž.* 9 (1968), 368–374.
- [19] RESHETNYAK, Y. G. The local structure of mappings with bounded distortion. *Sibirsk. Mat. Ž.* 10 (1969), 1311–1333.
- [20] RESHETNYAK, Y. G. The set of branch points of mappings with bounded distortion. *Sibirsk. Mat. Ž.* 11 (1970), 1333–1339.
- [21] RICKMAN, S. *Quasiregular mappings*, vol. 26 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1993.

- [22] ŠVERÁK, V. On regularity for the Monge-Ampère equations. Preprint, Heriot-Watt University, 1991.
- [23] ŠVERÁK, V. On Tartar's conjecture. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 10 (1993), 405–412.
- [24] SZÉKELYHIDI, JR., L. Rank-one convex hulls in $\mathbb{R}^{2 \times 2}$. *Calc. Var. Partial Differential Equations* 22, 3 (2005), 253–281.
- [25] ZHANG, K. On connected subsets of $M^{2 \times 2}$ without rank-one connections. *Proc. Roy. Soc. Edinburgh Sect. A* 127, 1 (1997), 207–216.
- [26] ZHANG, K. Neighborhoods of parallel wells in two dimensions that separate gradient Young measures. *SIAM J. Math. Anal.* 34, 5 (2003), 1207–1225 (electronic).
- [27] ZHANG, K. On separation of gradient Young measures. *Calc. Var. Partial Differential Equations* 17, 1 (2003), 85–103.

MATHEMATICAL INSTITUTE, OXFORD UNIVERSITY, 24-29 ST. GILES', OXFORD, OX1 3LB, UK

E-mail address: kirchhei@maths.ox.ac.uk

DEPARTEMENT MATHEMATIK, ETH ZÜRICH, CH-8092 ZÜRICH, SWITZERLAND

E-mail address: szekelyh@math.ethz.ch