

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

NON-LOCAL ANISOTROPIC DISPERSAL
WITH MONOSTABLE NONLINEARITY

by

Jerome Coville, Juan Davila, and Salome Martinez

Preprint no.: 58

2007



NON-LOCAL ANISOTROPIC DISPERSAL WITH MONOSTABLE NONLINEARITY

JÉRÔME COVILLE¹, JUAN DÁVILA², AND SALOMÉ MARTÍNEZ²

ABSTRACT. We study the travelling wave problem

$$J \star u - u - cu' + f(u) = 0 \quad \text{in } \mathbb{R}, \quad u(-\infty) = 0, \quad u(+\infty) = 1$$

with an asymmetric kernel J and a monostable nonlinearity. We prove the existence of a minimal speed, and under certain hypothesis the uniqueness of the profile for $c \neq 0$. For $c = 0$ we show examples of non-uniqueness.

1. Introduction and main results

During the past ten years, much attention has been drawn to the study of the following nonlocal equation

$$(1.1) \quad \frac{\partial U}{\partial t} = \mathcal{J} \star U - U + f(U) \quad \text{in } \mathbb{R}^n \times \mathbb{R}^+,$$

$$(1.2) \quad U(x, 0) = U_0(x)$$

where \mathcal{J} is a probability density on \mathbb{R}^n and f a given nonlinearity. Such kind of equations appears in various applications ranging from population dynamics to Ising models as seen in [2, 7, 13, 14, 16, 17, 20, 24, 25] among many references. Here we will only be concerned with probability densities \mathcal{J} which satisfy the following assumption:

$$\mathcal{J} \in C(\mathbb{R}^n), \quad \mathcal{J}(z) \geq 0, \quad \int_{\mathbb{R}^n} \mathcal{J}(z) dz = 1, \quad \int_{\mathbb{R}^n} |z| \mathcal{J}(z) dz < \infty,$$

and nonlinearities f of monostable type, e.g.

$$(f1) \quad f \in C^1(\mathbb{R}), \quad \text{which satisfies } f(0) = f(1) = 0, \quad f'(1) < 0, \quad f|_{(0,1)} > 0$$

and $f|_{\mathbb{R} \setminus [0,1]} \leq 0$.

Such nonlinearities are commonly used in population dynamics to describe the interaction (birth, death, ...) of a species in its environment as described in [15, 18]. Equation (1.1) can be seen as nonlocal analogue of the traditional reaction diffusion equation

$$(1.3) \quad \frac{\partial U}{\partial t} = \Delta U + f(U) \quad \text{in } \mathbb{R}^n \times \mathbb{R}^+,$$

$$(1.4) \quad U(x, 0) = U_0(x).$$

Our analysis in this paper will mainly focus on the travelling wave solutions of equation (1.1) which have proved to be a robust way to describe the longtime behavior of solutions $U(x, t)$ of (1.3),(1.4) with compactly initial data U_0 , as seen in [1, 15, 16, 18, 28, 29]. These particular type of solutions are of the form $U_e(x, t) :=$

$u(x.e + ct)$ where $e \in \mathbb{S}^{n-1}$ is a given unit vector, the velocity $c \in \mathbb{R}$ and the scalar function u satisfy

$$(1.5) \quad J \star u - u - cu' + f(u) = 0 \quad \text{in } \mathbb{R},$$

$$(1.6) \quad u(-\infty) = 0,$$

$$(1.7) \quad u(+\infty) = 1,$$

where $u(\pm\infty)$ denotes the limit of $u(x)$ as $x \rightarrow \pm\infty$ and J is the real function defined as

$$J(s) := \int_{\mathbb{R}^{n-1}} \mathcal{J}(s, y) dy$$

after a suitable rotation of the coordinates. Thus we shall assume that the kernel J satisfies

$$(j1) \quad J \in C(\mathbb{R}), J(z) \geq 0, \int_{\mathbb{R}} J(z) dz = 1, \int_{\mathbb{R}} |z| J(z) dz < \infty.$$

We will call a solution $u \in L^\infty(\mathbb{R})$ to (1.5)–(1.7) a travelling wave or travelling front if it is non-decreasing.

The first works to study travelling fronts in this setting are due to Schumacher [25] and Weinberger [28, 29] who constructed travelling fronts satisfying some exponential decay for J symmetric and particular monostable nonlinearities, the so called KPP nonlinearity, e.g

$$(f2) \quad f \text{ is monostable and satisfies } f(s) \leq f'(0)s.$$

Then, Harris, Hudson and Zinner [19] and more recently Carr and Chmaj [5], Chen and Guo [6] and Coville and Dupaigne [12] extended and completed the work of Schumacher to more general monostable nonlinearities and dispersal kernels J satisfying what is called in the literature the Mollison condition [21, 23, 24]:

$$(j2) \quad \exists \lambda > 0 \quad \text{such that} \quad \int_{-\infty}^{\infty} J(-z) e^{\lambda z} dz < +\infty.$$

More precisely, they show that

Theorem 1.1. [5, 6, 12, 19, 25] *Let f be a monostable nonlinearity, J be a symmetric function satisfying (j1)–(j2). Then there exists a constant $c^* > 0$ such that for all $c \geq c^*$, there exists an increasing function u , such that (u, c) is a solution of (1.5)–(1.7) and for any $c < c^*$, there exists no increasing solution of (1.5)–(1.7). Moreover, if in addition $f'(0) > 0$, then any bounded solution (u, c) of (1.5)–(1.7) is unique up to translation.*

Furthermore, as in the classical case, when the nonlinearity is KPP the critical speed c^* can be precisely evaluated by means of a formula.

Theorem 1.2. [5, 6, 19, 25, 29] *Let f be a KPP nonlinearity and J be a symmetric function satisfying (j1)–(j2). Then the critical speed c^* is given by*

$$c^* = \min_{\lambda > 0} \frac{1}{\lambda} \left(\int_{\mathbb{R}} J(x) e^{\lambda x} dx + f'(0) - 1 \right).$$

In Theorems 1.1 and 1.2 the dispersal kernel J is assumed to be symmetric. This corresponds to the situation where the dispersion of the species is isotropic. Since the dispersal of an individual can be influenced in many ways (wind, landscape, ...), it is natural to ask what happens when the kernel J is non-symmetric. In this direction, we have the following result:

Theorem 1.3. *Let f be a monostable nonlinearity satisfying (f1) and J be a dispersal kernel satisfying (j1). Assume further that there exists (w, κ) with $w \in C(\mathbb{R})$ a supersolution of (1.5)-(1.7) in the sense:*

$$(1.8) \quad \begin{aligned} J \star w - w - \kappa w' + f(w) &\leq 0 \quad \text{in } \mathbb{R}, \\ w(-\infty) &\geq 0, \\ w(+\infty) &\geq 1 \end{aligned}$$

and such that $w(x_0) < 1$ for some $x_0 \in \mathbb{R}$. Then there exists a critical speed $c^ \leq \kappa$, such that for all $c \geq c^*$ there exists a non decreasing solution (u, c) to (1.5)-(1.7) and for $c < c^*$ there exists no non-decreasing travelling wave with speed c .*

We emphasize that in the above theorem we do not require monotonicity of the supersolution w . The first consequence of Theorem 1.3 is to relate the existence of a minimal speed c^* and the existence of a travelling front for any speed $c \geq c^*$ to the existence of a supersolution. In other words, we have the following necessary and sufficient condition:

Corollary 1.4. *Let f and J be such that (f1) and (j1) holds. Then there exists a non decreasing solution with minimal speed (u, c^*) of (1.5)-(1.7) if and only if there exists a supersolution (w, κ) of (1.5)-(1.7).*

The existence of a supersolution in Theorem 1.3 is automatic under extra assumptions on J . For instance, we have

Theorem 1.5. *Let f be a monostable nonlinearity and J be continuous compactly supported which satisfies (j1). Assume further that there exists $a \in \mathbb{R}$ such that $\{a, -a\} \subset \text{supp}(J)$. Then there exists a critical speed c^* , such that for all $c \geq c^*$ there exists a non decreasing function u such that (u, c) is a solution of (1.5)-(1.7). Moreover, there is no non decreasing travelling wave with speed $c < c^*$.*

Next we examine the validity of Theorem 1.2 for nonsymmetric J . Let c^1 denote the following quantity

$$c^1 := \inf_{\lambda > 0} \frac{1}{\lambda} \left(\int_{\mathbb{R}} J(-x) e^{\lambda x} dx + f'(0) - 1 \right).$$

For $c \geq c^1$ we denote $\lambda(c)$ the unique minimal $\lambda > 0$ such that

$$-c\lambda + \int_{\mathbb{R}} J(-x) e^{\lambda x} dx + f'(0) - 1 = 0.$$

We generalize a result of Carr and Chmaj [5] to the case when J is nonsymmetric.

Theorem 1.6. *Let f be a monostable nonlinearity satisfying (f1), $f'(0) > 0$, $f \in C^{1,\gamma}$ near 0 and there is $m \geq 1$, $\delta > 0$, $A > 0$ such that*

$$(1.9) \quad |u - f(u)| \geq Au^m \quad \text{for all } 0 \leq u < \delta.$$

Let J be a dispersal kernel satisfying (j1), $J \in C^1$ and is compactly supported. Then $c^1 \leq c^$. Moreover, if u is a solution of (1.5), (1.6), $0 \leq u \leq 1$, $u \not\equiv 0$ then, when $c = c^1$*

$$(1.10) \quad 0 < \lim_{x \rightarrow -\infty} \frac{u(x)}{|x| e^{\lambda(c^*)x}} < \infty,$$

and when $c > c^1$

$$(1.11) \quad 0 < \lim_{x \rightarrow -\infty} \frac{u(x)}{e^{\lambda(c)x}} < \infty.$$

Observe that when J is symmetric, by Jensen's inequality $c^1 > 0$. On the other hand, it is not difficult to construct examples of nonsymmetric J such that $c^1 \leq 0$. Note also that in Theorem 1.6 we do not need to assume that the solution u to (1.5), (1.6) is monotone.

Corollary 1.7. *If f and J satisfy the hypotheses of Theorem 1.6 and f satisfies also (f2) then*

$$c^* = c^1.$$

Regarding the uniqueness of the profile of the travelling waves we prove:

Theorem 1.8. *Assume f and J satisfy the hypotheses of Theorem 1.6 and J satisfies:*

$$(MP) \quad \exists a < 0 < b \quad \text{such that} \quad J(a) > 0, \quad J(b) > 0.$$

Then for $c \neq 0$ the solution of the problem (1.5)-(1.7) is unique up to translation.

We notice if $c \neq 0$ then any solution to (1.5) is continuous. In the case $c = 0$, the same argument used to prove Theorem 1.8 gives uniqueness for continuous solutions of (1.5)-(1.7) provided that this problem admits a continuous solution (see Remark 6.4). In the case $c = 0$ one sufficient condition for a solution $0 \leq u \leq 1$ to (1.5) to be continuous is that

$$u - f(u) \text{ is strictly increasing in } [0, 1].$$

In Proposition 6.8 we give examples of f and non-symmetric J such that no solution of (1.5)-(1.7) is continuous, and this problem admits infinitely many solutions.

Our results also have implications in the study of solutions to

$$(1.12) \quad J \star u - u + f(u) = 0$$

which corresponds to (1.5) with velocity $c = 0$. In [11] it was shown that if $f(u)/u$ is decreasing and J is symmetric then any non-trivial bounded solution of (1.12) is identically 1. The symmetry of J was important in the argument and it was conjectured that if the kernel J is not even (1.12) may have more than one solution. For this discussion we shall assume that f and J satisfy the hypotheses of Theorem 1.6 and f also satisfies (f2). We observe that when the dispersal kernel is not even, the critical velocity c^* can be non-positive. If $c^* \leq 0$ we obtain that the equation (1.12) has a non-constant positive solution satisfying (1.6)-(1.7). Similarly, equation (1.12) has positive solutions satisfying

$$\lim_{x \rightarrow -\infty} u(x) = 1, \quad \lim_{x \rightarrow +\infty} u(x) = 0, \quad u \text{ is non-increasing}$$

if and only if $c_* < 0$ where

$$c_* = \min_{\lambda > 0} \frac{1}{\lambda} \left(\int_{\mathbb{R}} J(x) e^{\lambda x} dx + f'(0) - 1 \right).$$

Observe that by Jensen's inequality we have $c^* > 0$ or $c_* > 0$. In summary, besides $u \equiv 0$ and $u \equiv 1$ equation (1.12) has travelling wave solutions if $c^* \leq 0$ or $c_* \leq 0$. Under some additional conditions on f these are the only possible solutions, that is, we have a complete classification result for (1.12), in the sense that we do not require the boundary conditions at $\pm\infty$, continuity nor the monotonicity of the solutions. The next Theorem can be shown by slightly modifying the arguments for Theorem 2.1 in [5].

Theorem 1.9. *Suppose f and J satisfy the hypotheses of Theorem 1.6, J satisfies (MP) and $f'(r) \leq f'(0)$ for $r \in (0, 1)$. Then any solution $0 \leq u \leq 1$ of problem (1.12) is one of the following: 1) $u \equiv 0$ or $u \equiv 1$, 2) a non-decreasing travelling wave or 3) a non-increasing travelling wave. Moreover in cases 2) and 3) the profile is unique up to translation.*

Regarding Mollison's condition (j2) let us mention that recently Kot and Medlock in [21] have shown that for a one dimensional problem when the dispersal kernel J is even with a fat tails and $f(s) := s(1 - s)$, the solutions of the initial value problem (1.1) do not behave like travelling waves with constant speed but rather like what they called *accelerating* waves. Moreover, they predict the apparition of accelerating waves for (1.1). More precisely, supported by numerical evidence and analytical proof, they conjecture that (1.1) admits travelling wave solutions if and only if for some $\lambda > 0$

$$\int_{-\infty}^{+\infty} J(z)e^{\lambda z} dz < +\infty.$$

It appears from our analysis on non symmetric dispersal kernels, that the existence of travelling waves with constant speed is more related to

$$\int_0^{+\infty} J(z)e^{\lambda z} dz < +\infty \quad \text{for some } \lambda > 0$$

if we look at fronts propagating from the left to the right and

$$\int_0^{+\infty} J(-z)e^{\lambda z} dz < +\infty \quad \text{for some } \lambda > 0$$

if we look at fronts propagating from the right to the left. As a consequence, for asymmetric kernels, it may happen that in one direction, the solution behave like a front with finite speed and in the other like an accelerating wave.

The outline of this paper is the following. In Section 2, we recall some results on front solutions for ignition nonlinearities, then in Section 3 we construct increasing solution of (3.6) and prove Theorem 1.5. Section 4 is devoted to the proof of Theorem 1.3. Section 5 contains the proof of Theorem 1.6 and Corollary 1.7. In Section 6 we prove the uniqueness of the profile Theorem 1.8 and Theorem 1.9.

2. Approximation by ignition type nonlinearities

The proof of Theorem 1.3 essentially relies on some estimates and properties of the speed of fronts for problem (1.1) with ignition type nonlinearities f . We say that f is of ignition type if $f \in C^1([0, 1])$ and

(f3) there exists $\rho \in (0, 1)$ such that $f|_{[0, \rho]} \equiv 0, f|_{(\rho, 1)} > 0$ and $f(1) = 0$.

Consider the following problem

$$(2.1) \quad \begin{cases} J \star u - u - cu' + f(u) = 0 & \text{in } \mathbb{R} \\ u(-\infty) = 0 \\ u(+\infty) = 1, \end{cases}$$

where $c \in \mathbb{R}$ and f is either an ignition nonlinearity or a monostable nonlinearity.

The main result in this section is the following:

Proposition 2.1. *Let f be a monostable nonlinearity and assume that J is a non negative continuous function of unit mass. Assume further that there exists (w, κ) a super-solution of (1.5)-(1.7). Let $(f_k)_{k \in \mathbb{N}}$ be any sequence of ignition functions which converges pointwise to f and satisfies $\forall k \in \mathbb{N}, f_k \leq f_{k+1} \leq f$ and let c_k be the unique speed of fronts associated to (2.1). Then*

$$(2.2) \quad \lim_{k \rightarrow +\infty} c_k = c^*,$$

exists and is independent of the sequence f_k . Furthermore, $c^ \leq \kappa$, there exists a non decreasing solution (u, c^*) of (1.5)-(1.7) and for $c < c^*$ there are no non-decreasing solutions to (1.5)-(1.7).*

The fact that for (2.1) with ignition type nonlinearity there exists a unique speed of fronts has been recently established by one of the authors in [8, 9, 10] and holds also for the following perturbation of (2.1)

$$(2.3) \quad \begin{cases} \epsilon u'' + J \star u - u - cu' + f(u) & = 0 & \text{in } \mathbb{R} \\ u(-\infty) & = 0 \\ u(+\infty) & = 1, \end{cases}$$

where $\epsilon \geq 0, c \in \mathbb{R}$.

Theorem 2.2. ([10, Theorem 1.2] and [8, Theorem 3.2]) *Let f be an ignition nonlinearity and assume that J satisfies (j1). Then there exists a non decreasing solution (u, c) of (2.3). Furthermore the speed c is unique. Moreover, if (v, c') is a super solution of (2.3), then $c \leq c'$. The inequality becomes strict when v is not a solution of (2.3).*

We remark that in this results the supersolution v is not required to be monotone.

Corollary 2.3. *Let $f_1 \geq f_2, f_1 \not\equiv f_2$ be two ignition nonlinearities and assume that J is a non negative continuous function of unit mass with finite first moment. Then $c_1 > c_2$ where c_1 and c_2 are the corresponding unique speeds given by Theorem 2.2.*

We also recall some useful results on solutions of (2.3), which can be found in [10, 12].

Lemma 2.4. [10, Lemma 2.1]

Suppose f satisfies either (f1) and J satisfies (j1). Assume $\epsilon \geq 0, c \in \mathbb{R}$ and let $0 \leq u \leq 1$ be an increasing solution of (2.3). Then

$$f(l^\pm) = 0,$$

where l^\pm are the limits of u at $\pm\infty$.

Lemma 2.5. [10, Lemma 2.2] *Let f and J be as in Theorem 2.2. Then following holds*

$$\mu c^2 - \nu |c| \leq 0$$

where the constants μ, ν are defined by

$$\mu := \inf\{\rho, 1 - \rho\} \quad \nu := \int_{\mathbb{R}} J(z)|z| dz$$

Proof of Proposition 2.1. Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of ignition functions which converges pointwise to f and satisfies $\forall n \in \mathbb{N}, f_n \leq f_{n+1} \leq f$. Let (u_n, c_n) denote

the corresponding solution given by Theorem 2.2. By Corollary 2.3 $(c_n)_{n \in \mathbb{N}}$ is an increasing sequence. Next, we see that $c_n \leq \kappa$. Since w satisfies

$$J \star w - w - \kappa w' + f_n(w) \leq 0 \quad \text{in } \mathbb{R}$$

by Theorem 2.2 we get

$$c_n \leq \kappa.$$

Let us observe that we can normalize the sequence of solutions u_n by $u_n(0) = \frac{1}{2}$. Indeed, when $c^* = 0$ since $c_n < c^*$ the solution u_n is smooth. Since any translation of u_n is a solution of the problem and $u_n(-\infty) = 0$, $u_n(+\infty) = 1$ we can normalize it by $u_n(0) = \frac{1}{2}$. When $c^* \neq 0$, since $c_n \rightarrow c^*$ the sequence u_n is smooth for all n sufficiently large. Thus the same normalization can be also taken in this situation.

Since $(u_n)_{n \in \mathbb{N}}$ is an uniformly bounded sequence of increasing functions, using Helly's lemma there exists a subsequence which converges pointwise to a non decreasing function u . Moreover, u satisfies in the distribution sense

$$J \star u - u - c^* u' + f(u) = 0 \quad \text{in } \mathbb{R},$$

and by the monotonicity and the normalization of u_n

$$(2.4) \quad u(x) \leq \frac{1}{2} \quad \text{for all } x \leq 0, \quad u(x) \geq \frac{1}{2} \quad \text{for all } x \geq 0.$$

Observe that when $c^* \neq 0$, using C_{loc}^1 regularity, we get that $u \in C_{loc}^1$ and satisfies the above equation in a strong sense. Otherwise, when $c^* = 0$, a standard argument shows that u satisfies almost everywhere the equation

$$J \star u - u + f(u) = 0.$$

Observe that by (2.4) u is non trivial. It remains to show that u satisfies the right boundary conditions. Now, since u is non decreasing and bounded, the following limits are well defined:

$$l^- := \lim_{x \rightarrow -\infty} u(x)$$

$$l^+ := \lim_{x \rightarrow +\infty} u(x).$$

We get $l^+ = 1$ and $l^- = 0$ using Lemma 2.4, the definition of f and the monotonicity of u .

To finish we need to prove that c^* is independent of the sequence f_n . So consider another sequence \tilde{f}_n of ignition functions such that $\tilde{f}_n \leq \tilde{f}_{n+1} \leq f$ and $\tilde{f}_n \rightarrow f$ pointwise. Let $(\tilde{u}_n, \tilde{c}_n)$ denote the front solution and speed of (2.1) with nonlinearity \tilde{f}_n and let

$$\tilde{c} = \lim_{n \rightarrow \infty} \tilde{c}_n.$$

Since $u = \lim_{n \rightarrow \infty} u_n$ satisfies

$$J \star u - u - c^* u' + \tilde{f}_n(u) \leq 0$$

by Theorem 2.2 we have $\tilde{c}_n \leq c^*$. Hence $\tilde{c} \leq c^*$ and reversing the roles of f_n and \tilde{f}_n we get $c^* \leq \tilde{c}$.

Finally observe that for $c < c^*$ there is no monotone solution to (1.6)-(1.7). Otherwise this solution would be a supersolution of (2.1) with f_n instead of f . By Theorem 2.2 we would have $c_n \leq c$ for all n , which is a contradiction. \square

3. Construction of a solution of (1.5)-(1.7) when J is compactly supported

In this section we deal with the proof of Theorem 1.5. We proceed following the strategy developed in [12]. It is based on the vanishing viscosity technique, a priori estimates, the construction of adequate super and sub-solutions and the characterization of the critical speed obtained in Section 2. Let us first briefly explain how we proceed.

Step 1:

For convenience, let us first rewrite problem (2.3) the following way :

$$(3.1) \quad \begin{cases} \mathcal{M}[u] + f(u) = 0 & \text{in } \mathbb{R} \\ u(-\infty) = 0 \\ u(+\infty) = 1, \end{cases}$$

where the operator \mathcal{M} is defined for a given $\epsilon > 0, c \in \mathbb{R}$ by

$$(3.2) \quad \mathcal{M}[u] = \mathcal{M}(\epsilon, c)u = \epsilon u'' + J \star u - u - cu'.$$

For problem (3.1), for small ϵ , we construct a super solution which is independent of ϵ . More precisely we show the following

Lemma 3.1. *Let J and f be as in Theorem 1.5. Then there exists ϵ_0 and (w, κ) such that $\forall \epsilon \leq \epsilon_0$, (w, κ) is a super-solution of (3.1).*

Step 2: Using the above super solution and a standard approximation scheme, for fixed $0 < \epsilon \leq \epsilon_0$, we prove the following

Proposition 3.2. *Fix $0 < \epsilon \leq \epsilon_0$ and let J and f be as in Theorem 1.5. Then there exists $c^*(\epsilon)$ such that $\forall c \geq c^*(\epsilon)$, there exists an increasing function u_ϵ such that (u_ϵ, c) is a solution of (3.1). Moreover $c^*(\epsilon) \leq \kappa$ where (w, κ) is the super-solution of Lemma 3.1.*

Step 3: We study the singular limit $\epsilon \rightarrow 0$ and prove Theorem 1.5.

Some of the arguments developed in [12], on which this procedure is based, do not use the symmetry of J . Hence in some cases we will skip details in our proofs, making appropriate references to [12].

We divide this section in 3 subsections, each one devoted to one Step.

3.1. Step 1. Existence of a super-solution.

We start with the construction of a super-solution of (3.1) for speeds $c \geq \bar{\kappa}$ for some $\bar{\kappa} > 0$ which is independent of ϵ for $0 < \epsilon \leq 1$.

Lemma 3.3.

Assume J has compact support and let $\epsilon > 0$. There exists a real number $\bar{\kappa} > 0$ and an increasing function $\bar{w} \in C^2(\mathbb{R})$ such that, given any $c \geq \bar{\kappa}$ and $0 < \epsilon \leq 1$

$$\begin{cases} \mathcal{M}[\bar{w}] + f(\bar{w}) \leq 0 & \text{in } \mathbb{R}, \\ \bar{w}(-\infty) = 0, \\ \bar{w}(+\infty) = 1, \end{cases}$$

where $\mathcal{M} = \mathcal{M}(\epsilon, c)$ is defined by (3.2). Furthermore, $\bar{w}(0) = \frac{1}{2}$.

The construction of the super-solution is an adaptation of the one proposed in [12]. The essential difference lies in the computation of the super-solution in a neighborhood of $-\infty$.

Proof. As in [12], fix positive constants N, λ, δ such that $\lambda > \delta$.

Let $\bar{w} \in C^2(\mathbb{R})$ be a positive increasing function satisfying

- $\bar{w}(x) = e^{\lambda x}$ for $x \in (-\infty, -N]$,
- $\bar{w}(x) \leq e^{\lambda x}$ on \mathbb{R} ,
- $\bar{w}(x) = 1 - e^{-\delta x}$ for $x \in [N, +\infty)$,
- $\bar{w}(0) = \frac{1}{2}$.

Let $x_0 = e^{-\lambda N}$ and $x_1 = 1 - e^{-\delta N}$. We have $0 < x_0 < x_1 < 1$.

We now construct a positive function g defined on $(0, 1)$ which satisfies $g(\bar{w}) \geq f(\bar{w})$. Since f is smooth near 0 and 1, we have for c large enough, say $c \geq \kappa_0$,

$$(3.3) \quad \lambda(c - \lambda)s \geq f(s) \quad \text{for } s \in [0, x_0]$$

and

$$(3.4) \quad \delta(c - \delta)(1 - s) \geq f(s) \quad \text{for } s \in [x_1, 1].$$

Therefore we can achieve $g(s) \geq f(s)$ for s in $[0, 1]$, with g defined by:

$$(3.5) \quad g(s) = \begin{cases} \lambda(\kappa_0 - \lambda)s & \text{for } 0 \leq s \leq x_0 \\ l(s) & \text{for } x_0 < s < x_1 \\ \delta(\kappa_0 - \delta)(1 - s) & \text{for } x_1 \leq s \leq 1 \end{cases}$$

where l is any smooth positive function greater than f on $[x_0, x_1]$ such that g is of class C^1 .

According to (3.5), for $x \leq -N$ i.e. for $w \leq e^{-\lambda N}$, we have

$$\begin{aligned} \mathcal{M}[\bar{w}] + g(\bar{w}) &= \epsilon \bar{w}'' + J \star \bar{w} - \bar{w} - c \bar{w}' + g(\bar{w}) \\ &= \epsilon \lambda^2 e^{\lambda x} + J \star \bar{w} - e^{\lambda x} - \lambda c e^{\lambda x} + \lambda(\kappa_0 - \lambda) e^{\lambda x} \\ &\leq \epsilon \lambda^2 e^{\lambda x} + J \star e^{\lambda x} - e^{\lambda x} - \lambda c e^{\lambda x} + \lambda(\kappa_0 - \lambda) e^{\lambda x} \\ &\leq e^{\lambda x} \left[\int_{\mathbb{R}} J(-z) e^{\lambda z} dz - 1 - \lambda(c - \kappa_0) - \lambda^2(1 - \epsilon) \right] \\ &\leq 0, \end{aligned}$$

for c large enough, say

$$c \geq \kappa_1 = \frac{\int_{\mathbb{R}} J(-z) e^{\lambda z} dz - 1 + \lambda \kappa_0 - \lambda^2(1 - \epsilon)}{\lambda}.$$

In the open set $(x_1, +\infty)$, the computation of the super-solution is identical to the one in [12]. So, we end up with

$$\mathcal{M}[\bar{w}] + g(\bar{w}) \leq 0 \quad \text{in } (x_1, +\infty)$$

for c large enough, say $c \geq \kappa_2$.

Therefore, by taking $c \geq \sup\{\kappa_0, \kappa_1, \kappa_2\}$, we achieve

$$\begin{aligned} g(\bar{w}) \geq f(\bar{w}) \quad \text{and} \quad \mathcal{M}[\bar{w}] + g(\bar{w}) \leq 0 \\ \text{for } 0 \leq \bar{w} \leq e^{-\lambda N} \quad \text{and} \quad \bar{w} \geq 1 - e^{-\delta N}. \end{aligned}$$

For the remaining values of \bar{w} , i.e. for $x \in [-N, N]$, $\bar{w}' > 0$ and we may increase c further if necessary, to achieve

$$\mathcal{M}[\bar{w}] + g(\bar{w}) \leq 0 \text{ in } \mathbb{R}.$$

The result follows for

$$\bar{\kappa}(\epsilon) := \sup\{\kappa_0, \kappa_1, \kappa_2, \kappa_3\},$$

where

$$\kappa_3 = \sup_{x \in [-N, N]} \left\{ \frac{\epsilon |\bar{w}''| + |J \star \bar{w} - \bar{w}| + g(\bar{w})}{\bar{w}'} \right\}.$$

□

Now, note that $\bar{\kappa}(\epsilon)$ is a non-decreasing function of ϵ , therefore for all non-negative $\epsilon \leq 1$, $(\bar{w}, \bar{\kappa})$ with $\bar{\kappa} = \bar{\kappa}(1)$, will be a super solution of (3.1), which ends the Step 1.

Remark 3.4. *The above construction of a super-solution also works if we only assume that for some positive λ , the following holds*

$$\int_0^{+\infty} J(-z)e^{\lambda z} dz < +\infty.$$

3.2. Step 2. Construction of a solution when $\epsilon > 0$.

To prove Proposition 3.2 we follow the strategy used in [12] relying on the following approximation scheme.

We first prove existence and uniqueness of a monotone solution for

$$(3.6) \quad \begin{cases} \mathcal{S}[u] + f(u) = -h_r(x) & \text{in } \omega, \\ u(-r) = \theta, \\ u(+\infty) = 1, \end{cases}$$

where $\epsilon > 0$, $r \in \mathbb{R}$, $c \in \mathbb{R}$ and $\theta \in (0, 1)$ are given, and

$$(3.7) \quad \omega = (-r, +\infty),$$

$$(3.8) \quad \mathcal{S}[u] = \mathcal{S}(\epsilon, r, c)[u] = \epsilon u'' + \int_{-r}^{+\infty} J(x-y)u(y)dy - u - cu',$$

$$(3.9) \quad h_r(x) = \theta \int_{-\infty}^{-r} J(x-y)dy.$$

More precisely, we show

Proposition 3.5. *Assume f and J are as in Theorem 1.5. For any $\epsilon > 0$, $\theta \in [0, 1)$ $r > 0$ so that $\text{supp}J \subset (-r, +\infty)$ and $c \in \mathbb{R}$ there exists a unique positive increasing solution u_c of (3.6)*

To prove this proposition we use a construction introduced by one of the authors [9, 10] which consists first to obtain a solution of the following problem:

$$(3.10) \quad \begin{cases} \mathcal{L}[u] + f(u) + h_r + h_R = 0 & \text{for } x \in \Omega \\ u(-r) = \theta, \\ u(+R) = 1, \end{cases}$$

where $\Omega = (-r, +R)$ and $\mathcal{L} = \mathcal{L}(\epsilon, J, r, R, c)$, h_r and h_R are defined by

$$(3.11) \quad \begin{aligned} \mathcal{L}[u] &= \mathcal{L}(\epsilon, J, r, R, c)[u] = \epsilon u'' + \left[\int_{-r}^{+R} J(x-y)u(y)dy - u \right] - cu', \\ h_r(x) &= \theta \int_{-\infty}^{-r} J(x-y)dy, \\ h_R(x) &= \int_{+R}^{+\infty} J(x-y)dy. \end{aligned}$$

Namely, we have,

Proposition 3.6.

Assume f and J are as in Theorem 1.5. For any $\epsilon > 0$, $\theta \in [0, 1)$ $r < R$ so that $\text{supp}J \subset (-r, R)$ and $c \in \mathbb{R}$ there exists a unique positive increasing solution u_c of (3.10).

Proof. The construction of a solution uses the super- and sub-solution iterative scheme presented in [10]. To produce a solution, we just have to construct ordered sub and super-solutions. An easy computation shows that $\underline{u} = \theta$ and $\bar{u} = 1$ are respectively a sub and a super-solution of (3.10). Indeed,

$$\begin{aligned} \mathcal{L}[\underline{u}] + f(\underline{u}) + h_r + h_R &= \int_{-r}^R J(x-y)\theta dy - \theta + \theta \int_{-\infty}^{-r} J(x-y) dy \\ &\quad + \int_R^{+\infty} J(x-y) dy + f(\theta) \\ &= (1 - \theta) \int_R^{+\infty} J(x-y) dy + f(\theta) \geq 0 \end{aligned}$$

and

$$\begin{aligned} \mathcal{L}[\bar{u}] + f(\bar{u}) + h_r + h_R &= \int_{-r}^R J(x-y) dy - 1 + \theta \int_{-\infty}^{-r} J(x-y) dy \\ &\quad + \int_R^{+\infty} J(x-y) dy + f(1) \\ &= (\theta - 1) \int_{-\infty}^{-r} J(x-y) dy \leq 0 \end{aligned}$$

The uniqueness and the monotonicity of such solutions have been already established in [9], so we refer to this reference for interested reader. \square

We are now in a position to prove Proposition 3.5

Proof of Proposition 3.5. Let us now construct a solution of (3.6). Fix $\epsilon > 0$, $c \in \mathbb{R}$ and $r > 0$ such that $\text{supp}(J) \subset \omega$. Let $(R_n)_{n \in \mathbb{N}}$ be a sequence of real which converges to $+\infty$. Since J has compact support, without losing generality we may also assume that $\text{supp}(J) \subset (-r, R_n)$ for all $n \in \mathbb{N}$. Let us denote (u_n, c) the corresponding solution given by Proposition 3.6. Clearly, $h_{R_n} \rightarrow 0$ pointwise, as $n \rightarrow \infty$. Observe now that $(u_n)_{n \in \mathbb{N}}$ is a uniformly bounded sequence of increasing functions. Since $\epsilon > 0$, using local $C^{2,\alpha}$ estimates, up to a subsequence, u_n converges in $C_{loc}^{2,\alpha}$ to a non-decreasing function u . Therefore $u \in C^{2,\alpha}$ and satisfies

$$(3.12) \quad \begin{cases} \epsilon u'' + \int_{-r}^{+\infty} J(x-y)u(y) dy - u - cu' + f(u) + h_r = 0 & \text{in } \omega \\ u(-r) = \theta \end{cases}$$

To complete the construction of the solution, we prove that $u(+\infty) = 1$. Indeed, since u is uniformly bounded and non-decreasing, u achieves its limit at $+\infty$. Using Lemma 2.4 yields $u(+\infty) = 1$. \square

Proof of Proposition 3.2. By Lemma 3.3 there exists $\bar{\kappa}$ and a function \bar{w} which is a supersolution to (3.1) for any $c \geq \bar{\kappa}$ and any $0 < \epsilon \leq 1$. If $c \geq \bar{\kappa}$, following the approach in [12], we can take the limit as $r \rightarrow \infty$ in the problem (3.6) to obtain a solution of (3.1).

Finally one can also verify, see [12], that there exists a monotone solution u_ϵ with the following speed

$$c^*(\epsilon) := \inf\{c \mid (3.1) \text{ admits a monotone solution with speed } c\}.$$

The proof of these claims are straightforward adaptations of [12], since in this reference the author makes no use of the symmetry of J for this part of the proof, and essentially relies on the Maximum principle and Helly's Theorem. We point the interested reader to [12] for the details. \square

Remark 3.7. Note that from the previous comments we get the following uniform estimates

$$\forall 0 < \epsilon \leq \epsilon_0 \quad c^*(\epsilon) \leq \bar{\kappa}$$

3.3. Step 3. Proof of Theorem 1.5 .

We essentially use the ideas introduced in [12].

First, we remark that since J has a compact support, using the super-solution of Step 1, we get from Proposition 2.1 a monotone solution (u, c^*) of (1.5) – (1.7). Furthermore, there exists no monotone solution of (1.5) – (1.7) with speed $c < c^*$ and we have the following characterization :

$$\lim_{k \rightarrow \infty} c_k = c^*,$$

where c_k is the unique speed of fronts associated with an arbitrary sequence of ignition function $(f_k)_{k \in \mathbb{N}}$ which converges pointwise to f and satisfies $\forall k \in \mathbb{N}, f_k \leq f_{k+1} \leq f$.

Also observe that from Remark 3.7 we have a uniform bound from above on $c^*(\epsilon)$.

Lemma 3.8. For all $\epsilon \leq \epsilon_0$ we have $c^*(\epsilon) \leq \bar{\kappa}$.

For any speed $c \geq \bar{\kappa} > 0$, there exists a monotone solution (u_ϵ, c) of (3.1) for any $\epsilon \leq \epsilon_0$. Normalizing the functions by $u_\epsilon(0) = \frac{1}{2}$ and letting $\epsilon \rightarrow 0$, using Helly's Theorem, a priori bounds and some regularity we end up with a solution (u, c) of (1.5) - (1.7). Repeating this limiting process for any speed $c \geq \bar{\kappa}$, we end up with a monotone solution of (1.5) - (1.7) for any speed $c \geq \bar{\kappa}$.

Define now the following critical speed

$$c^{**} = \inf\{c \mid \forall c' \geq c \text{ (1.5) – (1.7) has a positive monotone solution of speed } c'\}.$$

Remark 3.9. Observe that from the uniform bounds we easily see that

$$(3.13) \quad c^{**} \leq \liminf_{\epsilon \rightarrow 0} c^*(\epsilon).$$

Obviously, we have $c^* \leq c^{**} \leq \bar{\kappa}$. To complete the proof of Theorem 1.5, we are then led to prove that $c^{**} = c^*$. To prove this equality, we use some properties of the speed of the following approximated problem

$$(3.14) \quad \begin{cases} \epsilon u'' + J \star u - u - cu' + f\eta_\theta(u) = 0 & \text{in } \mathbb{R}, \\ u(-\infty) = 0, \\ u(+\infty) = 1, \end{cases}$$

where $\theta > 0$, $\eta_\theta(u) = \eta(u/\theta)$ and $\eta \in C^\infty(\mathbb{R})$ is such that

$$0 \leq \eta \leq 1, \quad \eta' \geq 0, \quad \eta(s) = 0 \text{ for } s \leq 1, \quad \eta(s) = 1 \text{ for } s \geq 2.$$

Then η_θ has the following properties

- $\eta_\theta \in C^\infty(\mathbb{R})$,
- $0 \leq \eta_\theta \leq 1$,
- $\eta_\theta(s) \equiv 0$ for $s \leq \theta$ and $\eta_\theta(s) \equiv 1$ for $s \geq 2\theta$,
- if $0 < \theta_1 \leq \theta_2$ then $\eta_{\theta_1} \leq \eta_{\theta_2}$.

For (3.14), we have the following results:

Lemma 3.10. *Let c^θ be the unique speed of front solutions to (2.1) and nonlinearity $f\eta_\theta$. Let $c_\epsilon^\theta, c^*(\epsilon)$ be respectively the unique (minimal) speed solution of (3.1) with the nonlinearity $f\eta_\theta$ and f . Then the following holds :*

- a) For fixed $\theta > 0$, $\lim_{\epsilon \rightarrow 0} c_\epsilon^\theta = c^\theta$.
- b) For fixed ϵ so that $\epsilon_0 \geq \epsilon > 0$, $\lim_{\theta \rightarrow 0} c_\epsilon^\theta = c^*(\epsilon)$.

Assuming for the moment that the above Lemma holds we conclude the proof of Theorem 1.5.

Proof of Theorem 1.5. Assume by contradiction that $c^* < c^{**}$. Then choose c such that $c^* < c < c^{**}$. By (3.13) we may fix $\epsilon_0 > 0$ small such that

$$(3.15) \quad c < c^*(\epsilon) \quad \forall \epsilon \in (0, \epsilon_0).$$

Now consider any sequence $\bar{\theta}_n \rightarrow 0$. Since $c^{\bar{\theta}_n} < c$, using Lemma 3.10 a) there exists $0 < \epsilon_n < \epsilon_0$, $\epsilon_n \rightarrow 0$ such that

$$(3.16) \quad c_{\epsilon_n}^{\bar{\theta}_n} < c.$$

Then, using the continuity of the map $\theta \mapsto c_{\epsilon_n}^\theta$, (3.16), (3.15) and Lemma 3.10 b) we conclude that there exists $0 < \theta_n < \bar{\theta}_n$ such that

$$c = c_{\epsilon_n}^{\theta_n}.$$

Note that $\theta_n \rightarrow 0$. Let u_n be the associated solution to (3.1) with $\epsilon = \epsilon_n$, speed c and nonlinearity $f\eta_{\theta_n}$. We normalize u_n by $u_n(0) = 1/2$. Using Helly's theorem we get a solution \bar{u} of (1.5)-(1.7) with speed c . This contradicts the definition of c^{**} . \square

Proof of Lemma 3.10. The first limit, as $\epsilon \rightarrow 0$ when $\theta > 0$ is fixed, has been already obtained in [10], so we refer to this reference for a detailed proof. The second limit, for fixed $\epsilon > 0$, is obtained using a similar argument as in the proof of Proposition 2.1 to obtain the characterization of c^* . \square

4. Construction of solution in the general case: Proof of Theorem 1.3

In this section we prove Theorem 1.3 and construct solutions of (1.5)-(1.7) only assuming that there exists a super-solution (w, κ) of (1.5)-(1.7).

The construction of solutions uses a standard procedure of approximation of J by kernels J_n with compact support and the characterization of the minimal speed c^* obtained in Section 2.

Let us describe briefly our proof. From Proposition 2.1, there exists a monotone solution (u, c^*) of (1.5)-(1.7) with critical speed. Then we construct monotone solution of (1.5)-(1.7) for any $c > c^*$, $c \neq 0$, using a sequence $(J_n)_{n \in \mathbb{N}}$ of approximated kernels and the same type of arguments developed in the Step 3 of the above section. Let us first construct the approximated kernel and get some uniform lower bounds for c_n^* .

4.1. The approximated kernel and related problems.

First, let j_0 be a positive symmetric function defined by

$$(4.1) \quad j_0(x) = \begin{cases} e^{\frac{1}{x^2-1}} & \text{for } x \in (-1, 1) \\ 0 & \text{elsewhere} \end{cases}$$

Now, let $(\chi_n)_{n \in \mathbb{N}}$ be the following sequence of ‘‘cut-off’’ function:

- $\chi_n \in C_0^\infty(\mathbb{R})$,
- $0 \leq \chi_n \leq 1$,
- $\chi_n(s) \equiv 1$ for $|s| \leq n$ and $\chi_n(s) \equiv 0$ for $|s| \geq 2n$.

Define

$$J_n := \frac{1}{m_n} \left(\frac{j_0}{n} + J(z)\chi_n(z) \right),$$

where $m_n := \frac{1}{n} \int_{\mathbb{R}} j_0(z) dz + \int_{\mathbb{R}} J\chi_n(z) dz$. Observe that since $\int_{\mathbb{R}} j_0 > 0$, J_n is well defined and $J_n(z) \rightarrow J(z)$ pointwise.

Since J_n satisfies the assumption of Theorem 1.5, there exists for each $n \in \mathbb{N}$ a critical speed c_n^* for the problem (4.2) below :

$$(4.2) \quad \begin{cases} J_n \star u - u - cu' + f(u) = 0 & \text{in } \mathbb{R} \\ u(-\infty) = 0 \\ u(+\infty) = 1. \end{cases}$$

Before going to the proof of Theorem 1.3, we prove some *a-priori* estimates on c_n^* . Namely we have the following,

Proposition 4.1.

Let c_n^* be the critical speed defined above, then there exists a positive constant κ_1 such that

$$-\kappa_1 \leq c_n^*.$$

Proof. Let f_θ be a fixed function of ignition type such that $f_\theta \leq f$. Using Theorem 2.2, we have $c_n^\theta \leq c_n^*$. To obtain our desired bound, we just have to bound from below c_n^θ . The later is obtained using Lemma 2.5. Indeed, for each $n \in \mathbb{N}$, we have

$$\mu(c_n^\theta)^2 - \nu_n |c_n^\theta| \leq 0,$$

with $\nu_n := \int_{\mathbb{R}} J_n(z)|z| dz$ and μ is independent of n . Since $\nu_n \leq \bar{\nu} := \sup_{n \in \mathbb{N}} \{\nu_n\} < \infty$, we end up with

$$\mu(c_n^\theta)^2 - \bar{\nu} |c_n^\theta| \leq 0.$$

Hence,

$$|c_n^\theta| \leq \kappa_1.$$

□

Let us also recall some properties of the following approximated problem :

$$(4.3) \quad \begin{cases} J_n \star u - u - cu' + f\eta_\theta(u) = 0 & \text{in } \mathbb{R}, \\ u(-\infty) = 0, \\ u(+\infty) = 1, \end{cases}$$

where $\theta > 0$ and η_θ is such that

- $\eta_\theta \in C_0^\infty(\mathbb{R})$,
- $0 \leq \eta_\theta \leq 1$,
- $\eta_\theta(s) \equiv 0$ for $s \leq \theta$ and $\eta_\theta(s) \equiv 1$ for $s \geq 2\theta$.

For such kind of problem we have,

Lemma 4.2. *Let c^θ and c_n^θ be the unique speed solution of (2.1) with the nonlinearity $f\eta_\theta$ and respectively the kernel J and J_n and let c_n^* be the critical speed solution of (4.3) with the nonlinearity f and the kernel J_n . Then the following holds :*

- a) For fixed θ , $\lim_{n \rightarrow \infty} c_n^\theta = c^\theta$.
- b) For a fixed n , then $\lim_{\theta \rightarrow 0} c_n^\theta = c_n^*$.

Part b) of this Lemma is contained in Proposition 2.1. Part a) can be proved using similar arguments as in Proposition 2.1.

4.2. Construction of the solutions: Proof of Theorem 1.3.

We are now in position to prove Theorem 1.3. From Proposition 2.1, we already know that there exists a travelling front to (1.5)-(1.7) with a critical speed c^* . To complete the proof, we have to construct non decreasing solution for any speed $c \geq c^*$. We emphasize that since (w, κ) is not a super-solution of (1.5)-(1.7) with the approximated kernel J_n , there is no uniform upper bound directly available for the speed c_n^* and the argumentation in the above section cannot directly be applied.

From Proposition 4.1, we have the following dichotomy: either $\liminf (c_n^*)_{n \in \mathbb{N}} < +\infty$ or $\liminf (c_n^*)_{n \in \mathbb{N}} = +\infty$. We prove that in both situations there exists a front solution for any speed $c \geq c^*$.

Case 1: $\liminf (c_n^*)_{n \in \mathbb{N}} < +\infty$

In this case, the same argument as in Theorem 1.5 in Section 3.3 works. Indeed, up to a subsequence $c_n^* \rightarrow \tilde{c}$ and we must have $c^* \leq \tilde{c}$. To prove that $c^* = \tilde{c}$ we proceed as in Section 3.3, using Lemma 4.2 instead of Lemma 3.10.

□

Let now turn our attention to the other situation.

Case 2: $\liminf (c_n^*)_{n \in \mathbb{N}} = +\infty$

In this case $\lim_{n \rightarrow \infty} c_n^* = +\infty$ we argue as follows. Fix $c > c^*$, $c \neq 0$ where c^* is defined by Proposition 2.1. We will show that for such c there is a monotone solution to (1.5)-(1.7). When $c^* \leq 0$ and $c = 0$ then a standard limiting procedure will show that a monotone solution exists with this speed.

Again, by Theorem 2.2 and Proposition 2.1, we have $c^\theta < c^*$ for every positive θ . Therefore,

$$\forall \theta > 0, \quad c^\theta < c^* < c.$$

Fix $\theta > 0$. Since $c_n^\theta \rightarrow c^\theta$, one has on one hand $c_n^\theta < c$ for $n \geq n_0$ for some integer n_0 . On the other hand, $c_n^* \rightarrow +\infty$, thus there exists an integer n_1 such that $c < c_n^*$ for all $n \geq n_1$. Therefore, we may achieve for $n \geq \sup\{n_0, n_1\}$,

$$c_n^\theta < c < c_n^*.$$

From this last inequality, and according to Theorem 2.2 and Lemma 4.2, for each $n \geq \sup\{n_0, n_1\}$ there exists a positive $\theta(n) \leq \theta$ such that $c = c_n^{\theta(n)}$.

Let u_n be the non decreasing solution of (4.2) associated with $\theta(n)$. Since $\theta(n)$ is bounded, we can extract a subsequence still denoted $(\theta(n))_{n \in \mathbb{N}}$ which converges to some $\bar{\theta}$. We claim that

Claim. $\bar{\theta} = 0$

Assume for the moment that the claim is proved. Using the translation invariance, we may assume that for all n , $u_n(0) = \frac{1}{2}$. Using now that u_n is uniformly bounded and Helly's theorem, up to a subsequence $u_n \rightarrow u$ pointwise, where u is a solution of (1.5)-(1.7) with speed c .

In this way we get a non trivial solution of (1.5)-(1.7) for any speed $c \geq c^*$. \square

Let us now turn our attention to the proof of the above claim.

Proof of the Claim. We argue by contradiction. If not, then $\bar{\theta} > 0$ and the speed $c^{\bar{\theta}}$ of the corresponding non decreasing front solution of (2.1) satisfies

$$c^{\bar{\theta}} < c^* < c.$$

Let now consider, u_n the solution associated with $\theta(n)$, normalized by $u_n(0) = \theta(n)$. Using uniform *a priori* estimates, Helly's theorem we can extract a converging sequence of function and get a solution u with speed c of the following

$$J \star u - u - cu' + f_{\bar{\theta}}(u) = 0 \quad \text{in } \mathbb{R}.$$

Using the arguments developed in [10, Section 5.1] to prove Theorem 1.2 of that reference, one can show that u satisfies the boundary conditions

$$u(+\infty) = 1, \quad u(-\infty) = 0.$$

According Proposition 2.1, we get the contradiction

$$c = c^{\bar{\theta}} < c^* < c.$$

Hence $\bar{\theta} = 0$. \square

5. CHARACTERIZATION OF THE MINIMAL SPEED AND ASYMPTOTIC BEHAVIOR

Throughout this section we will assume the hypotheses of Theorem 1.6, namely f satisfies (f1), $f'(0) > 0$, $f \in C^{1,\gamma}$ near 0 and (1.9), and J satisfies (j1), $J \in C^1$ and is compactly supported.

Let us consider the following equation

$$(5.1) \quad \begin{aligned} J \star u - u - cu' + f(u) &= 0 \quad \text{in } \mathbb{R}, \\ \lim_{x \rightarrow -\infty} u(x) &= 0. \end{aligned}$$

We need to establish some estimates on bounded solutions of (5.1) that we constantly use along this section.

Lemma 5.1. *Let u be a no-negative bounded solution of (5.1), then the following holds:*

- (i) $\int_y^x \int_{\mathbb{R}} J(s-t)[u(t) - u(s)] dt ds = \int_0^1 \int_{\mathbb{R}} J(-z)z[u(x+z\eta) - u(y+z\eta)] dz d\eta$
- (ii) $f(u) \in L^1(\mathbb{R})$,
- (iii) $u, J \star u \in L^1(\mathbb{R}^-)$
- (iv) $v(x) := \int_{-\infty}^x u(s) ds$ satisfies $v(x) \leq K(1 + |x|)$ for some positive K and $v(x) \in L^1(\mathbb{R}^-)$.

Proof. We start with the proof of (i). Let $(u_n)_n$ be a sequence of smooth (C^1) function which converge pointwise to u . Using the Fundamental Theorem of Calculus and Fubini's Theorem, we have

$$\begin{aligned} \int_y^x \int_{\mathbb{R}} J(s-t)[u_n(t) - u_n(s)] dt ds &= \int_y^x \int_0^1 \int_{\mathbb{R}} J(-z)z u_n'(s+z\eta) dz d\eta ds \\ &= \int_0^1 \int_{\mathbb{R}} J(-z)z [u_n(x+z\eta) - u_n(y+z\eta)] dz d\eta \end{aligned}$$

Since $|J(-z)z u_n(y+\eta z)| \leq K|J(-z)z| \in L^1(\mathbb{R} \times [0, 1])$ and u_n converges pointwise to u , passing to the limit in the above equation yields

$$\int_y^x \int_{\mathbb{R}} J(s-t)[u(t) - u(s)] dt ds = \int_0^1 \int_{\mathbb{R}} J(-z)z [u(x+z\eta) - u(y+z\eta)] dz d\eta.$$

To obtain (ii), we argue as follow. Integrating (5.1) from y to x , it follows that

$$(5.2) \quad c(u(x) - u(y)) - \int_y^x \int_{\mathbb{R}} J(s-t)[u(t) - u(s)] dt ds = \int_y^x f(u(s)) ds.$$

Using (i), we end up with

$$(5.3) \quad c(u(x) - u(y)) - \int_0^1 \int_{\mathbb{R}} J(-z)z [u(x+z\eta) - u(y+z\eta)] dz d\eta = \int_y^x f(u(s)) ds.$$

Again, since $|J(-z)z u(y+\eta z)| \leq K|J(-z)z| \in L^1(\mathbb{R} \times [0, 1])$, we can pass to the limit $y \rightarrow -\infty$ in the above equation using Lebesgue dominated convergence Theorem. Therefore, we end up with

$$cu(x) - \int_0^1 \int_{\mathbb{R}} J(-z)z u(x+z\eta) dz d\eta = \int_{-\infty}^x f(u(s)) ds.$$

Thus,

$$\int_{-\infty}^x f(u(s)) ds \leq K(|c| + \int_{\mathbb{R}} J(z)|z| dz)$$

which proves (ii). From equation (5.2), we have

$$c(u(x) - u(y)) - \int_y^x f(u(s)) ds + \int_y^x u(s) ds = \int_y^x J \star u(s) ds.$$

Thus $J \star u \in L^1(\mathbb{R}^-)$ will immediately follows from $u \in L^1(\mathbb{R}^-)$ and (ii). Observe now that since $f'(0) > 0$, and $u(-\infty) = 0$, for $x \ll -1$, we have $f(u) > \alpha u$ for some positive constant α . Therefore,

$$\alpha \int_{-\infty}^x u(s) ds \leq \int_{-\infty}^x f(u(s)) ds$$

and (iii) is proved.

To obtain (iv) we argue as follow. From (i-iii), v is a well defined non decreasing function such that $v(-\infty) = 0$. Moreover, v is smooth provide u is continuous. By definition of v , we easily see that $v(x) \leq C(|x| + 1)$ for all $x \in \mathbb{R}$. Indeed, we have

$$v(x) \leq \int_{-\infty}^0 u(s) ds + \int_0^{|x|} u(s) ds \leq K(1 + |x|),$$

where $K = \sup\{\int_{-\infty}^0 u(s) ds; \|u\|_{L^\infty(\mathbb{R})}\}$.

Now, integrating (5.1) on $(-\infty, x)$, we easily see that

$$(5.4) \quad cv'(x) = J \star v(x) - v(x) + \int_{-\infty}^x f(u(s)) ds.$$

Since $f'(0) > 0$ we can choose $R \ll -1$ so that for $s \leq R$, $f(u(s)) \geq \alpha u(s)$ for some $\alpha > 0$. Fixing now $x < R$ and integrating (5.4) between y and x , we obtain

$$(5.5) \quad c(v(x) - v(y)) \geq \int_y^x (J \star v(s) - v(s)) ds + \alpha \int_y^x v(s) ds.$$

Proceeding as above, we get that $v \in L^1(-\infty, R)$. □

Following the idea of Carr and Chmaj [5], we now derive some asymptotic behavior of the non negative bounded solution u of (5.1) More precisely, we show the following

Lemma 5.2. *Let u be a non negative bounded continuous solution of (5.1). Then there exists two positive constants M, β , such that $v(x) = \int_{-\infty}^x u(s) ds$ satisfies :*

$$(5.6) \quad v(x) \leq Me^{\beta x},$$

Proof. The proof uses ideas from [12]. Let first show that some positive constants C, R , we have

$$(5.7) \quad \int_{-\infty}^{-R} v(x) e^{-\beta x} dx < C,$$

for some $\beta > 0$ small.

Consider $R > 0$ and $\beta > 0$ constants to be chosen later. Let $\zeta \in C^\infty(\mathbb{R})$ be a non-negative non-decreasing function such that $\zeta \equiv 0$ in $(-\infty, -2]$ and $\zeta \equiv 1$ in $[-1, \infty)$. For $N \in \mathbb{N}$, let $\zeta_N = \zeta(x/N)$. Multiplying (5.4) by $e^{-\beta x} \zeta_N$ and integrating over \mathbb{R} , we get

$$(5.8) \quad \int_{\mathbb{R}} (J \star v - v)(e^{-\beta x} \zeta_N) dx - \int_{\mathbb{R}} cv'(e^{-\beta x} \zeta_N) dx + \int_{\mathbb{R}} \int_{-\infty}^x f(u(s)) ds (e^{-\beta x} \zeta_N) dx = 0$$

Note that by the monotonicity of ζ_N we have

$$\begin{aligned} \int_{\mathbb{R}} J \star v(x) \zeta_N(x) e^{-\beta x} dx &= \int_{\mathbb{R}} \int_{\mathbb{R}} J(x-y) e^{-\beta x} \zeta_N(x) v(y) dz dy \\ &= \int_{\mathbb{R}} \int_{\mathbb{R}} J(z) e^{-\beta(z+y)} \zeta_N(z+y) v(y) dz dy \\ &\geq \int_{\mathbb{R}} v(y) e^{-\beta y} \left(\int_{-R}^{\infty} J(z) e^{-\beta z} \zeta_N(y-R) dz \right) dy. \end{aligned}$$

Therefore, we have

$$(5.9) \quad \int_{\mathbb{R}} (J \star v - v)(e^{-\beta x} \zeta_N) dx \geq \int v(x) e^{-\beta x} \left(\int_{-R}^{\infty} J(z) e^{-\beta z} dz \zeta_N(x-R) - \zeta_N(x) \right) dx,$$

Let us now choose our adequate $R > 0$. First pick $0 < \alpha < f'(0)$ and $R > 0$ so large that

$$(5.10) \quad f(u)(x) \geq \alpha u(x) \quad \text{for } x \leq -R.$$

Next, one can increase R further if necessary so that $\int_{-R}^{\infty} J(y) dy > (1 - \alpha/2)$. By continuity we obtain for some $\beta_0 > 0$ and all $0 < \beta < \beta_0$,

$$(5.11) \quad \int_{-R}^{\infty} J(y) e^{-\beta y} dy \geq (1 - \alpha/2) e^{\beta R}.$$

Collecting (5.9) and (5.11), we then obtain

$$\begin{aligned} \int_{\mathbb{R}} (J \star v - v)(e^{-\beta x} \zeta_N) &\geq \int_{\mathbb{R}} v(x) e^{-\beta x} \left((1 - \alpha/2) e^{\beta R} \zeta_N(x-R) - \zeta_N(x) \right) dx \\ &\geq (1 - \alpha/2) \int_{\mathbb{R}} v(x+R) e^{-\beta x} \zeta_N(x) dx - \int_{\mathbb{R}} v(x) e^{-\beta x} \zeta_N(x) dx \\ (5.12) \quad &\geq -\alpha/2 \int_{\mathbb{R}} v(x) e^{-\beta x} \zeta_N(x) dx, \end{aligned}$$

where we used the monotone behavior of v in the last inequality.

We now estimate the second term in (5.8) :

$$(5.13) \quad \begin{aligned} \int_{\mathbb{R}} v' \zeta_N e^{-\beta x} dx &= \beta \int_{\mathbb{R}} v \zeta_N e^{-\beta x} - \int_{\mathbb{R}} v \zeta_N' e^{-\beta x} dx \\ &\leq \beta \int_{\mathbb{R}} v \zeta_N e^{-\beta x}. \end{aligned}$$

Finally using (5.10), the last term in (5.8) satisfies

$$(5.14) \quad \begin{aligned} \int_{\mathbb{R}} \left(\int_{-\infty}^x f(u(s)) ds \right) \zeta_N e^{-\beta x} dx &= \int_{-\infty}^{-R} \left(\int_{-\infty}^x f(u(s)) ds \right) \zeta_N e^{-\beta x} dx - C \\ &\geq \alpha \int_{-\infty}^{-R} v \zeta_N e^{-\beta x} dx - C. \end{aligned}$$

By (5.8), (5.12), (5.13) and (5.14), we then obtain

$$\begin{aligned}
|c|\beta \int_{\mathbb{R}} u \zeta_N e^{-\beta x} dx &\geq \alpha \int_{-\infty}^{-R} u \zeta_N e^{-\beta x} dx - C - \alpha/2 \int_{\mathbb{R}} v \zeta_N e^{-\beta x} dx \\
(\alpha/2 - |c|\beta) \int_{-\infty}^{-R} u \zeta_N e^{-\beta x} dx &\leq \tilde{C}.
\end{aligned}$$

Choosing $\beta < \alpha/(2|c|)$ and letting $N \rightarrow \infty$ proves (5.7).

Using the monotonicity of v we can conclude that

$$(5.15) \quad v(x) \leq C e^{\beta x},$$

for some constant C . Indeed, if (5.15) does not hold, then for a sequence $x_n \rightarrow -\infty$ we have $v(x_n) \geq n e^{\beta x_n}$. Extracting a subsequence if necessary, we can assume that $x_{n+1} < x_n - 1$, thus since v is increasing we have

$$\begin{aligned}
\int_{-\infty}^{x_0} v(x) e^{-\beta x} dx &\geq \sum_{n \geq 1} \int_{x_n}^{x_{n-1}} n e^{\beta x_n} e^{-\beta x} dx \\
&\geq \sum_{n \geq 1} n \frac{1 - e^{-\beta(x_n - x_{n-1})}}{\beta} \\
&\geq \sum_{n \geq 1} n \frac{1 - e^{-\beta}}{\beta} = \infty
\end{aligned}$$

which is a contradiction. \square

In the next result we establish that the bounded solution u of (5.1) also decays exponentially as $x \rightarrow -\infty$.

Lemma 5.3. *Suppose that u is bounded solution of (5.1). If for some $M, \beta > 0$ we have that $v(x) \leq M e^{\beta x}$ for all x then there exists $M_1, \alpha > 0$ such that*

$$(5.16) \quad u(x) \leq M_1 e^{\alpha x} \text{ for all } x \in \mathbb{R}.$$

Proof. When $c \neq 0$ then by (5.4) we have the following estimates

$$\begin{aligned}
|c|u(x) &= \left| J \star v - v + \int_{\infty}^x f(u(s)) ds \right| \\
&\leq J \star v + v + \int_{\infty}^x \frac{f(u(s))}{u(s)} u(s) ds \\
&\leq J \star v + (K + 1)v
\end{aligned}$$

where K is the Lipschitz constant of f . Now since

$$J \star v(x) \leq C \int_{\mathbb{R}} J(x-y) e^{\beta y} \leq C' e^{\beta x},$$

we easily see that (5.6) holds.

When $c = 0$ the estimate does not directly comes from (5.4) and we have to distinguish several cases.

Let first observe that for $x < 0$ since u is bounded by some constant C , $J \star u$ satisfies the following

$$\begin{aligned} J \star u(x) &= \int_{-\infty}^{\frac{\alpha}{\beta}x} J(x-y)u(y) dy + \int_{\frac{\alpha}{\beta}x}^{+\infty} J(x-y)u(y) dy \\ &\leq \|J\|_{\infty} \int_{-\infty}^{\frac{\alpha}{\beta}x} u(y) dy + C \int_{x(\frac{\alpha}{\beta}-1)}^{\infty} J(-z) dz \\ &\leq \|J\|_{\infty} v\left(\frac{\alpha}{\beta}x\right) + Ce^{(\beta-\alpha)x} \int_{x(\frac{\alpha}{\beta}-1)}^{\infty} J(-z)e^{\beta z} dz \end{aligned}$$

Choosing $\alpha = \frac{\beta}{2}$ in the above equation, we end up with

$$(5.17) \quad J \star u(x) \leq Ce^{\frac{\beta}{2}x},$$

for some constant C . Observe also that since f is smooth and $f(0) = 0$, we have for small $\epsilon > 0$ and $s > 0$ small,

$$\left| \frac{f(s)}{s} - f'(0) \right| \leq \epsilon.$$

Therefore from (5.1), for $\epsilon > \text{small}$ there exists $K(\epsilon) > 0$ such that for $x < -K(\epsilon)$ we have

$$(5.18) \quad u(1 - f'(0) + \epsilon) \geq J \star u = u\left(1 - \frac{f(u)}{u}\right) \geq u(1 - f'(0) - \epsilon).$$

Observe now that if $f'(0) > 1$, we get a contradiction. Indeed, choose ϵ so that $(1 - f'(0) + \epsilon) < 0$, then we have the following contradiction when $x < -K(\epsilon)$

$$0 > u(1 - f'(0) + \epsilon) \geq J \star u \geq 0.$$

Thus, when $f'(0) > 1$, there is no positive solution of (5.1) with zero speed.

Let us now look at the other cases. Assume now that $f'(0) < 1$ and choose ϵ small so that $(1 - f'(0) - \epsilon) > 0$ then from (5.18) for $x < -K(\epsilon)$ there exists a positive constant C so that

$$u \leq CJ \star u \leq Ce^{\frac{\beta}{2}x}.$$

Finally, when $f'(0) = 1$ recall that f satisfies (1.9). Thus, for $x \ll -1$

$$J \star u(x) = u - f(u) \geq Au^m,$$

where $A > 0$, $m \geq 1$. Using (5.17), yields

$$u \leq \frac{C}{A} e^{\frac{\beta}{2m}x}.$$

□

Remark 5.4. From the above proof, we easily conclude that for any $0 < \alpha < \bar{\alpha}$, where $\bar{\alpha}$ depends only on β and γ , there exists $M_1 > 0$ such that (5.16) holds.

As in [5], for u a solution of (5.1) we define the function $U(\lambda) = \int_{\mathbb{R}} e^{-\lambda x} u(x) dx$ which by Lemma 5.3 is defined and analytic in the strip $0 < \text{Re } \lambda < \alpha$. Note that

$$\int_{\mathbb{R}} J \star u(x) e^{-\lambda x} = \int_{\mathbb{R}} u(y) e^{-\lambda y} dy \int_{\mathbb{R}} J(-z) e^{\lambda z} dz$$

and using integration by parts

$$c \int_{\mathbb{R}} u' e^{-\lambda x} dx = \lambda c \int_{\mathbb{R}} u(y) e^{-\lambda y} dy.$$

Using the above identities, if we multiply (5.1) by $e^{-\lambda x}$ and integrate in \mathbb{R} we obtain

$$(5.19) \quad U(\lambda)(-c\lambda + m(\lambda)) = \int_{\mathbb{R}} e^{-\lambda x} (f'(0)u(x) - f(u(x))) dx,$$

where the function $m(\lambda) = \int_{\mathbb{R}} J(-x) e^{-\lambda x} dx + f'(0) - 1$ is analytic in \mathbb{C} .

Let c^1 be the following quantity

$$c^1 := \min_{\lambda > 0} \frac{1}{\lambda} \left(\int_{\mathbb{R}} J(-x) e^{\lambda x} dx + f'(0) - 1 \right).$$

Proposition 5.5. *If $c < c^1$ then (5.1) does not have any solution.*

Proof. Since $u > 0$ we deduce, from a property of Laplace transform (Theorem 5b, p. 58 [30]) and Lemma 5.3, that the function $U(\lambda)$ is analytic in $0 < \operatorname{Re} \lambda < B$, where $B \geq \alpha$, and $U(\lambda)$ has a singularity at $\lambda = B$. Observe that if $c < c^1$ then for some $\delta > 0$

$$(5.20) \quad -c\lambda + m(\lambda) > \delta\lambda, \quad \text{for all } \lambda > 0.$$

Observe that since $f \in C^{1,\gamma}$ near 0 and using Lemma 5.3 we have that for some constant $C > 0$

$$\begin{aligned} \int_{\mathbb{R}} e^{-\lambda x} |f'(0)u(x) - f(u(x))| dx &= \int_{\infty}^{-K} e^{-\lambda x} |f'(0)u(x) - f(u(x))| dx \\ &\quad + \int_{-K}^{+\infty} e^{-\lambda x} |f'(0)u(x) - f(u(x))| dx \\ &\leq \int_{\infty}^{-K} e^{-\lambda x} |Au^{1+\gamma} + o(u^{1+\gamma})| dx + C \int_{-K}^{+\infty} e^{-\lambda x} u(x) dx \\ &\leq C \int_{\mathbb{R}} e^{-\lambda x} u^{1+\gamma}(x) dx \\ &\leq C \int_{\mathbb{R}} e^{(-\lambda+\gamma\alpha)x} u(x) dx. \end{aligned}$$

From the above computation, it follows that $\int_{\mathbb{R}} e^{-\lambda x} |f'(0)u(x) - f(u(x))| dx$ is analytic in the region $0 < \operatorname{Re} \lambda < B + \gamma\alpha$. Since $\gamma > 0$, using the equation (5.19), we get $U(\lambda)$ defined and analytic for $0 < \operatorname{Re} \lambda < B + \gamma\alpha$. Bootstrapping this argumentation we can extend analytically $U(\lambda)$ to $\operatorname{Re} \lambda > 0$. Then for all $\lambda > 0$

$$\int_{\mathbb{R}} e^{-\lambda x} |f'(0)u(x) - f(u(x))| dx \leq (f'(0) + k) \int_{\mathbb{R}} e^{-\lambda x} u(x) = CU(\lambda).$$

Therefore for all $\lambda > 0$, using (5.19), it follows that $-c\lambda + m(\lambda) \leq C$ contradicting (5.20). \square

Remark 5.6. We should point out that the above proposition holds as well if the kernel J instead of being compactly supported, is only assumed to satisfy:

$$\exists M, \lambda_0 > 0 \quad \text{such that} \quad \int_0^{+\infty} J(-x)e^{\lambda_0 x} \leq M.$$

Let us first establish the exact asymptotic behavior, as $x \rightarrow -\infty$, of a solution u of (5.1). We proceed as follows. First, we obtain the exact behavior of $v = \int_{-\infty}^x u(s)ds$, proceeding as in [5] and then we conclude the behavior of u .

For $c \geq c^1$ we denote $\lambda(c)$ the unique minimal $\lambda > 0$ such that $-c\lambda + m(\lambda) = 0$. It can be easily verified that $\lambda(c)$ is a simple root of $-c\lambda + m(\lambda)$ if $c > c^1$, and it is a double root when $c = c^1$.

Proof of Theorem 1.6. Since there is a monotone solution (u, c^*) of (1.5)-(1.7) with critical speed, it is a bounded solution of (5.1). Thus by Proposition 5.5 $c^* \geq c^1$.

It remains to prove (1.10) and (1.11). The proof follows from a modified version of Ikehara's Theorem (see [30]). We define $F(\lambda) = \int_{-\infty}^0 v(y)e^{-\lambda y}$. Since v is monotone, we can obtain the appropriate asymptotic behavior of v if F has the representation

$$(5.21) \quad F(\lambda) = \frac{H(\lambda)}{(\lambda - \alpha)^{k+1}},$$

with H analytic in the strip $0 < \operatorname{Re} \lambda \leq \alpha$, and $k = 0$ when $c > c^*$, $k = 1$ when $c = c^*$.

Using (5.4), we have that

$$\int_{-\infty}^0 v(x)e^{-\lambda x} dx = \frac{\int_{-\infty}^{\infty} \int_{-\infty}^x f(u(s)) - f'(0)u(s) ds e^{-\lambda x} dx}{c\lambda - m(\lambda)} - \int_0^{\infty} v(x)e^{-\lambda x},$$

thus, using that either $c \neq 0$ or $f'(0) < 1$ holds, we have that by Lemma 5.3, (5.21) holds replacing u by v with $\alpha = \lambda(c)$ described above, since it can be checked that $-c\lambda + m(\lambda)$ has only two real roots which are simple when $c > c^1$ and double when $c = c^1$.

It remains to conclude that (5.21) holds for u . First suppose that $c = c^1$ and denote $\lambda = \lambda(c^1)$. If $c \neq 0$ then using (5.4) we have that

$$cu = J \star v(x) - (1 - f'(0))v(x) + \int_{-\infty}^x f(u(s)) - f'(0)u(s) ds,$$

By Remark 5.4 and since f is $C^{1,\gamma}$ near 0 we have that

$$(5.22) \quad \frac{\int_{-\infty}^x f(u(s)) - f'(0)u(s) ds}{|x|e^{-\lambda(c^1)x}} \rightarrow 0,$$

as $|x| \rightarrow -\infty$. Therefore, we just have to prove that

$$(5.23) \quad \lim_{x \rightarrow -\infty} \frac{J \star v(x) - (1 - f'(0))v(x)}{|x|e^{\lambda(c^1)x}} = L \neq 0.$$

Observe that since v satisfies (1.10) we have that for $\eta = \lim_{x \rightarrow -\infty} \frac{v(x)}{|x|e^{\lambda(c^1)x}}$ and $\operatorname{supp} J \subset [-k, k]$ we have

$$\frac{J \star v}{|x|e^{\lambda(c^1)x}} = \frac{1}{|x|} \int_{-k}^k J(-z)(\eta + O(1/x))e^{\lambda(c^1)z}(|x| + z) dz,$$

therefore

$$\frac{J \star v(x) - (1 - f'(0))v(x)}{|x|e^{\lambda(c^1)x}} \rightarrow \eta m(\lambda(c^1)) = \eta c^1 \lambda(c^1) \neq 0,$$

which gives the desired result.

When $c^1 = 0$, we proceed in a slightly different way. Observe that in this case $f'(0) < 1$

$$(5.24) \quad (1 - f'(0))u = J \star u + f(u) - f'(0)u,$$

and by Remark 5.4 and since $f \in C^{1,\gamma}$ near 0 we have that (5.22) holds. Also, by (j2) we have that $J \star u = J' \star v$ and

$$\begin{aligned} \frac{J' \star v}{|x|e^{\lambda(c^1)x}} &= \frac{1}{|x|} \int_{-k}^k J'(-z)(\eta + O(1/x))e^{\lambda(c^1)z}(|x| + z)dz \\ &= \eta \int_{\mathbb{R}} J(-z)e^{\lambda(c^1)z} dz + O(1/x), \end{aligned}$$

with $\eta > 0$ as above. Hence, we obtain the desired result.

Finally, the case $c > c^1$ is analogous. \square

Proof of Corollary 1.7. Observe now that in the case of a KPP nonlinearity f , the function $w := e^{\lambda x}$ is a super-solution of (1.5)-(1.7), provided that $\lambda > 0$ is chosen such that $-c\lambda + m(\lambda) = 0$. The existence of such $\lambda > 0$ is guaranteed since $c \geq c^1$. The existence of a monotone travelling wave for any $c \geq c^1$ is then provided by Theorem 1.3. Therefore $c^* \leq c^1$ and we conclude $c^* = c^1$. \square

6. UNIQUENESS OF THE PROFILE

In this section we deal with the uniqueness up to translation of solution of (1.5)–(1.7). Our proof follows ideas of [8] and is mainly based on the sliding methods introduced by Berestycki and Nirenberg [3].

In the sequel, given a function $u : \mathbb{R} \rightarrow \mathbb{R}$ and $\tau \in \mathbb{R}$ we define its translation by τ as

$$(6.1) \quad u_\tau(x) = u(x + \tau)$$

and sometimes we shall write $u^\tau(x) = u(\tau + x)$.

Let L denote the operator

$$Lu = J \star u - u - cu'.$$

Proposition 6.1. (*Nonlinear Comparison Principle*)

Let J satisfy (j1), (MP) and let f be a monostable nonlinearity so that $f'(1) < 0$. Let u and v be two continuous functions in \mathbb{R} such that

$$(6.2) \quad Lu + f(u) \leq 0 \quad \text{on } \mathbb{R}$$

$$(6.3) \quad Lv + f(v) \geq 0 \quad \text{on } \mathbb{R}$$

$$(6.4) \quad \lim_{x \rightarrow -\infty} u(x) \geq 0, \quad \lim_{x \rightarrow -\infty} v(x) \leq 0$$

$$(6.5) \quad \lim_{x \rightarrow +\infty} u(x) \geq 1, \quad \lim_{x \rightarrow +\infty} v(x) \leq 1.$$

Assume further that either u or v is monotone and that $u \geq v$ in some interval $(-\infty, K)$. Then there exists $\tau \in \mathbb{R}$ such that $u_\tau \geq v$ in \mathbb{R} . Moreover, either $u_\tau > v$ in \mathbb{R} or $u_\tau \equiv v$.

Remark 6.2. Observe that by the Maximum Principle and since $f(s) \geq 0 \quad \forall s \leq 0$, the supersolution u is necessarily positive. Similarly, since $f(s) \leq 0 \quad \forall s \geq 1$, the Maximum Principle implies that $v < 1$.

Proof of Proposition 6.1. Note that if $\inf_{\mathbb{R}} u \geq \sup_{\mathbb{R}} v$, the theorem trivially holds. In the sequel, we assume that $\inf_{\mathbb{R}} u < \sup_{\mathbb{R}} v$.

Let $\epsilon > 0$ be such that

$$(6.6) \quad f'(p) \leq 0 \quad \text{for } 1 - \epsilon < p < 1.$$

Now fix $0 < \delta \leq \frac{\epsilon}{2}$ and choose $M > 0$ sufficiently large so that

$$(6.7) \quad 1 - u(x) < \frac{\delta}{2} \quad \forall x > M$$

$$(6.8) \quad v(x) < \frac{\delta}{2} \quad \forall x < -M$$

$$(6.9) \quad \text{and } v(x) \leq u(x) \quad \forall x < -M.$$

Step 1. There exists a constant D such that for every $b \geq D$

$$(6.10) \quad u(x + b) > v(x) \quad \forall x \in [-M - 1 - b, M + 1].$$

Indeed, since $u > 0$ in \mathbb{R} and $\lim_{x \rightarrow +\infty} u(x) \geq 1$ we have

$$c_0 := \inf_{[-M-1, \infty)} u > 0$$

Since $\lim_{x \rightarrow -\infty} v(x) \leq 0$ there is $L > 0$ large such that

$$v(x) < c_0 \quad \forall x \leq -L.$$

Then for all $b > 0$

$$u(x + b) > v(x) \quad \forall x \in [M - 1 - b, -L].$$

Now, since $\sup_{[-L, M+1]} v < 1$ and $\lim_{x \rightarrow +\infty} u(x) \geq 1$ we deduce (6.10).

Step 2. There exists $b \geq D$ such that

$$(6.11) \quad u(x + b) + \frac{\delta}{2} > v(x) \quad \forall x \in \mathbb{R}.$$

If not then we have,

$$(6.12) \quad \forall b \geq D \quad \text{there exists } x(b) \text{ such that } u(x(b) + b) + \frac{\delta}{2} \leq v(x(b)).$$

Since u is nonnegative and v satisfies (6.4) there exists a positive constant A such that

$$(6.13) \quad u(x + b) + \frac{\delta}{2} > v(x) \quad \text{for all } b > 0 \text{ and } x \leq -A.$$

Take now a sequence $(b_n)_{n \in \mathbb{N}}$ which tends to $+\infty$. Let $x(b_n)$ be the point defined by (6.12). Thus we have for that sequence

$$(6.14) \quad u(x(b_n) + b_n) + \frac{\delta}{2} \leq v(x(b_n)).$$

According to (6.13) we have $x(b_n) \geq -A$. Therefore the sequence $x(b_n) + b_n$ converges to $+\infty$. Pass to the limit in (6.14) to get

$$1 + \frac{\delta}{2} \leq \lim_{n \rightarrow +\infty} u(x(b_n) + b_n) + \frac{\delta}{2} \leq \limsup_{n \rightarrow +\infty} v(x(b_n)) \leq 1,$$

which is a contradiction. This proves our claim (6.11).

Step 3 We observe that as a consequence of (6.10) and (6.11), and using that either u or v is monotone we in fact have

$$(6.15) \quad \begin{aligned} u(x+b) &\geq v(x) \quad \forall x \leq M+1 \\ u(x+b) + \frac{\delta}{2} &> v(x) \quad \forall x \geq M+1. \end{aligned}$$

Indeed, it only remains to verify that $u(x+b) > v(x)$ for $x \leq M-1-b$. If u is monotone from (6.9) we have $u(x+b) > u(x) > v(x)$ for $x < -M$. If v is monotone $u(x) > v(x) > v(x-b)$ for $x < -M$.

Step 4. Now we claim that

$$(6.16) \quad u(x+b) \geq v(x) \quad \forall x \in \mathbb{R}.$$

To prove this, consider

$$(6.17) \quad a^* = \inf\{a > 0 \mid u(x+b) + a \geq v(x) \forall x \in \mathbb{R}\}$$

which is well defined by (6.11).

If $a^* = 0$ then (6.16) follows. Suppose $a^* > 0$. Then, since

$$\lim_{x \rightarrow \pm\infty} u(x+b) + a^* - v(x) \geq a^* > 0,$$

there exists $x_0 \in \mathbb{R}$ such that $u(x_0+b) + a^* = v(x_0)$.

Let $w(x) := u(x+b) + a^* - v(x)$ and note that

$$(6.18) \quad 0 = w(x_0) = \min_{\mathbb{R}} w(x).$$

Observe that w also satisfies the following equations:

$$(6.19) \quad Lw \leq f(v(x)) - f(u(x+b))$$

$$(6.20) \quad w(+\infty) \geq a^*$$

$$(6.21) \quad w(-\infty) \geq a^*.$$

Since $w \geq 0$, $w \not\equiv 0$ using the strong maximum principle at x_0 we have

$$(6.22) \quad Lw(x_0) > 0.$$

By (6.15) we necessarily have $x_0 > M+1$.

At x_0 we have

$$(6.23) \quad f(u(x_0+b) + a^*) - f(u(x_0+b)) \leq 0,$$

since f is non-increasing for $s \geq 1 - \epsilon$, $a^* > 0$ and $1 - \epsilon < 1 - \frac{\delta}{2} \leq u$ for $x > M$. Combining (6.19), (6.22) and (6.23) yields the contradiction

$$0 < Lw(x_0) \leq f(u(x_0+b) + a^*) - f(u(x_0+b)) \leq 0.$$

Step 5. Finally it remains to prove that either $u_\tau > v$ or $u_\tau \equiv v$. Let $w := u_\tau - v$, then either $w > 0$ or $w(x_0) = 0$ at some point $x_0 \in \mathbb{R}$. In the latter case we have $w(x) \geq w(x_0) = 0$ and

$$(6.24) \quad 0 \leq Lw(x_0) \leq f(v(x_0)) - f(u(x_0+\tau)) = f(v(x_0)) - f(v(x_0)) = 0.$$

Then using the maximum principle, we obtain $w \equiv 0$, which means $u_\tau \equiv v$. \square

Proposition 6.3. *Let J satisfy (j1), (MP) and let f be a monostable nonlinearity so that $f'(1) < 0$. Let u_1 and u_2 be respectively super and sub-solutions of (1.5)-(1.7) which are continuous. If $u_1 \geq u_2$ in some interval $(-\infty, K)$ and either u_1 or u_2 is monotone then $u_1 \geq u_2$ everywhere. Moreover either $u_1 > u_2$ or $u_1 \equiv u_2$.*

Proof. Assume first that $\inf_{\mathbb{R}} u_1 < \sup_{\mathbb{R}} u_2$. Otherwise there is nothing to prove. Without losing generality we can assume that u_1 is monotonic. Using Theorem 6.1, $u_1^\tau \geq u_2$ for some $\tau \in \mathbb{R}$, so the following quantity is well defined

$$\tau^* := \inf\{\tau \in \mathbb{R} | u_1^\tau \geq u_2\}$$

We claim that

$$(6.25) \quad \tau^* \leq 0$$

Observe that by showing that $\tau^* \leq 0$, we end the proof. To prove (6.25) we argue by contradiction. Assume that $\tau^* > 0$, then since u_i are a continuous functions, we will have $u_1^{\tau^*} \geq u_2$ in \mathbb{R} . Let $w := u_1^{\tau^*} - u_2 \geq 0$. Since $\tau^* > 0$ and u_1 is monotone then $w > 0$ in $(-\infty, K)$. Now observe that $w > 0$ in \mathbb{R} or $w(x_0) = 0$ for some point x_0 in \mathbb{R} . In the latter case

$$0 \leq (J \star w - w)(x_0) \leq f(u_2(x_0)) - f(u_1^{\tau^*}(x_0)) = 0.$$

Thus, using the maximum principle, $w \equiv 0$, which contradicts that $w > 0$ in $(-\infty, K)$. Now since u_1 is monotonic and $\tau^* > 0$ for small $\epsilon > 0$, we have $u_1^{\tau^* - \epsilon} > u_2$ in $(-\infty, M)$. Arguing as in Step 4 of the proof of Proposition 6.1 we deduce $u_1^{\tau^* - \epsilon} > u_2$ in \mathbb{R} which contradicts the definition of τ^* . \square

Remark 6.4. *With minor modifications the proofs of the Propositions 6.1 and 6.3 hold if only one of the functions u_1 or u_2 is continuous. For the proof of this statement we need the strong maximum principle for solutions in L^∞ , which can be found in [11]:*

Theorem 6.5. *Assume J satisfies (j1), (MP) and let $c \in L^\infty(\mathbb{R})$. If $u \in L^\infty(\mathbb{R})$ satisfies $u \leq 0$ a.e. and $J \star u - u + c(x)u \geq 0$ a.e. in \mathbb{R} , then $\text{ess sup}_K u < 0$ for all compact $K \subset \mathbb{R}$ or $u = 0$ a.e. in \mathbb{R} .*

Proof of Theorem 1.8. The case of $c \neq c^1$ and $c = c^1$ being similar, we present only the case $c \neq c^1$. Let u_1 and u_2 be two solution of (1.5)-(1.7) with the same speed $c \neq 0$. Since $c \neq 0$ the functions u_i are uniformly continuous. From Theorem 1.3, we can assume that u_1 is a monotonic function. Since, u_i solve the same equation and u_1 is monotone, using the translation invariance of the equation and (1.11) we see that up to a translation

$$(6.26) \quad u_1 = e^{\lambda(c)x} + o(e^{\lambda(c)x}) \quad \text{as } x \rightarrow -\infty$$

$$(6.27) \quad u_2 = e^{\lambda(c)x} + o(e^{\lambda(c)x}) \quad \text{as } x \rightarrow -\infty.$$

Let us first recall the following notation, $u^\tau(\cdot) := u(\cdot + \tau)$. Then, by monotonicity of u_1 and (6.26)- (6.27) for some positive τ we have $u_1^\tau \geq u_2$ in some interval $(-\infty, -K)$. Using Proposition 6.3, it follows that $u_1^\tau \geq u_2$ for possibly a new τ . Define now the following quantity:

$$\tau^* := \inf\{\tau > 0 | u_1^\tau \geq u_2\}$$

Observe that from the above argument τ^* is well defined. We claim

Claim 6.6. $\tau^* = 0$.

Observe that proving the claim ends the proof of the uniqueness up to translation of the solution. Indeed, assume for a moment that the claim is proved then we end up with $u_1 \geq u_2$. Observe now that in the above argumentation the role of u_1 and u_2 can be interchanged, so we easily see that we have $u_1 \leq u_2 \leq u_1$ which ends the proof of the uniqueness. \square

Let us now prove the Claim.

Proof of the Claim. If not, then $\tau^* > 0$. Let $w := u_1^{\tau^*} - u_2 \geq 0$. Then either there exists a point x_0 where $w(x_0) = 0$ or $w > 0$. In the first case, at x_0 , w satisfies:

$$0 \leq J \star w(x_0) - w(x_0) = f(u_2(x_0)) - f(u_1^{\tau^*}(x_0)) = 0$$

Using the strong maximum principle, it follows that $w \equiv 0$. Thus $u_1^{\tau^*} \equiv u_2$, which contradicts (6.26)–(6.27). Therefore, $u_1^{\tau^*} > u_2$. Using (6.26), since $\tau^* > 0$ we have for $u_1^{\tau^*}$ the following behavior near $-\infty$.

$$u_1^{\tau^*} := e^{\tau^*} e^{\lambda(c)x} + o(e^{\lambda(c)x}).$$

Therefore, for some $\epsilon > 0$ small, we still have $u_1^{\tau^* - \epsilon} \geq u_2$ in some neighborhood $(-\infty, -K)$ of $-\infty$. Using Theorem 6.3, we end up with $u_1^{\tau^* - \epsilon} \geq u_2$ everywhere, contradicting the definition of τ^* . Hence, $\tau^* = 0$. \square

Regarding Theorem 1.9 we need the following result:

Lemma 6.7. *Assume that J and f satisfy (j1), (j2), (MP) and (f1), (f2) respectively. Let $0 \leq u \leq 1$ be a solution to (1.5).*

a) Then

$$\lim_{x \rightarrow -\infty} u(x) = 0 \quad \text{or} \quad \lim_{x \rightarrow -\infty} u(x) = 1,$$

and

$$\lim_{x \rightarrow \infty} u(x) = 0 \quad \text{or} \quad \lim_{x \rightarrow \infty} u(x) = 1.$$

b) If $u(-\infty) = 1$ and $u(+\infty) = 1$ then $u \equiv 1$.

Note that in this lemma we do not assume that u is continuous.

Proof.

a) Let $0 \leq u \leq 1$ be a solution to (1.12). We first note that by (5.3) any bounded solution u of (1.5) satisfies

$$(6.28) \quad \int_{-\infty}^{\infty} f(u) du < \infty.$$

Let $g(u) = u - f(u)$ and note that

$$(6.29) \quad J \star u = g(u) \quad \text{in } \mathbb{R},$$

and that the hypotheses on f imply $g'(u) \geq g'(0)$ and $g(u) \leq u$ for $u \in [0, 1]$.

If $f'(0) < 1$ then $g'(0) > 0$ and then g is strictly increasing. This together with (6.29) implies that u is uniformly continuous and using (6.28) we see that $u(-\infty) = 0$ or $u(-\infty) = 1$ and the same at $+\infty$ which is the desired conclusion. Therefore in the sequel we assume $f'(0) \geq 1$, that is, $g'(0) \leq 0$.

Since both limits at $-\infty$ and $+\infty$ are analogous we concentrate on the case $x \rightarrow -\infty$.

We will establish the conclusion of part a) by proving

$$(6.30) \quad \liminf_{x \rightarrow -\infty} J \star u(x) = 0 \implies \lim_{x \rightarrow -\infty} u(x) = 0,$$

and

$$(6.31) \quad \liminf_{x \rightarrow -\infty} J \star u(x) > 0 \implies \lim_{x \rightarrow -\infty} u(x) = 1.$$

We start with (6.30). Suppose that $f'(0) > 1$. Then there is $\delta > 0$ such that $g(u) < 0$ for $u \in (0, \delta)$ and from (6.29) we deduce that $u(x) \geq \delta$ for all x , so regarding (6.30) there is nothing to prove.

Suppose $f'(0) = 1$. Then g is non-decreasing and by (1.9) we have, for some $A > 0, m \geq 1, \delta_1 > 0$

$$(6.32) \quad g(u) \geq Au^m \quad \forall 0 \leq u \leq \delta_1.$$

Assume that $\liminf_{x \rightarrow -\infty} J \star u(x) = 0$ and let us show first that

$$(6.33) \quad \lim_{x \rightarrow -\infty} J \star u(x) = 0.$$

Otherwise, set $\bar{l} = \limsup_{x \rightarrow -\infty} J \star u(x) > 0$. Choose $l \in (0, \bar{l})$ such that $g'(l) > 0$ and then pick a sequence $x_n \rightarrow -\infty$ such that $J \star u(x_n) = g(l)$ for all n . Then there is some $\sigma > 0$ such that for $x \in (x_n - \sigma, x_n + \sigma)$ we have $f(u(x)) \geq c > 0$ for some uniform c . This contradicts (6.28) and we deduce (6.33). This combined with (6.32) implies that $\lim_{x \rightarrow -\infty} u(x) = 0$, and this establishes (6.30).

We prove now (6.31). Let us assume

$$\underline{l} := \liminf_{x \rightarrow -\infty} J \star u(x) > 0.$$

Since $J \star u = g(u) \leq u$ it is enough to show that

$$(6.34) \quad \lim_{x \rightarrow -\infty} J \star u(x) = 1.$$

Assume the contrary, that is

$$(6.35) \quad 0 < \underline{l} < 1.$$

Observe that

$$\liminf_{x \rightarrow -\infty} u(x) > 0.$$

This is direct if $f'(0) > 1$ and follows from (6.29), (6.32) and $\underline{l} > 0$ if $f'(0) = 1$. Therefore $\limsup_{x \rightarrow -\infty} u(x) = 1$, otherwise (6.28) can not hold. Hence

$$(6.36) \quad \limsup_{x \rightarrow -\infty} J \star u(x) = 1.$$

Chose now $\alpha \in (\underline{l}, 1)$ a regular value of the function g . By (6.35), (6.36) and the continuity of $J \star u$ there exists a sequence $x_n \rightarrow -\infty$ such that $J \star u(x_n) = \alpha$. Note that the set $\{u \in [0, 1] / g(u) = \alpha\}$ is discrete and hence finite and does not contain 0 nor 1. Hence, for sufficiently small $\epsilon > 0$ we have $\{u \in [0, 1] / \alpha - \epsilon < g(u) < \alpha + \epsilon\} \subseteq [\epsilon, 1 - \epsilon]$. Since $J \star u$ is uniformly continuous there is $\sigma > 0$ such that for $x \in (x_n - \sigma, x_n + \sigma)$ we have $\epsilon \leq u(x_n) \leq 1 - \epsilon$. This contradicts the integrability condition (6.28), and we deduce the validity of (6.34).

b) Assume that $\lim_{x \rightarrow \infty} u(x) = \lim_{x \rightarrow -\infty} u(x) = 1$ and set $\gamma^* = \sup\{0 < \gamma < 1 / u > \gamma\}$. For the sake of contradiction assume that u is nonconstant. Then

$0 < \gamma^* < 1$. Since $f(\gamma^*) > 0$ we have that $v = u - \gamma^* \geq 0$ satisfies

$$(6.37) \quad J \star v - v - cv' + \frac{f(u) - f(\gamma^*)}{u - \gamma^*} (u - \gamma^*) < 0.$$

If $c \neq 0$ then v reaches its global minimum at some $x_0 \in \mathbb{R}$ which satisfies $v(x_0) = 0$. Thus, evaluating (6.37) at x_0 we obtain a contradiction. If $c = 0$ we reach again a contradiction applying Theorem 6.5. \square

Proof of Theorem 1.9. Assume $0 \leq u \leq 1$ is a solution of (1.12) such that $u \not\equiv 0$ and $u \not\equiv 1$. By Lemma 6.7 $u(-\infty) = 0$ or $u(+\infty) = 0$. Then we may apply Theorem 1.6 and deduce the exact asymptotic behavior of u at either $-\infty$ or $+\infty$ and that $c^* \leq 0$ or $c_* \leq 0$. Let u_0 denote a non-decreasing travelling wave with speed $c = 0$ if $c^* \leq 0$ or a non-increasing one if $c_* \leq 0$. Then, by slightly modifying the proof of Theorem 2.1 in [5] we deduce that for a suitable translation we have $u^\tau \equiv u_0$. In particular the profile of the travelling wave u_0 is unique. \square

Next we address the issues of non-uniqueness and discontinuities of solutions when $c = 0$. We consider f such that

$$(6.38) \quad f \text{ is smooth, } 0 < f'(0) < 1, f'(1) < 0 \text{ and } f \text{ is KPP.}$$

We are interested in the case where $u - f(u)$ is not monotone, and for simplicity we shall assume that setting

$$g(u) = u - f(u)$$

there exists $0 < \alpha < \beta < 1$ such that

$$(6.39) \quad \begin{aligned} g'(u) &> 0 \quad \forall u \in [0, \alpha] \cup (\beta, 1] \\ g'(u) &< 0 \quad \forall u \in (\alpha, \beta). \end{aligned}$$

Proposition 6.8. *Assume f satisfies (6.38), (6.39). Then there exists J such that no solution of (1.5)-(1.7) is continuous, and this problem admits infinitely many solutions.*

Proof. Let us choose $J \in C^1$, with compact support and satisfying (j1) and (MP), and such that $c^1 \leq 0$. Then by Corollary 1.7 we have $c^* = c^1 \leq 0$. Thus there exists a monotone travelling wave solution u_1 of (1.5)-(1.7) with speed $c = 0$. If (1.5)-(1.7) has a continuous solution u_2 , then by Theorem 1.8 and Remark 6.4 we have $u_1 \equiv u_2$. Hence u_1 is monotone and continuous. Then $J \star u_1$ is monotone which implies that $u_1 - f(u_1)$ is monotone in \mathbb{R} . This is impossible if u_1 is continuous and $u - f(u)$ is not monotone.

For the construction of infinitely many solutions we follow closely the work of [2]. Since $g'(0) > 0$ and $g'(1) > 0$ there are $a < b$ such that

$$\begin{aligned} g \text{ is increasing in } [0, a], g \text{ is increasing in } [b, 1] \\ g(a) = g(b) \text{ and } g \text{ is not monotone in } [a, b]. \end{aligned}$$

Define

$$\tilde{g}(u) = \begin{cases} g(u) & \text{if } u \in [0, a] \text{ or } u \in [b, 1] \\ g(a) & \text{if } u \in [a, b] \end{cases}$$

Let $g_n : [0, 1] \rightarrow \mathbb{R}$ be smooth such that $g_n \rightarrow g$ uniformly in $[0, 1]$, $g_n \equiv g$ in a neighborhood of 0 and 1, $g'_n > 0$ and $u - g_n(u)$ is KPP. Then by Corollary 1.7 the problem (1.5)- (1.7) with nonlinearity $f_n = u - g_n(u)$ has critical speed $c^* \leq 0$ independent of n , and hence there exists a monotone solution u_n

$$J \star u_n = g_n(u_n), \quad u_n(-\infty) = 0, \quad u_n(+\infty) = 1.$$

Notice that any solution to this problem is continuous and hence we may choose

$$u_n(0) = a.$$

By Helly's theorem there is a subsequence which converges pointwise to a solution u of the following problem

$$J \star u = \tilde{g}(u) \quad \text{in } \mathbb{R}.$$

Remark that $u(0) = a$, and $u(-\infty) = 0$, $u(+\infty) = 1$ by Lemma 2.4. Note that u is continuous in $(-\infty, 0]$ since $u \leq a$ in $(-\infty, 0]$ and g is strictly increasing in $[0, a]$.

We will show that u has a discontinuity at 0 and $u(0^+) = b$. As in [2], choose $\delta_n > 0$ such that $u_n(\delta_n) = b$. Let $\delta = \liminf \delta_n$ and note that $u \geq b$ in (δ, ∞) . Let us show that $\delta = 0$. If not, then $\tilde{g}(u(x)) = g(a)$ for $x \in (0, \delta)$ and this implies $J \star u = \text{const}$ in $(0, \delta)$. Then for $0 < \tau < \delta/2$ we have $J \star (u - u(\cdot - \tau)) \geq 0$ and vanishes in a nonempty interval. By the maximum principle $u \equiv u(\cdot - \tau)$ and this implies that u is constant, which is a contradiction. Thus $\delta = 0$ and u has a jump discontinuity at 0. Hence u is a solution to (1.5)- (1.7). We conclude that $u(0^+) = b$ because $J \star u$ is continuous. \square

Acknowledgements. This work has been partly supported by the Ecos-Conicyt project C05E04, Fondecyt 1050754, Fondecyt 1050725, Nucleus Millenium P04-069-F, Information and Randomness and by FONDAP grant for Applied Mathematics, Chile. This works also has been partly supported by the Max Planck Institut for mathematics in the science of Leipzig.

REFERENCES

- [1] D. G. Aronson, and H. Weinberger, *Multidimensional nonlinear diffusion arising in population genetics*, Adv. in Math. 30, (1978), no. 1, 33–76.
- [2] P. Bates, P. Fife, X. Ren and X. Wang, *Traveling waves in a convolution model for phase transitions*, Arch. Rational Mech. Anal. 138, (1997), no. 2, 105–136.
- [3] Berestycki, H. and Nirenberg, L., *On the method of moving planes and the sliding method*, Bol. Soc. Brasil. Mat. (N.S.), 22, 1991, 1, 1–37.
- [4] H. Berestycki and L. Nirenberg, *Travelling fronts in cylinders*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 9, (1992), 5, 497–572,
- [5] J. Carr and A. Chmaj, *Uniqueness of travelling waves for nonlocal monostable equations*, Proc. Amer. Math. Soc. 132, (2004), no. 8, 2433–2439.
- [6] X. Chen and J.-S. Guo, *Uniqueness and existence of traveling waves for discrete quasilinear monostable dynamics*, Math. Ann. 326, (2003), no. 1, 123–146.
- [7] R. Coutinho and B. Fernandez, *Fronts in extended systems of bistable maps coupled via convolutions*, Nonlinearity 17, (2004), no. 1, 23–47.
- [8] J. Coville, *On uniqueness and monotonicity of solutions of non-local reaction diffusion equation*, Ann. Mat. Pura Appl. (4) 185, 2006, 3, 461–485
- [9] J. Coville, *Maximum principle, sliding techniques and applications to nonlocal equations*, Submitted.
- [10] J. Coville, *Travelling fronts in asymmetric nonlocal reaction diffusion equation: The bistable and ignition case*, Submitted
- [11] J. Coville, J. Dávila and S. Martínez, *Existence and uniqueness of solutions to a non-local equation with monostable nonlinearity*. Submitted.

- [12] J. Coville and L. Dupaigne, *On a nonlocal reaction diffusion equation arising in population dynamics*, Proc. Roy. Soc. Edinburgh Sect. A, To appear.
 - [13] A. De Masi, T. Gobron and E. Presutti, *Travelling fronts in non-local evolution equations*, Arch. Rational Mech. Anal. 132, (1995), no. 2, 143–205.
 - [14] G. Ermentrout and J. McLeod, *Existence and uniqueness of travelling waves for a neural network*, Proc. Roy. Soc. Edinburgh Sect. A 123, (1993), no. 3, 461–478.
 - [15] P. Fife, *Mathematical aspects of reacting and diffusing systems*, Lecture Notes in Biomathematics 28, Springer-Verlag, Berlin, 1979.
 - [16] P. Fife, *An integrodifferential analog of semilinear parabolic PDEs*, Partial differential equations and applications, Lecture Notes in Pure and Appl. Math. 177, 137–145, Dekker, New York, 1996.
 - [17] N. Fournier and S. Méléard, *A microscopic probabilistic description of a locally regulated*, Ann. Appl. Probab. 14, (2004), no. 4, 1880–1919.
 - [18] Gilding, Brian H. and Kersner, Robert, *Travelling waves in nonlinear diffusion-convection reaction*, Progress in Nonlinear Differential Equations and their Applications, 60, Birkhäuser Verlag, Basel, 2004.
 - [19] G. Harris, W. Hudson and B. Zinner, *Traveling wavefronts for the discrete Fisher's equation*, J. Differential Equations 105, (1993), no. 1, 46–62.
 - [20] V. Hutson, S. Martínez, K. Mischaikow and G. Vickers, *The evolution of dispersal*, J. Math. Biol. 47, (2003), no. 6, 483–517.
 - [21] Kot, Mark and Medlock, Jan, *Spreading disease: integro-differential equations old and new*, Math. Biosci. 184, 2003, 2, 201–222.
 - [22] Lucia, Marcello and Muratov, Cyrill B. and Novaga, Matteo, *Linear vs. nonlinear selection for the propagation speed of the solutions of scalar reaction-diffusion equations invading an unstable equilibrium*, Comm. Pure Appl. Math., 57, 2004, 5, 616–636.
 - [23] Mollison, J. D. *The rate of spatial propagation of simple epidemics*, Proc. 6th Berkeley Symp. on Math. Statist. and Prob. Vol 3, University of California press, Berkeley, 1972 579.
 - [24] Murray, J. D. *Mathematical biology*, Biomathematics 19, Second Edition, Springer-Verlag, Berlin, 1993.
 - [25] K. Schumacher, *Travelling-front solutions for integro-differential equations. I*, J. Reine Angew. Math. 316, (1980), 54–70.
 - [26] van Saarloos, Wim, *Front propagation into unstable states: marginal stability as a dynamical mechanism for velocity selection*, Phys. Rev. A (3), 37, 1988, 1, 211–229.
 - [27] van Saarloos, Wim, *Front propagation into unstable states. II. Linear versus nonlinear marginal stability and rate of convergence*, Phys. Rev. A (3), 39, 1989, 12, 6367–6390.
 - [28] H. F. Weinberger, *Long-time behavior of a class of biological models*, SIAM J. Math. Anal., 13, 1982, 3, 353–396.
 - [29] H. F. Weinberger, *Asymptotic behavior of a model in population genetics*, 47–96. Lecture Notes in Math., Vol. 648, 1978,
 - [30] D. V. Widder, *The Laplace transform*, Princeton University Press, Princeton, NJ 1941.
- E-mail address: coville@mis.mpg.de
E-mail address: jdavila@dim.uchile.cl
E-mail address: samartin@dim.uchile.cl

¹MAX PLANCK INSTITUT FOR MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, D-04103 LEIPZIG, DEUTSCHLAND

² DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CENTRO DE MODELAMIENTO MATEMÁTICO, UMI 2807 CNRS, UNIVERSIDAD DE CHILE, BLANCO ENCALADA 2120 - 5 PISO, SANTIAGO - CHILE