

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

Multiple valued functions and integral currents

by

Camillo De Lellis and Emanuele Spadaro

Preprint no.: 54

2013



MULTIPLE VALUED FUNCTIONS AND INTEGRAL CURRENTS

CAMILLO DE LELLIS AND EMANUELE SPADARO

ABSTRACT. We prove several results on Almgren's multiple valued functions and their links to integral currents. In particular, we give a simple proof of the fact that a Lipschitz multiple valued map naturally defines an integer rectifiable current; we derive explicit formulae for the boundary, the mass and the first variations along certain specific vector-fields; and exploit this connection to derive a delicate reparametrization property for multiple valued functions. These results play a crucial role in our new proof of the partial regularity of area minimizing currents [5, 6, 7].

0. INTRODUCTION

It is known since the pioneering work of Federer and Fleming [10] that one can naturally associate an integer rectifiable current to the graph of a Lipschitz function in the Euclidean space, integrating forms over the corresponding submanifold, endowed with its natural orientation. It is then possible to derive formulae for the boundary of the current, its mass and its first variations along smooth vector-fields. Moreover, all these formulae have important Taylor expansions when the current is sufficiently flat. In this paper we provide elementary proofs for the corresponding facts in the case of Almgren's multiple valued functions (see [4] for the relevant definitions).

The connection between multiple valued functions and integral currents is crucial in the analysis of the regularity of area minimizing currents for two reasons. On the one hand, it provides the necessary tools for the approximation of currents with graphs of multiple valued function. This is a fundamental idea for the study of the regularity of minimizing currents in the classical "single-valued" case, and it also plays a fundamental role in the proof of Almgren's partial regularity result (cf. [1, 5]). In this perspective, explicit expressions for the mass and the first variations are necessary to derive the right estimates on the main geometric quantities involved in the regularity theory (cf. [5, 6, 7]). On the other hand, the connection can be exploited to infer interesting conclusions about the multiple valued functions themselves.

This point of view has been taken fruitfully in many problems for the case of classical functions (see, for instance, [11, 12] and the references therein), and has been recently exploited in the multiple valued setting in [3, 14]. The prototypical example of interest here is the following: let $f : \mathbb{R}^m \supset \Omega \rightarrow \mathbb{R}^n$ be a Lipschitz map and $\text{Gr}(f)$ its graph. If the Lipschitz constant of f is small and we change coordinates in \mathbb{R}^{m+n} with an orthogonal transformation close to the identity, then the set $\text{Gr}(f)$ is the graph of a Lipschitz function \tilde{f} over some domain $\tilde{\Omega}$ also in the new system of coordinates. In fact it is easy to see that there exist suitable maps Ψ and Φ such that $\tilde{f}(x) = \Psi(x, f(\Phi(x)))$. In the multiple valued

case, it remains still true that $\text{Gr}(f)$ is the graph of a new Lipschitz map \tilde{f} in the new system of coordinates, but we are not aware of any elementary proof of such statement, which has to be much more subtle because simple relations as the one above cannot hold. It turns out that the structure of $\text{Gr}(f)$ as integral current gives a simple approach to this and similar issues. Several natural estimates can then be proved for \tilde{f} , although more involved and much harder. The last section of the paper is dedicated to these questions; more careful estimates obtained in the same vein will also be given in [6], where they play a crucial role.

Most of the conclusions of this paper are already established, or have a counterpart, in Almgren's monograph [1], but we are not always able to point out precise references to statements therein. However, also when this is possible, our proofs have an independent interest and are in our opinion much simpler. More precisely, the material of Sections 1 and 2 is covered by [1, Sections 1.5-1.7], where Almgren deals with general flat chains. This is more than what is needed in [5, 6, 7], and for this reason we have chosen to treat only the case of integer rectifiable currents. Our approach is anyway simpler and, instead of relying, as Almgren does, on the intersection theory of flat chains, we use rather elementary tools. For the theorems of Section 3 we cannot point out precise references, but Taylor expansions for the area functional are ubiquitous in [1, Chapters 3 and 4]. The theorems of Section 4 do not appear in [1], as Almgren seems to consider only some particular classes of deformations (the "squeeze" and "squash", see [1, Chapter 5]), while we derive fairly general formulas. Finally, it is very likely that the conclusions of Section 5 appear in some form in the construction of the center manifold of [1, Chapter 4], but we cannot follow the intricate arguments and notation of that chapter. In any case, our approach to "reparametrizations" of multiple valued maps seems more flexible and powerful, capable of further applications, because, as it was first realized in [4], we can use tools from metric analysis and metric geometry developed in the last 20 years.

Acknowledgments The research of Camillo De Lellis has been supported by the ERC grant agreement RAM (Regularity for Area Minimizing currents), ERC 306247. The authors are warmly thankful to Bill Allard for several enlightening conversations and his constant enthusiastic encouragement; and very grateful to Luca Spolaor and Matteo Focardi for carefully reading a preliminary version of the paper and for their very useful comments. Camillo De Lellis is also very thankful to the University of Princeton, where he has spent most of his sabbatical completing this and the papers [5, 6, 7].

1. Q -VALUED PUSH-FORWARDS

We use the notation $\langle \cdot, \cdot \rangle$ for: the euclidean scalar product, the naturally induced inner products on p -vectors and p -covectors and the duality pairing of p -vectors and p -covectors; we instead restrict the use of the symbol \cdot to matrix products. Given a C^1 m -dimensional submanifold $\Sigma \subset \mathbb{R}^N$, a function $f : \Sigma \rightarrow \mathbb{R}^k$ and a vector field X tangent to Σ , we denote by $D_X f$ the derivative of f along X , that is $D_X f(p) = (f \circ \gamma)'(0)$ whenever γ is a smooth curve on Σ with $\gamma(0) = p$ and $\gamma'(0) = X(p)$. When $k = 1$, we denote by ∇f the vector field tangent to Σ such that $\langle \nabla f, X \rangle = D_X f$ for every tangent vector field X . For general

k , $Df|_x : T_x\Sigma \rightarrow \mathbb{R}^k$ will be the linear operator such that $Df|_x \cdot X(x) = D_X f(x)$ for any tangent vector field X . We write Df for the map $x \mapsto Df|_x$ and sometimes we will also use the notation $Df(x)$ in place of $Df|_x$. Having fixed an orthonormal base e_1, \dots, e_m on $T_x\Sigma$ and letting (f_1, \dots, f_k) be the components of f , we can write $\nabla f_i = \sum_{j=1}^m a_{ij} e_j$ and $|Df|$ for the usual Hilbert-Schmidt norm:

$$|Df|^2 = \sum_{j=1}^m |D_{e_j} f|^2 = \sum_{i=1}^k |\nabla f_i|^2 = \sum_{i,j} a_{ij}^2.$$

All the notation above is extended to the differential of Lipschitz multiple valued functions at points where they are differentiable in the sense of [4, Definition 1.19]: although the definition in there is for euclidean domains, its extension to C^1 submanifolds $\Sigma \subset \mathbb{R}^N$ is done, as usual, using coordinate charts.

We will keep the same notation also when $f = Y$ is a vector field, i.e. takes values in \mathbb{R}^N , the same Euclidean space where Σ is embedded. In that case we define additionally $\text{div}_\Sigma Y := \sum_i \langle D_{e_i} Y, e_i \rangle$. Moreover, when Y is tangent to Σ , we introduce the covariant derivative $D_\Sigma Y|_x$, i.e. a linear map from $T_x\Sigma$ into itself which gives the tangential component of $D_X Y$. Thus, if we denote by $\mathbf{p}_x : \mathbb{R}^N \rightarrow T_x\Sigma$ the orthogonal projection onto $T_x\Sigma$, we have $D_\Sigma Y|_x = \mathbf{p}_x \cdot DY(x)$. It follows that $D_\Sigma Y \cdot X = \nabla_X Y$, where we use ∇ for the connection (or covariant differentiation) on Σ compatible with its structure as Riemannian submanifold of \mathbb{R}^N . Such covariant differentiation is then extended in the usual way to general tensors on Σ .

When dealing with C^2 submanifolds Σ of \mathbb{R}^N we will denote by A the following tensor: $A|_x$ as a bilinear map on $T_x\Sigma \times T_x\Sigma$ taking values on $T_x\Sigma^\perp$ (the orthogonal complement of $T_x\Sigma$) and if X and Y are vector fields tangent to Σ , then $A(X, Y)$ is the normal component of $D_X Y$, which we will denote by $D_X^\perp Y$. A is called second fundamental form by some authors (cf. [13, Section 7], where the tensor is denoted by B) and we will use the same terminology, although in differential geometry it is more customary to call A “shape operator” and to use “second fundamental form” for scalar products $\langle A(X, Y), \eta \rangle$ with a fixed normal vector field (cf. [8, Chapter 6, Section 2] and [15, Vol. 3, Chapter 1]). In addition, H will denote the trace of A (i.e. $H = \sum_i A(e_i, e_i)$ where e_1, \dots, e_m is an orthonormal frame tangent to Σ) and will be called *mean curvature*.

1.1. Push-forward through multiple valued functions of C^1 submanifolds. In what follows we consider an m -dimensional C^1 submanifold Σ of \mathbb{R}^N and use the word *measurable* for those subsets of M which are \mathcal{H}^m -measurable. Any time we write an integral over (a measurable subset of) Σ we understand that this integral is taken with respect to the \mathcal{H}^m measure. We recall the following lemma which, even if not stated explicitly in [4], is contained in several arguments therein.

Lemma 1.1 (Decomposition). *Let $M \subset \Sigma$ be measurable and $F : M \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ Lipschitz. Then there are a countable partition of M in bounded measurable subsets M_i ($i \in \mathbb{N}$) and Lipschitz functions $f_i^j : M_i \rightarrow \mathbb{R}^n$ ($j \in \{1, \dots, Q\}$) such that*

- (a) $F|_{M_i} = \sum_{j=1}^Q \llbracket f_i^j \rrbracket$ for every $i \in \mathbb{N}$ and $\text{Lip}(f_i^j) \leq \text{Lip}(F) \forall i, j$;

- (b) $\forall i \in \mathbb{N}$ and $j, j' \in \{1, \dots, Q\}$, either $f_i^j \equiv f_i^{j'}$ or $f_i^j(x) \neq f_i^{j'}(x) \forall x \in M_i$;
(c) $\forall i$ we have $DF(x) = \sum_{j=1}^Q \llbracket Df_i^j(x) \rrbracket$ for a.e. $x \in M_i$.

Proof. The proof is by induction on Q . For $Q = 1$ it is obvious. Assume the statement for any $Q^* < Q$ and fix F and M . Note that, without loss of generality, we can assume that M is bounded. We set $M_0 := \{x : \exists y = y(x) \in \mathbb{R}^n \text{ with } F(x) = Q \llbracket y \rrbracket\}$. Clearly, M_0 is measurable because it is the counterimage of a closed subset of $\mathcal{A}_Q(\mathbb{R}^n)$. Moreover, $y : M_0 \rightarrow \mathbb{R}^n$ is Lipschitz. We then set $f_0^j = y$ for every $j \in \{1, \dots, Q\}$. Next, consider $x \notin M_0$. By [4, Proposition 1.6] there exist a ball B_x , two positive numbers Q_1 and Q_2 , and two Lipschitz Q_l -valued functions $g_l : M \cap B_x \rightarrow \mathcal{A}_{Q_l}(\mathbb{R}^n)$ for $l = 1, 2$ such that $F|_{B_x \cap M} = \llbracket g_1 \rrbracket + \llbracket g_2 \rrbracket$. We can apply the inductive hypothesis to g_1 and g_2 , and conclude that $F|_{B_x \cap M}$ can be reduced to the form as in (a) and (b) when restricted to a (suitably chosen) countable partition of $M \cap B_x$ into measurable sets. Since Σ is paracompact, we can find a countable cover $\{B_{x_i}\}_i$ of $M \setminus M_0$, from which (a) and (b) follow. The last statement can be easily verified at every Lebesgue point $x \in M_i$ where F and all the f_i^j 's are differentiable. \square

When $F : M \subset \Sigma \rightarrow \mathbb{R}^n$ is a proper Lipschitz function and $\Sigma \subset \mathbb{R}^N$ is oriented, the current $S = F_{\#} \llbracket M \rrbracket$ in \mathbb{R}^n is given by

$$S(\omega) = \int_M \langle \omega(F(x)), DF(x)_{\#} \vec{e}(x) \rangle d\mathcal{H}^m(x) \quad \forall \omega \in \mathcal{D}^m(\mathbb{R}^n),$$

where $\vec{e}(x) = e_1(x) \wedge \dots \wedge e_m(x)$ is the orienting m -vector of Σ and

$$DF(x)_{\#} \vec{e} = (DF|_x \cdot e_1) \wedge \dots \wedge (DF|_x \cdot e_m),$$

(cf. [13, Remark 26.21(3)]; as usual $\mathcal{D}^m(\Omega)$ denotes the space of smooth m -forms compactly supported in Ω). Using the Decomposition Lemma 1.1 it is possible to extend this definition to multiple valued functions. To this purpose, we give the definition of *proper* multiple valued functions.

Definition 1.2 (Proper Q -valued maps). A measurable $F : M \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ is called *proper* if there is a measurable selection F^1, \dots, F^Q as in [4, Definition 1.1] (i.e. $F = \sum_i \llbracket F^i \rrbracket$) such that $\bigcup_i \overline{(F^i)^{-1}(K)}$ is compact for every compact $K \subset \mathbb{R}^n$. It is then obvious that if there exists such a selection, then *every* measurable selection shares the same property.

We warn the reader that the terminology might be slightly misleading, as the condition above is effectively *stronger* than the usual properness of maps taking values in the metric space $(\mathcal{A}_Q(\mathbb{R}^n), \mathcal{G})$, even when F is continuous: the standard notion of *properness* would not ensure the well-definition of the multiple-valued push-forward.

Definition 1.3 (Q -valued push-forward). Let $\Sigma \subset \mathbb{R}^N$ be a C^1 oriented manifold, $M \subset \Sigma$ a measurable subset and $F : M \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ a proper Lipschitz map. Then, we define the push-forward of M through F as the current $\mathbf{T}_F = \sum_{i,j} (f_i^j)_{\#} \llbracket M_i \rrbracket$, where M_i and f_i^j are

as in Lemma 1.1: that is,

$$\mathbf{T}_F(\omega) := \sum_{i \in \mathbb{N}} \sum_{j=1}^Q \underbrace{\int_{M_i} \langle \omega(f_i^j(x)), Df_i^j(x) \sharp \vec{e}(x) \rangle d\mathcal{H}^m(x)}_{T_{ij}(\omega)} \quad \forall \omega \in \mathcal{D}^m(\mathbb{R}^n). \quad (1.1)$$

We first want to show that T is well-defined. Since F is proper, we easily deduce that

$$|T_{ij}(\omega)| \leq \text{Lip}(F) \|\omega\|_\infty \mathcal{H}^m((f_i^j)^{-1}(\text{spt}(\omega))) < \infty.$$

On the other hand, upon setting $F^j(x) := f_i^j(x)$ for $x \in M_i$, we have $\cup_i (f_i^j)^{-1}(\text{spt}(\omega)) = (F^j)^{-1}(\text{spt}(\omega))$ and $(f_i^j)^{-1}(\text{spt}(\omega)) \cap (f_{i'}^j)^{-1}(\text{spt}(\omega)) = \emptyset$ for $i \neq i'$, thus leading to

$$\sum_{i,j} |T_{ij}(\omega)| \leq \text{Lip}(F) \|\omega\|_\infty \sum_{j=1}^Q \mathcal{H}^m((F^j)^{-1}(\text{spt}(\omega))) < +\infty.$$

Therefore, we can pass the sum inside the integral in (1.1) and, by Lemma 1.1, get

$$\mathbf{T}_F(\omega) = \int_M \sum_{l=1}^Q \langle \omega(F^l(x)), DF^l(x) \sharp \vec{e}(x) \rangle d\mathcal{H}^m(x) \quad \forall \omega \in \mathcal{D}^m(\mathbb{R}^n). \quad (1.2)$$

In particular, recalling the standard theory of rectifiable currents (cf. [13, Section 27]) and the area formula (cf. [13, Section 8]), we have achieved the following proposition.

Proposition 1.4 (Representation of the push-forward). *The definition of the action of \mathbf{T}_F in (1.1) does not depend on the chosen partition M_i nor on the chosen decomposition $\{f_i^j\}$, (1.2) holds and, hence, \mathbf{T}_F is a (well-defined) integer rectifiable current given by $\mathbf{T}_F = (\text{Im}(F), \Theta, \vec{\tau})$ where:*

- (R1) $\text{Im}(F) = \bigcup_{x \in M} \text{spt}(F(x)) = \bigcup_{i \in \mathbb{N}} \bigcup_{j=1}^Q f_i^j(M_i)$ is an m -dimensional rectifiable set;
- (R2) $\vec{\tau}$ is a Borel unitary m -vector orienting $\text{Im}(F)$; moreover, for \mathcal{H}^m -a.e. $p \in \text{Im}(F)$, we have $Df_i^j(x) \sharp \vec{e}(x) \neq 0$ for every i, j, x with $f_i^j(x) = p$ and

$$\vec{\tau}(p) = \pm \frac{Df_i^j(x) \sharp \vec{e}(x)}{|Df_i^j(x) \sharp \vec{e}(x)|}; \quad (1.3)$$

- (R3) for \mathcal{H}^m -a.e. $p \in \text{Im}(F)$, the (Borel) multiplicity function Θ equals

$$\Theta(p) := \sum_{i,j,x: f_i^j(x)=p} \left\langle \vec{\tau}, \frac{Df_i^j(x) \sharp \vec{e}(x)}{|Df_i^j(x) \sharp \vec{e}(x)|} \right\rangle.$$

1.2. Push-forward of Lipschitz submanifolds. As for the classical push-forward, Definition 1.3 can be extended to domains Σ which are Lipschitz submanifolds using the fact that such Σ can be ‘‘chopped’’ into C^1 pieces. Recall indeed the following fact.

Theorem 1.5 ([13, Theorem 5.3]). *If Σ is a Lipschitz m -dimensional oriented submanifold, then there are countably many C^1 m -dimensional oriented submanifolds Σ_i which cover \mathcal{H}^m -a.s. Σ and such that the orientations of Σ and Σ_i coincide on their intersection.*

Definition 1.6 (Q -valued push-forward of Lipschitz submanifolds). Let $\Sigma \subset \mathbb{R}^N$ be a Lipschitz oriented submanifold, $M \subset \Sigma$ a measurable subset and $F : M \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ a proper Lipschitz map. Consider the $\{\Sigma_i\}$ of Theorem 1.5 and set $F_i := F|_{M \cap \Sigma_i}$. Then, we define the push-forward of M through F as the integer rectifiable current $\mathbf{T}_F := \sum_i \mathbf{T}_{F_i}$.

The aboved definition can be extended to Q -valued pushforwards of general rectifiable currents in a straightforward way: however this will never be used in the papers [5, 6, 7] and thus goes beyond the scope of our work. The following conclusion is a simple consequence of Theorem 1.5 and classical arguments in geometric measure theory (cf. [13, Section 27]).

Lemma 1.7. *Let M, Σ and F be as in Definition 1.6 and consider a Borel unitary m -vector \vec{e} orienting Σ . Then \mathbf{T}_F is a well-defined integer rectifiable current for which all the conclusions of Proposition 1.4 hold.*

As for the classical push-forward, \mathbf{T}_F is invariant under bilipschitz change of variables.

Lemma 1.8 (Bilipschitz invariance). *Let $F : \Sigma \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ be a Lipschitz and proper map, $\Phi : \Sigma' \rightarrow \Sigma$ a bilipschitz homeomorphism and $G := F \circ \Phi$. Then, $\mathbf{T}_F = \mathbf{T}_G$.*

Proof. The lemma follows trivially from the corresponding result for classical push-forwards (see [9, 4.1.7 & 4.1.14]), the Decomposition Lemma 1.1 and the definition of Q -valued push-forward. \square

We will next use the area formula to compute explicitly the mass of \mathbf{T}_F . Following standard notation, we will denote by $\mathbf{J}F^j(x)$ the Jacobian determinant of DF^j , i.e. the number

$$|DF^j(x)_\# \vec{e}| = \sqrt{\det((DF^j(x))^T \cdot DF^j(x))}$$

Lemma 1.9 (Q -valued area formula). *Let Σ, M and $F = \sum_j \llbracket F^j \rrbracket$ be as in Definition 1.6. Then, for any bounded Borel function $h : \mathbb{R}^n \rightarrow [0, \infty[$, we have*

$$\int h(p) d\|\mathbf{T}_F\|(p) \leq \int_M \sum_j h(F^j(x)) \mathbf{J}F^j(x) d\mathcal{H}^m(x). \quad (1.4)$$

Equality holds in (1.4) if there is a set $M' \subset M$ of full measure for which

$$\langle DF^j(x)_\# \vec{e}(x), DF^i(y)_\# \vec{e}(y) \rangle \geq 0 \quad \forall x, y \in M' \text{ and } i, j \text{ with } F^i(x) = F^j(y). \quad (1.5)$$

If (1.5) holds the formula is valid also for bounded real-valued Borel h with compact support.

Proof. Let $h : \mathbb{R}^n \rightarrow [0, \infty[$ be a Borel function. Consider a decomposition as in the Decomposition Lemma 1.1 and the integer rectifiable currents T_{ij} of (1.1). By the classical area formula, see [13, Remark 27.2], we have

$$\int h(p) d\|T_{ij}\|(p) \leq \int_{M_i} h(f_i^j(x)) \mathbf{J}f_i^j(x) d\mathcal{H}^m(x). \quad (1.6)$$

Summing this inequality over i and j and using Lemma 1.1(c), we easily conclude (1.4). When (1.5) holds, we can choose $\vec{\tau}$ of Proposition 1.4 such that the identity (1.3) has always the + sign. Define $\Theta_{ij}(p) := \mathcal{H}^0(\{x : f_i^j(x) = p\})$. We then conclude from Proposition

1.4(R3) that $\Theta(p) = \sum_{i,j} \Theta_{ij}(p)$ for \mathcal{H}^m -a.e. $p \in \text{Im}(F)$. On the other hand, again by [13, Remark 27.2], equality holds in (1.6) and, moreover, we have the identities $\|T_{ij}\| = \Theta_{ij} \mathcal{H}^m \llcorner \text{Im}(f_i^j)$, $\|\mathbf{T}_F\| = \Theta \mathcal{H}^m \llcorner \text{Im}(F)$. This easily implies the second part of the lemma and hence completes the proof. \square

A particular class of push-forwards are given by graphs.

Definition 1.10 (*Q-graphs*). Let Σ, M and $f = \sum_i \llbracket f_i \rrbracket$ be as in Definition 1.6. Define the map $F : M \rightarrow \mathcal{A}_Q(\mathbb{R}^{N+n})$ as $F(x) := \sum_{i=1}^Q \llbracket (x, f_i(x)) \rrbracket$. \mathbf{T}_F is the *current associated to the graph* $\text{Gr}(f)$ and will be denoted by \mathbf{G}_f .

Observe that, if Σ, f and F are as in Definition 1.10, then the condition (1.5) is always trivially satisfied. Moreover, when $\Sigma = \mathbb{R}^m$ the well-known Cauchy-Binet formula gives

$$(\mathbf{J}F^j)^2 = 1 + \sum_{k=1}^m \sum_{A \in M^k(DF^j)} (\det A)^2,$$

where $M^k(B)$ denotes the set of all $k \times k$ minors of the matrix B . Lemma 1.9 gives then the following corollary in the case of *Q-graphs*

Corollary 1.11 (*Area formula for Q-graphs*). *Let $\Sigma = \mathbb{R}^m$, $M \subset \mathbb{R}^m$ and f be as in Definition 1.10. Then, for any bounded compactly supported Borel $h : \mathbb{R}^{m+n} \rightarrow \mathbb{R}$, we have*

$$\int h(p) d\|\mathbf{G}_f\|(p) = \int_M \sum_i h(x, f_i(x)) \left(1 + \sum_{k=1}^m \sum_{A \in M^k(DF^j)} (\det A)^2 \right)^{\frac{1}{2}} dx. \quad (1.7)$$

2. BOUNDARIES

In the classical theory of currents, when Σ is a Lipschitz manifold with Lipschitz boundary and $F : \Sigma \rightarrow \mathbb{R}^N$ is Lipschitz and proper, then $\partial(F_{\#} \llbracket \Sigma \rrbracket) = F_{\#} \llbracket \partial \Sigma \rrbracket$ (see [9, 4.1.14]). This result can be extended to multiple-valued functions.

Theorem 2.1 (*Boundary of the push-forward*). *Let Σ be a Lipschitz submanifold of \mathbb{R}^N with Lipschitz boundary, $F : \Sigma \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ a proper Lipschitz function and $f = F|_{\partial \Sigma}$. Then, $\partial \mathbf{T}_F = \mathbf{T}_f$.*

The main building block is the following small variant of [4, Homotopy Lemma 1.8].

Lemma 2.2. *There is $c(Q, m) > 0$ such that, for every closed cube $C \subset \mathbb{R}^m$ centered at x_0 and every $F \in \text{Lip}(C, \mathcal{A}_Q(\mathbb{R}^n))$, we can find $G \in \text{Lip}(C, \mathcal{A}_Q(\mathbb{R}^n))$ satisfying:*

- (i) $G|_{\partial C} = F|_{\partial C} =: f$, $\text{Lip}(G) \leq c \text{Lip}(F)$ and $\|\mathcal{G}(F, G)\|_{L^\infty} \leq c \text{Lip}(F) \text{diam}(C)$;
- (ii) *there are Lipschitz multi-valued maps G_j and f_j (with $j \in \{1, \dots, J\}$) such that $G = \sum_{j=1}^J \llbracket G_j \rrbracket$, $f = \sum_{j=1}^J \llbracket f_j \rrbracket$ and $\mathbf{G}_{G_j} = \llbracket (x_0, a_j) \rrbracket \ast \mathbf{G}_{f_j}$ for some $a_j \in \mathbb{R}^n$.*

Proof. The proof of (i) is contained in [4, Lemma 1.8]. Concerning (ii), the proof is contained in the inductive argument of [4, Lemma 1.8], it suffices to complement the arguments there with the following fact: if $C = [-1, 1]^m$, $u \in \text{Lip}(\partial C, \mathcal{A}_Q(\mathbb{R}^n))$ and

$G(x) = \sum_i \left[\|x\| u_i \left(\frac{x}{\|x\|} \right) \right]$ is the “cone-like” extension of u to C (where $\|x\| = \sup_i |x_i|$), then $\mathbf{G}_G = \llbracket 0 \rrbracket \times \mathbf{G}_u$. The proof of this claim is a simple consequence of the Decomposition Lemma 1.1 and the very definition of \mathbf{G}_u . Consider, indeed, a countable measurable partition $\cup_i M_i = \partial C$ and Lipschitz functions u_i^j with $\sum_j \llbracket u_i^j \rrbracket = u|_{M_i}$. According to our definitions, $\mathbf{G}_u = \sum_{i,j} (u_i^j)_\# \llbracket M_i \rrbracket =: \sum_{i,j} T_{ij}$. Consider now for each i the set $R_i := \{\lambda x : x \in M_i, \lambda \in]0, 1]\}$ and define $G_i^j(\lambda x) := \lambda u_i^j(x)$ for every $x \in M_i$ and $\lambda \in]0, 1]$. The sets R_i are a measurable decomposition of $C \setminus \{0\}$ and we have $\sum_j \llbracket G_i^j \rrbracket = G|_{R_i}$. Therefore, setting $S_{ij} := (G_i^j)_\# \llbracket R_i \rrbracket$, we have $\mathbf{G}_G = \sum_{i,j} S_{ij}$. On the other hand, by the classical theory of currents $S_{ij} = \llbracket 0 \rrbracket \times T_{ij}$ (see [9, Section 4.1.11]). Since $\sum_{i,j} (\mathbf{M}(S_{ij}) + \mathbf{M}(T_{ij})) < \infty$, the desired claim follows. \square

Proof of Theorem 2.1. The proof is by induction on the dimension m . Since every Lipschitz manifold can be triangulated and the statement is invariant under bilipschitz homeomorphisms, it suffices to prove the theorem when $\Sigma = [0, 1]^m$. Next, given a classical Lipschitz map $\Phi : \mathbb{R}^N \rightarrow \mathbb{R}^k$, let $\Phi \circ F$ be the multiple-valued map $\sum_i \llbracket \Phi(F_i) \rrbracket$ (cf. [4, Section 1.3.1]). If F is a classical Lipschitz map, then $\mathbf{T}_{\Phi \circ F} = \Phi_\# F_\# \llbracket \Sigma \rrbracket = \Phi_\# \mathbf{T}_F$ (cf. [9, 4.1.14]). The same identity holds for Q -valued map, as the Decomposition Lemma 1.1 easily reduces it to the single-valued case. Then, if $\mathbf{p} : \mathbb{R}^m \times \mathbb{R}^{m+n} \rightarrow \mathbb{R}^{m+n}$ is the orthogonal projection on the second components, we have $\mathbf{p}_\# \mathbf{G}_F = \mathbf{T}_F$. Given the classical commutation of boundary and (single-valued) push-forward (see [9, Section 4.1.14]) we are then reduced to proving the identity $\partial \mathbf{G}_F = \mathbf{G}_f$.

We turn therefore to the case \mathbf{G}_F . The starting step $m = 1$ is an obvious corollary of the Lipschitz selection principle [4, Proposition 1.2]. Indeed, for $F \in \text{Lip}([0, 1], \mathcal{A}_Q(\mathbb{R}^n))$, there exist functions $F_i \in \text{Lip}([0, 1], \mathbb{R}^n)$ such that $F = \sum_i \llbracket F_i \rrbracket$. Therefore, $\mathbf{T}_F = \sum_i \mathbf{T}_{F_i}$ and

$$\partial \mathbf{T}_F = \sum_i \partial \mathbf{T}_{F_i} = \sum_i (\llbracket F_i(1) \rrbracket - \llbracket F_i(0) \rrbracket) = \mathbf{T}_f.$$

For the inductive argument, consider the dyadic decomposition at scale 2^{-l} of $[0, 1]^m$:

$$[0, 1]^m = \bigcup_{k \in \{0, \dots, 2^l - 1\}^m} Q_{k,l}, \quad \text{with } Q_{k,l} = 2^{-l}(k + [0, 1]^m).$$

In each $Q_{k,l}$, let $u_{k,l}$ be the cone-like extension given by Lemma 2.2 of $f_{k,l} := F|_{\partial Q_{k,l}}$. Denote by u_l the Q -function on $[0, 1]^m$ which coincides with $u_{k,l}$ on each $Q_{k,l}$. Obviously the u_l 's are equi-Lipschitz and converge uniformly to F by Lemma 2.2 (i). Set $T_l := \mathbf{G}_{u_l} = \sum_k \mathbf{G}_{u_{k,l}}$. By the inductive hypothesis $\partial \mathbf{G}_{f_{k,l}} = 0$. Since $\partial(\llbracket p \rrbracket \times T) = T - \llbracket p \rrbracket \times \partial T$ (see [13, Section 26]), Lemma 2.2 implies $\partial \mathbf{G}_{u_{k,l}} = \mathbf{G}_{f_{k,l}}$. Considering that the boundary faces common to adjacent cubes come with opposite orientations, we conclude $\partial T_l = \mathbf{G}_f$. By Corollary 1.11, $\limsup_l (\mathbf{M}(T_l) + \mathbf{M}(\partial T_l)) < \infty$ and so the compactness theorem for integral currents (see [13, Theorem 27.3]) guarantees the existence of an integral current T which is the weak limit of a subsequence of $\{T_l\}$ (not relabeled). It suffices therefore to show that:

- (C) if $\Omega \subset \mathbb{R}^m$ is an open set and u_l is a sequence of Lipschitz Q -valued maps on Ω such that u_l converge uniformly to some F and $T_l := \mathbf{G}_{u_l}$ converge to an integral current T , then $T = \mathbf{G}_F$.

We will prove (C) by induction over Q : the case $Q = 1$ is classical (see for instance [11, Theorem 2, Section 3.1 in Chapter 3] and [11, Proposition 2, Section 2.1 in Chapter 3]). We assume (C) holds for every $Q^* < Q$ and want to prove it for Q . Fix a sequence as in (C). Clearly T is supported in the rectifiable set $\text{Gr}(F)$. Fix an orthonormal basis e_1, \dots, e_m of \mathbb{R}^m and extend it to an orthonormal basis of \mathbb{R}^{m+n} with positive orientation. Set $\vec{e} = e_1 \wedge \dots \wedge e_m$. Thanks to the Lipschitz regularity of F , $\text{Gr}(F)$ can be oriented by m -planes $\vec{\tau}$ with the property that $\langle \vec{\tau}, \vec{e} \rangle \geq c > 0$, where the constant c depends on $\text{Lip}(F)$. We have $T = (\text{Gr}(F), \vec{\tau}, \bar{\Theta})$ and $\mathbf{G}_F = (\text{Gr}(F), \vec{\tau}, \Theta)$: we just need to show that $\Theta = \bar{\Theta}$ \mathcal{H}^m -a.e. on $\text{Gr}(F)$.

As observed in Lemma 1.1 there is a closed set M_0 and a Lipschitz function f_0 such that:

- $F(x) = Q \llbracket f_0(x) \rrbracket$ for every $x \in M_0$;
- F “splits” locally on $\Omega' = \Omega \setminus M_0$ into (Lipschitz) functions taking less than Q values.

Using the induction hypothesis, it is trivial to verify that $T \llcorner \Omega' \times \mathbb{R}^n = \mathbf{G}_F \llcorner \Omega' \times \mathbb{R}^n$. Thus we just need to show that $\bar{\Theta}(x, f_0(x)) = \Theta(x, f_0(x))$ for \mathcal{H}^m -a.e. $x \in M_0$. Consider the orthogonal projection $\mathbf{p} : \mathbb{R}^{m+n} \rightarrow \mathbb{R}^m$. By the well-known formula for the pushforward of currents (see [9, Lemma 4.1.25]), we have $\mathbf{p}_\# T = \bar{\Theta}' \llbracket \Omega \rrbracket$ and $\mathbf{p}_\# \mathbf{G}_F = \Theta' \llbracket \Omega \rrbracket$, where

$$\bar{\Theta}'(x) = \sum_{(x,y) \in \text{Gr}(F)} \bar{\Theta}(x, y) \quad \text{and} \quad \Theta'(x) = \sum_{(x,y) \in \text{Gr}(F)} \Theta(x, y).$$

Therefore $\bar{\Theta}'(x) = \bar{\Theta}(x, f_0(x))$ and $\Theta'(x) = \Theta(x, f_0(x))$ for \mathcal{H}^m -a.e. $x \in M_0$. On the other hand, by the definition of \mathbf{G}_F and the very same formula for the push-forward (i.e. [9, Lemma 4.1.25]) it is easy to see that $\mathbf{p}_\# \mathbf{G}_F = Q \llbracket \Omega \rrbracket = \mathbf{p}_\# T_l$. Since $\mathbf{p}_\# T_l$ converges to $\mathbf{p}_\# T$, we conclude that $\Theta' \equiv Q \equiv \bar{\Theta}'$ \mathcal{H}^m -a.e. on Ω , which in turn implies $\Theta(x, f_0(x)) = \bar{\Theta}(x, f_0(x))$ for a.e. $x \in M_0$. This completes the proof of the inductive step. \square

3. TAYLOR EXPANSION OF THE AREA FUNCTIONAL

In this section we compute the Taylor expansion of the area functional in several forms. To this aim, we fix the following notation and hypotheses.

Assumption 3.1. We consider the following:

- (M) an open submanifold $\mathcal{M} \subset \mathbb{R}^{m+n}$ of dimension m with $\mathcal{H}^m(\mathcal{M}) < \infty$, which is the graph of a function $\varphi : \mathbb{R}^m \supset \Omega \rightarrow \mathbb{R}^n$ with $\|\varphi\|_{C^3} \leq \bar{c}$; A and H will denote, respectively, the second fundamental form and the mean curvature of \mathcal{M} ;
- (U) a regular tubular neighborhood \mathbf{U} of \mathcal{M} , i.e. the set of points $\{x + y : x \in \mathcal{M}, y \perp T_x \mathcal{M}, |y| < c_0\}$, where the thickness c_0 is sufficiently small so that the nearest point projection $\mathbf{p} : \mathbf{U} \rightarrow \mathcal{M}$ is well defined and C^2 ; the thickness is supposed to be larger than a fixed geometric constant;

(N) a Q -valued map $F : \mathcal{M} \rightarrow \mathcal{A}_Q(\mathbb{R}^{m+n})$ of the form

$$\sum_{i=1}^Q \llbracket F_i(x) \rrbracket = \sum_{i=1}^Q \llbracket x + N_i(x) \rrbracket,$$

where $N : \mathcal{M} \rightarrow \mathcal{A}_Q(\mathbb{R}^{m+n})$ satisfies $x + N_i(x) \in \mathbf{U}$, $N_i(x) \perp T_x \mathcal{M}$ for every x and $\text{Lip}(N) \leq \bar{c}$.

We recall the notation $\boldsymbol{\eta} \circ F := \frac{1}{Q} \sum_i F_i$, for every multiple valued function $F = \sum_i \llbracket F_i \rrbracket$.

Theorem 3.2 (Expansion of $\mathbf{M}(\mathbf{T}_F)$). *If \mathcal{M} , F and N are as in Assumption 3.1 and \bar{c} is smaller than a geometric constant, then*

$$\begin{aligned} \mathbf{M}(\mathbf{T}_F) &= Q \mathcal{H}^m(\mathcal{M}) - Q \int_{\mathcal{M}} \langle H, \boldsymbol{\eta} \circ N \rangle + \frac{1}{2} \int_{\mathcal{M}} |DN|^2 \\ &\quad + \int_{\mathcal{M}} \sum_i \left(P_2(x, N_i) + P_3(x, N_i, DN_i) + R_4(x, DN_i) \right), \end{aligned} \quad (3.1)$$

where P_2, P_3 and R_4 are C^1 functions with the following properties:

(i) $n \mapsto P_2(x, n)$ is a quadratic form on the normal bundle of \mathcal{M} satisfying

$$|P_2(x, n)| \leq C |A(x)|^2 |n|^2 \quad \forall x \in \mathcal{M}, \forall n \perp T_x \mathcal{M}; \quad (3.2)$$

(ii) $P_3(x, n, D) = \sum_i L_i(x, n) Q_i(x, D)$, where $n \mapsto L_i(x, n)$ are linear forms on the normal bundle of \mathcal{M} and $D \mapsto Q_i(x, D)$ are quadratic forms on the space of $(m+n) \times (m+n)$ -matrices, satisfying

$$|L_i(x, n)| |Q_i(x, D)| \leq C |A(x)| |n| |D|^2 \quad \forall x \in \mathcal{M}, \forall n \perp T_x \mathcal{M}, \forall D;$$

(iii) $|R_4(x, D)| = |D|^3 L(x, D)$, for some function L with $\text{Lip}(L) \leq C$, which satisfies $L(x, 0) = 0$ for every $x \in \mathcal{M}$ and is independent of x when $A \equiv 0$.

Moreover, for any Borel function $h : \mathbb{R}^{m+n} \rightarrow \mathbb{R}$,

$$\left| \int h d\|\mathbf{T}_F\| - \int_{\mathcal{M}} \sum_i h \circ F_i \right| \leq C \int_{\mathcal{M}} \left(\sum_i |A| |h \circ F_i| |N_i| + \|h\|_{\infty} (|DN|^2 + |A| |N|^2) \right), \quad (3.3)$$

and, if $h(p) = g(\mathbf{p}(p))$ for some g , we have

$$\left| \int h d\|\mathbf{T}_F\| - Q \int_{\mathcal{M}} (1 - \langle H, \boldsymbol{\eta} \circ N \rangle + \frac{1}{2} |DN|^2) g \right| \leq C \int_{\mathcal{M}} (|A|^2 |N|^2 + |DN|^4) |g|. \quad (3.4)$$

In particular, as a simple corollary of the theorem above, we have the following.

Corollary 3.3 (Expansion of $\mathbf{M}(\mathbf{G}_f)$). *Assume $\Omega \subset \mathbb{R}^m$ is an open set with bounded measure and $f : \Omega \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ a Lipschitz map with $\text{Lip}(f) \leq \bar{c}$. Then,*

$$\mathbf{M}(\mathbf{G}_f) = Q |\Omega| + \frac{1}{2} \int_{\Omega} |Df|^2 + \int_{\Omega} \sum_i \bar{R}_4(Df_i), \quad (3.5)$$

where $\bar{R}_4 \in C^1$ satisfies $|\bar{R}_4(D)| = |D|^3 \bar{L}(D)$ for \bar{L} with $\text{Lip}(\bar{L}) \leq C$ and $\bar{L}(0) = 0$.

Proof. The corollary is reduced to Theorem 3.2 by simply setting $\mathcal{M} = \Omega \times \{0\}$,

$$N = \sum_i \llbracket N_i(x) \rrbracket := \sum_i \llbracket (0, f_i(x)) \rrbracket \quad \text{and} \quad F(x) = \sum_i \llbracket F_i(x) \rrbracket = \sum_i \llbracket (x, f_i(x)) \rrbracket .$$

Since in this case A vanishes, (3.1) gives precisely (3.5). \square

Proof of Theorem 3.2. We will in fact prove the statement for $\mathbf{M}(\mathbf{T}_{F|_V})$, where V is any Borel subset of \mathcal{M} . Under this generality, by the decomposition Lemma 1.1, it is enough to consider the case $F|_V = \sum_i G_i$, where each $G_i = F_i|_V = x + N_i|_V$ is a (one-valued!) Lipschitz map. Next observe that (1.5) obviously holds if \bar{c} is sufficiently small. Therefore,

$$\mathbf{M}(\mathbf{T}_{F|_V}) = \sum_i \mathbf{M}((F_i)_\# \llbracket V \rrbracket) ,$$

and, since $\boldsymbol{\eta} \circ N = \frac{1}{Q} \sum_i N_i$, the formula (3.1) follows from summing the corresponding identities

$$\begin{aligned} \mathbf{M}((F_i)_\# \llbracket V \rrbracket) &= \mathcal{H}^m(V) + \int_V \langle H, N_i \rangle + \frac{1}{2} \int_V |DN_i|^2 \\ &\quad + \int_V \left(P_2(x, N_i) + P_3(x, N_i, DN_i) + R_4(x, DN_i) \right) . \end{aligned} \quad (3.6)$$

To simplify the notation we drop the subscript i in the proof of (3.6). Using the area formula, we have that

$$\mathbf{M}(F_\# \llbracket V \rrbracket) = \int_V |DF_\# \vec{\xi}| d\mathcal{H}^m ,$$

where $\vec{\xi} = \xi_1 \wedge \dots \wedge \xi_m$ is the simple m -vector associated to an orthonormal frame on $T\mathcal{M}$. By simple multilinear algebra $|DF_\# \vec{\xi}| = \sqrt{\det M}$, where M is the $m \times m$ matrix given by

$$\begin{aligned} M_{jk} &= \langle DF \cdot \xi_j, DF \cdot \xi_k \rangle = \langle \xi_j + DN \cdot \xi_j, \xi_k + DN \cdot \xi_k \rangle \\ &= \delta_{jk} + \underbrace{\langle DN \cdot \xi_j, \xi_k \rangle + \langle DN \cdot \xi_k, \xi_j \rangle}_{a_{jk}} + \underbrace{\langle DN \cdot \xi_j, DN \cdot \xi_k \rangle}_{b_{jk}} . \end{aligned} \quad (3.7)$$

Set $a = (a_{jk})$, $b = (b_{jk})$ and denote by $M_2(a+b)$ and $M_3(a+b)$, respectively, the sum of all 2×2 and that of all 3×3 minors of the matrix $(a+b)$; similarly denote by $R(a+b)$ the sum of all $k \times k$ minors with $k \geq 4$. Then,

$$\det M = 1 + \text{tr}(a+b) + M_2(a+b) + M_3(a+b) + R(a+b) . \quad (3.8)$$

Observe that the entries of a are linear in DN and those of b are quadratic. Thus,

$$M_2(a+b) = M_2(a) + M_2(b) + C_2(a, b) , \quad (3.9)$$

$$M_3(a+b) = M_3(a) + C_4(a, b) , \quad (3.10)$$

where $C_2(a, b)$ is a linear combination of terms of the form $a_{jk} b_{lm}$ and $C_4(a, b)$ is a polynomial in the entries of DN satisfying the inequality $|C_4(a, b)| \leq C|DN|^4$. Recall the

Taylor expansion $\sqrt{1+\tau} = 1 + \frac{\tau}{2} - \frac{\tau^2}{8} + \frac{\tau^3}{16} + g(\tau)$, where g is an analytic function with $|g(\tau)| \leq |\tau|^4$. With the aid of (3.8), (3.9) and (3.10) we reach the following conclusion:

$$|DF_{\sharp}\vec{\xi}| = 1 + \frac{\operatorname{tr}(a+b) + M_2(a) + C_2(a,b) + M_3(a)}{2} - \frac{(\operatorname{tr} a)^2 + 2 \operatorname{tr} a \operatorname{tr} b + 2 \operatorname{tr} a M_2(a)}{8} + \frac{(\operatorname{tr} a)^3}{16} + R_4, \quad (3.11)$$

where R_4 is an analytic function of the entries of DN which satisfies $|R_4(DN)| \leq C|DN|^4$. Observe next that $\operatorname{tr} b = \sum_k \langle DN \cdot \xi_k, DN \cdot \xi_k \rangle = |DN|^2$. Moreover,

$$\langle DN \cdot \xi_j, \xi_k \rangle = \nabla_{\xi_j}(\langle N, \xi_k \rangle) - \langle N, \nabla_{\xi_j} \xi_k \rangle = -\langle N, A(\xi_j, \xi_k) \rangle.$$

Thus, by the symmetry of the second fundamental form, we have

$$a_{jk} = -2\langle A(\xi_j, \xi_k), N \rangle \quad \text{and} \quad \operatorname{tr} a = -2\langle H, N \rangle.$$

We then can rewrite

$$|DF_{\sharp}\vec{\xi}| = 1 - \langle H, N \rangle + \frac{|DN|^2}{2} + \underbrace{\frac{M_2(a)}{2} - \frac{(\operatorname{tr} a)^2}{8}}_{P_2} + \underbrace{\frac{C_2(a,b) + M_3(a)}{2} - \frac{\operatorname{tr} a (\operatorname{tr} b + M_2(a))}{4}}_{P_3} + \frac{(\operatorname{tr} a)^3}{16} + R_4. \quad (3.12)$$

Integrating (3.12) we reach (3.6). It remains to show that P_2 , P_3 and R_4 satisfy (i), (ii) and (iii). If $A = 0$, then \mathcal{M} is flat and the frame ξ_1, \dots, ξ_m can be chosen constant, so that R_4 will not depend on x . Next, each b_{jk} is a quadratic polynomial in the entries of DN , with coefficients which are C^2 functions of x . Instead each a_{jk} can be seen as a linear function in DN with coefficients which are C^2 functions of x , but also as a linear function L_{jk} of N , with a C^1 dependence on x . In the latter case we have the bound $|L_{jk}(x, n)| \leq |A(x)||n|$. Therefore the claims in (i) and (ii) follow easily. Finally, since R_4 is an analytic function of the entries of DN satisfying $|R_4(DN)| \leq C|DN|^4$, the representation in (iii) follows from the elementary consideration that $\frac{R_4(D)}{|D|^3}$ is a Lipschitz function vanishing at the origin.

Finally, observe that the argument above implies (3.4) when g is the indicator function of any measurable set and the general case follows from standard measure theory. The identity (3.3) follows easily from the same formulas for $|DF_{\sharp}\vec{\xi}|$, using indeed cruder estimates. \square

3.1. Taylor expansion for the excess in a cylinder. The last results of this section concerns estimates of the excess in different systems of coordinates, in particular with respect to tilted planes and curvilinear coordinates.

Proposition 3.4 (Expansion of a curvilinear excess). *There exist a dimensional constant $C > 0$ such that, if \mathcal{M} , F and N are as in Assumption 3.1 with \bar{c} small enough, then*

$$\left| \int |\vec{\mathbf{T}}_F(x) - \vec{\mathcal{M}}(\mathbf{p}(x))|^2 d\|\mathbf{T}_F\|(x) - \int_{\mathcal{M}} |DN|^2 \right| \leq C \int_{\mathcal{M}} (|A|^2|N|^2 + |DN|^4), \quad (3.13)$$

where $\vec{\mathbf{T}}_F$ and $\vec{\mathcal{M}}$ are the unit m -vectors orienting \mathbf{T}_F and $T\mathcal{M}$, respectively.

Proof. Let $p \in \mathcal{M}$ and define $\vec{\mathcal{M}}(p) = \xi_1 \wedge \dots \wedge \xi_m$ for some orthonormal frame ξ_1, \dots, ξ_m for $T\mathcal{M}$ and

$$\vec{\mathbf{T}}_F(F_i(p)) = \frac{\vec{\zeta}_i}{|\vec{\zeta}_i|} \quad \text{with} \quad \vec{\zeta}_i = (\xi_1 + DN_i|_p \cdot \xi_1) \wedge \dots \wedge (\xi_m + DN_i|_p \cdot \xi_m).$$

Our assumptions imply $\mathbf{p}(F_i(p)) = p$. Using the Q -valued area formula and obvious computations we get

$$\int |\vec{\mathbf{T}}_F - \vec{\mathcal{M}} \circ \mathbf{p}|^2 d\|\mathbf{T}_F\|(x) = \int_{\mathcal{M}} \sum_i \left| \frac{\vec{\zeta}_i}{|\vec{\zeta}_i|} - \vec{\mathcal{M}} \right|^2 |\zeta_i| = \int_{\mathcal{M}} 2 \left(\sum_i |\zeta_i| - \sum_i \langle \zeta_i, \vec{\mathcal{M}} \rangle \right).$$

As already computed in the proof of Theorem 3.2,

$$\sum_i |\zeta_i| = Q - Q \langle H, \boldsymbol{\eta} \circ N \rangle + \frac{|DN|^2}{2} + O(|A|^2|N|^2 + |DN|^4).$$

If we next define $B_{jk}^i := \langle \xi_j, \xi_k + DN_i \cdot \xi_k \rangle = \delta_{jk} - \langle N_i, A(\xi_j, \xi_k) \rangle$, we then get

$$\sum_i \langle \zeta_i, \vec{\mathcal{M}} \rangle = \sum_i \det B^i = Q - Q \langle H, \boldsymbol{\eta} \circ N \rangle + O(|A|^2|N|^2).$$

Hence the claimed formula follows easily. \square

Next we compute the excess of a Lipschitz graph with respect to a tilted plane.

Theorem 3.5 (Expansion of a cylindrical excess). *There exist dimensional constants $C, c > 0$ with the following property. Let $f : \mathbb{R}^m \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ be a Lipschitz map with $\text{Lip}(f) \leq c$. For any $0 < s$, set $A := \int_{B_s} D(\boldsymbol{\eta} \circ f)$ and denote by $\vec{\tau}$ the oriented unitary m -dimensional simple vector to the graph of the linear map $y \mapsto A \cdot y$. Then, we have*

$$\left| \int_{C_s} \left| \vec{\mathbf{G}}_f - \vec{\tau} \right|^2 d\|\mathbf{G}_f\| - \int_{B_s} \mathcal{G}(Df, Q[[A]])^2 \right| \leq C \int_{B_s} |Df|^4. \quad (3.14)$$

Proof of Theorem 3.5. Arguing as in the previous proofs, thanks to Lemma 1.1, we can write $f = \sum_i \llbracket f_i \rrbracket$ and process local computations (when needed) as if each f_i were Lipschitz. Moreover, we have that

$$\vec{\tau} = \frac{\vec{\xi}}{|\vec{\xi}|} \quad \text{with} \quad \vec{\xi} = (e_1 + A e_1) \wedge \dots \wedge (e_m + A e_m).$$

Here and for the rest of this proof, we identify \mathbb{R}^m and \mathbb{R}^n with the subspaces $\mathbb{R}^m \times \{0\}$ and $\{0\} \times \mathbb{R}^n$ of \mathbb{R}^{m+n} , respectively: this justifies the notation $e_j + A e_j$ for $e_j \in \mathbb{R}^m$ and $A e_j \in \mathbb{R}^n$. Next, we recall that

$$|\xi| = \sqrt{\langle \xi, \xi \rangle} = \sqrt{\det(\delta_{ij} + \langle A e_i, A e_j \rangle)} = 1 + \frac{1}{2}|A|^2 + O(|A|^4).$$

By Corollary 1.11 we also have

$$\begin{aligned} E &:= \int_{\mathbf{C}_s} \left| \vec{\mathbf{G}}_f - \vec{\tau} \right|^2 d\|\mathbf{G}_f\| = 2\mathbf{M}(\mathbf{G}_f) - 2 \int \langle \vec{\mathbf{G}}_f, \vec{\tau} \rangle d\|\mathbf{G}_f\| \\ &= 2Q|B_s| + \int_{B_s} (|Df|^2 + O(|Df|^4)) - 2 \int \sum_i \langle (e_1 + Df_i e_1) \wedge \dots \wedge (e_m + Df_i e_m), \vec{\tau} \rangle. \end{aligned}$$

On the other hand $\langle A e_j, e_k \rangle = 0 = \langle Df_i e_j, e_k \rangle$. Therefore,

$$\begin{aligned} \langle (e_1 + Df_i e_1) \wedge \dots \wedge (e_m + Df_i e_m), \vec{\tau} \rangle &= |\xi|^{-1} \det(\delta_{jk} + \langle Df_i e_j, A e_k \rangle) \\ &= \left(1 + \frac{|A|^2}{2} + O(|A|^4) \right)^{-1} (1 + Df_i : A + O(|Df|^2 |A|^2)). \end{aligned}$$

Recalling that $|A| \leq C s^{-m} \int |Df| \leq C (s^{-m} \int |Df|^4)^{\frac{1}{4}}$, we then conclude

$$\begin{aligned} E &= \int_{B_s} |Df|^2 + Q|B_s| |A|^2 - 2 \int_{B_s} \sum_i Df_i : A + O\left(\int_{B_s} |Df|^4\right) \\ &= \int_{B_s} \sum_i |Df_i - A|^2 + O\left(\int_{B_s} |Df|^4\right) = \int_{B_s} \mathcal{G}(Df, Q[[A]])^2 + O\left(\int_{B_s} |Df|^4\right). \quad \square \end{aligned}$$

4. FIRST VARIATIONS

In this section we compute the first variations of the currents induced by multiple valued maps. These formulae are ultimately the link between the stationarity of area minimizing currents and the partial differential equations satisfied by suitable approximations. We use here the following standard notation: given a current T in \mathbb{R}^N and a vector field $X \in C^1(\mathbb{R}^N, \mathbb{R}^N)$, we denote the first variation of T along X by $\delta T(X) := \frac{d}{dt} \Big|_{t=0} \mathbf{M}(\Phi_{t\#} T)$, where $\Phi :]-\eta, \eta[\times U \rightarrow \mathbb{R}^N$ is any C^1 isotopy of a neighborhood U of $\text{spt}(T)$ with $\Phi(0, x) = x$ for any $x \in U$ and $\frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \Phi_\varepsilon = X$ (in what follows we will often use Φ_ε for the map $x \mapsto \Phi(\varepsilon, x)$). It would be more appropriate to use the notation $\delta T(\Phi)$ (see, for instance, [9, Section 5.1.7]), but since the currents considered in this paper are rectifiable, it is well known that the first variation depends only on X and is given by the formula

$$\delta T(X) = \int \text{div}_{\vec{T}} X d\|T\|, \quad (4.1)$$

where $\text{div}_{\vec{T}} X = \sum_i \langle D_{e_i} X, e_i \rangle$ for any orthonormal frame e_1, \dots, e_m with $e_1 \wedge \dots \wedge e_m = \vec{T}$ (see [9, 5.1.8] and cf. [13, Section 2.9]). We begin with the expansion for the first variation of graphs. In what follows, $A : B$ will denote the usual Hilbert Schmidt scalar product of two $k \times j$ matrices.

Theorem 4.1 (Expansion of $\delta\mathbf{G}_f(X)$). *Let $\Omega \subset \mathbb{R}^m$ be a bounded open set and $f : \Omega \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ a map with $\text{Lip}(f) \leq \bar{c}$. Consider a function $\zeta \in C^1(\Omega \times \mathbb{R}^m, \mathbb{R}^n)$ and the corresponding vector field $\chi \in C^1(\Omega \times \mathbb{R}^n, \mathbb{R}^{m+n})$ given by $\chi(x, y) = (0, \zeta(x, y))$. Then,*

$$\left| \delta\mathbf{G}_f(\chi) - \int_{\Omega} \sum_i (D_x \zeta(x, f_i) + D_y \zeta(x, f_i) \cdot Df_i) : Df_i \right| \leq C \int_{\Omega} |D\zeta| |Df|^3. \quad (4.2)$$

The next two theorems deal with general \mathbf{T}_F as in Assumption 3.1. However we restrict our attention to “outer and inner variations”, where we borrow our terminology from the elasticity theory and the literature on harmonic maps. Outer variations result from deformations of the normal bundle of \mathcal{M} which are the identity on \mathcal{M} and map each fiber into itself, whereas inner variations result from composing the map F with isotopies of \mathcal{M} .

Theorem 4.2 (Expansion of outer variations). *Let \mathcal{M} , \mathbf{U} , \mathbf{p} and F be as in Assumption 3.1 with \bar{c} sufficiently small. If $\varphi \in C_c^1(\mathcal{M})$ and $X(p) := \varphi(\mathbf{p}(p))(p - \mathbf{p}(p))$, then*

$$\delta\mathbf{T}_F(X) = \int_{\mathcal{M}} \left(\varphi |DN|^2 + \sum_i (N_i \otimes D\varphi) : DN_i \right) - \underbrace{Q \int_{\mathcal{M}} \varphi \langle H, \boldsymbol{\eta} \circ N \rangle}_{\text{Err}_1} + \sum_{i=2}^3 \text{Err}_i \quad (4.3)$$

where

$$|\text{Err}_2| \leq C \int_{\mathcal{M}} |\varphi| |A|^2 |N|^2 \quad (4.4)$$

$$|\text{Err}_3| \leq C \int_{\mathcal{M}} \left(|\varphi| (|DN|^2 |N| |A| + |DN|^4) + |D\varphi| (|DN|^3 |N| + |DN| |N|^2 |A|) \right). \quad (4.5)$$

Let Y be a C^1 vector field on $T\mathcal{M}$ with compact support and define X on \mathbf{U} setting $X(p) = Y(\mathbf{p}(p))$. Let $\{\Psi_\varepsilon\}_{\varepsilon \in]-\eta, \eta]}$ be any isotopy with $\Psi_0 = \text{id}$ and $\frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \Psi_\varepsilon = Y$ and define the following isotopy of \mathbf{U} : $\Phi_\varepsilon(p) = \Psi_\varepsilon(\mathbf{p}(p)) + (p - \mathbf{p}(p))$. Clearly $X = \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \Phi_\varepsilon$.

Theorem 4.3 (Expansion of inner variations). *Let \mathcal{M} , \mathbf{U} and F be as in Assumption 3.1 with \bar{c} sufficiently small. If X is as above, then*

$$\delta\mathbf{T}_F(X) = \int_{\mathcal{M}} \left(\frac{|DN|^2}{2} \text{div}_{\mathcal{M}} Y - \sum_i DN_i : (DN_i \cdot D_{\mathcal{M}} Y) \right) + \sum_{i=1}^3 \text{Err}_i, \quad (4.6)$$

where

$$\text{Err}_1 = -Q \int_{\mathcal{M}} \left(\langle H, \boldsymbol{\eta} \circ N \rangle \text{div}_{\mathcal{M}} Y + \langle D_Y H, \boldsymbol{\eta} \circ N \rangle \right), \quad (4.7)$$

$$|\text{Err}_2| \leq C \int_{\mathcal{M}} |A|^2 (|DY| |N|^2 + |Y| |N| |DN|), \quad (4.8)$$

$$|\text{Err}_3| \leq C \int_{\mathcal{M}} \left(|Y| |A| |DN|^2 (|N| + |DN|) + |DY| (|A| |N|^2 |DN| + |DN|^4) \right). \quad (4.9)$$

4.1. Proof of Theorem 4.1. Set $\Phi_\varepsilon(x, y) := (x, y + \varepsilon \zeta(x, y))$. For ε sufficiently small Φ_ε is a diffeomorphism of $\Omega \times \mathbb{R}^n$ into itself. Moreover, $\frac{d}{d\varepsilon} \Phi_\varepsilon|_{\varepsilon=0} = \chi$. Let $f_\varepsilon = \sum_i \llbracket f_i + \varepsilon \zeta(x, f_i) \rrbracket$. Since $(\Phi_\varepsilon)_\# \mathbf{G}_f = \mathbf{G}_{f_\varepsilon}$, we can apply Corollary 3.3 to compute

$$\begin{aligned} \delta \mathbf{G}_f(\chi) &= \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \mathbf{M}(\mathbf{G}_{f_\varepsilon}) \stackrel{(3.5)}{=} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \frac{1}{2} \int \sum_i (|D(f_i + \varepsilon \zeta)|^2 + \bar{R}_4(D(f_i + \varepsilon \zeta))) \\ &= Q \int \sum_i (D_x \zeta(x, f_i) + D_y \zeta(x, f_i) \cdot Df_i) : Df_i + \int \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \bar{R}_4(Df_i + \varepsilon D\zeta). \end{aligned}$$

Since $\bar{R}_4(M) = |M|^3 L(M)$ for some Lipschitz L with $L(0) = 0$, we can estimate as follows:

$$\left| \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \bar{R}_4(M + \varepsilon \zeta) \right| \leq CL(M) |M|^2 |D\zeta| + C |M|^3 \text{Lip}(L) |D\zeta| \leq C |M|^3 |D\zeta|,$$

thus concluding the proof.

4.2. Proof of Theorem 4.2. Consider the map $\Phi_\varepsilon(p) = p + \varepsilon X(p)$. If ε is sufficiently small, Φ_ε maps \mathbf{U} diffeomorphically in a neighborhood of \mathcal{M} and we obviously have $\delta \mathbf{T}_F(X) = \frac{d}{d\varepsilon} \mathbf{M}((\Phi_\varepsilon)_\# \mathbf{T}_F)|_{\varepsilon=0}$. Next set $F_\varepsilon(x) = \sum_i \llbracket x + N_i(x)(1 + \varepsilon \varphi(x)) \rrbracket$ and observe that $(\Phi_\varepsilon)_\# \mathbf{T}_F = \mathbf{T}_{F_\varepsilon}$. Thus we can apply Theorem 3.2 to get:

$$\begin{aligned} \delta \mathbf{T}_F(X) &= \int_{\mathcal{M}} \left(\varphi |DN|^2 + \sum_i (N_i \otimes D\varphi) : DN_i \right) - \underbrace{\int_{\mathcal{M}} Q \varphi \langle H, \boldsymbol{\eta} \circ N \rangle}_{=:\text{Err}_1} \\ &\quad + \underbrace{\int_{\mathcal{M}} \sum_i \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} P_2(x, N_i(1 + \varepsilon \varphi))}_{=:\text{Err}_2} \\ &\quad + \underbrace{\int_{\mathcal{M}} \sum_i \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} (P_3(x, N_i(1 + \varepsilon \varphi), D(N_i(1 + \varepsilon \varphi))) + R_4(x, D(N_i(1 + \varepsilon \varphi))))}_{=:\text{Err}_3}. \end{aligned}$$

Since $n \mapsto P_2(x, n)$ is a quadratic form, we have $P_2(x, N_i(1 + \varepsilon \varphi)) = (1 + \varepsilon \varphi)^2 P_2(x, N_i)$ and thus (4.4) follows from (3.2). Next, by Theorem 3.2(ii), we have the bound

$$\left| \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} P_3(x, N_i(1 + \varepsilon \varphi), D(N_i(1 + \varepsilon \varphi))) \right| \leq C |A(x)| (|D\varphi| |N_i|^2 |DN_i| + |\varphi| |N_i| |DN_i|^2).$$

Finally, taking into account Theorem 3.2(iii):

$$\left| \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} R_4(x, D(N_i(1 + \varepsilon \varphi))) \right| \leq C (|DN_i|^3 + |DN_i|^3 \text{Lip}(L)) (|N_i| |D\varphi| + |DN_i| |\varphi|).$$

Putting together the last two inequalities we get (4.5).

4.3. **Proof of Theorem 4.3.** Set $F_\varepsilon(x) = \sum_i \llbracket x + N_i(\Psi_\varepsilon^{-1}(x)) \rrbracket$. Clearly, $\Phi_{\varepsilon\sharp} \mathbf{T}_F = \mathbf{T}_{F_\varepsilon}$. Fix an orthonormal frame e_1, \dots, e_m on $T\mathcal{M}$ and let $\vec{e} = e_1 \wedge \dots \wedge e_m$. By Lemma 1.9,

$$\delta \mathbf{T}_F(X) = \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \mathbf{M}(\mathbf{T}_{F_\varepsilon}) = \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int_{\mathcal{M}} \sum_i |(DF_{\varepsilon,i})\sharp \vec{e}|.$$

Fix $i \in \{1, \dots, Q\}$. Using the chain rule [4, Proposition 1.12], we have:

$$(DF_{\varepsilon,i})\sharp \vec{e} = w_1(\varepsilon, x) \wedge \dots \wedge w_m(\varepsilon, x) =: \vec{w}(\varepsilon, x),$$

where $w_j(\varepsilon, x) = e_j(x) + DN_i|_{\Psi_\varepsilon^{-1}(x)} \cdot D\Psi_\varepsilon^{-1}|_x \cdot e_j(x)$. Set $v_j(\varepsilon, x) = w_j(\varepsilon, \Psi_\varepsilon(x))$. Since Ψ_0 is the identity, we obviously have $\vec{v}(0, \cdot) = DF_{i\sharp} \vec{e}$. If we denote by $\mathbf{J}\Psi_\varepsilon(x)$ the Jacobian determinant of the transformation Ψ_ε , we can change variable in the integral to conclude:

$$\begin{aligned} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int_{\mathcal{M}} |(DF_{\varepsilon,i})\sharp \vec{e}| &= \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \int_{\mathcal{M}} |\vec{v}(\varepsilon, x)| \mathbf{J}\Psi_\varepsilon(x) \\ &= \int_{\mathcal{M}} |(DF_i)\sharp \vec{e}| \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} \mathbf{J}\Psi_\varepsilon + \int_{\mathcal{M}} |\vec{v}(0, x)|^{-1} \langle \partial_\varepsilon \vec{v}(0, x), \vec{v}(0, x) \rangle \\ &= \underbrace{\int_{\mathcal{M}} |(DF_i)\sharp \vec{e}| \operatorname{div}_{\mathcal{M}} Y}_{I_{i,1}} + \underbrace{\int_{\mathcal{M}} \langle \partial_\varepsilon \vec{v}(0, x), (DF_i)\sharp \vec{e} \rangle}_{I_{i,2}} \\ &\quad + \underbrace{\int_{\mathcal{M}} \langle \partial_\varepsilon \vec{v}(0, x), (DF_i)\sharp \vec{e} \rangle (|DF_i\sharp \vec{e}|^{-1} - 1)}_{I_{i,3}}. \end{aligned}$$

Thus, $\delta \mathbf{T}_F(X) = \sum_i I_{i,1} + \sum_i I_{i,2} + \sum_i I_{i,3} =: I_1 + I_2 + I_3$ and we will next estimate these three terms separately.

Step 1. Estimate on I_1 . By the Q -valued area formula of Lemma 1.9 and (3.4) in Theorem 3.2,

$$I_1 = Q \int_{\mathcal{M}} \operatorname{div}_{\mathcal{M}} Y + \frac{1}{2} \int_{\mathcal{M}} |DN|^2 \operatorname{div}_{\mathcal{M}} Y - Q \int \langle H, \boldsymbol{\eta} \circ N \rangle \operatorname{div}_{\mathcal{M}} Y + \operatorname{Err}$$

where $|\operatorname{Err}| \leq C \int_{\mathcal{M}} (|A|^2 |N|^2 + |DN|^4) |\operatorname{div}_{\mathcal{M}} Y|$. Since $\int_{\mathcal{M}} \operatorname{div}_{\mathcal{M}} Y = 0$ (recall that $Y \in C_c^1(\mathcal{M})$), we easily conclude that

$$I_1 = \frac{1}{2} \int_{\mathcal{M}} |DN|^2 \operatorname{div}_{\mathcal{M}} Y - Q \int \langle H, \boldsymbol{\eta} \circ N \rangle \operatorname{div}_{\mathcal{M}} Y + \sum_{j=2}^3 \operatorname{Err}_j, \quad (4.10)$$

where the Err_j 's satisfy the estimates (4.8) and (4.9).

Step 2. Estimate on I_2 . Set

$$\zeta_i(x) := \langle \partial_\varepsilon \vec{v}(0, x), (DF_i)\sharp \vec{e} \rangle = \langle \partial_\varepsilon \vec{v}(0, x), \vec{v}(0, x) \rangle = \frac{1}{2} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} |\vec{v}(\varepsilon, x)|^2.$$

Since $|\vec{v}(\varepsilon, x)|^2$ is independent of the orthonormal frame chosen, having fixed a point $x \in \mathcal{M}$, we can impose $D_{\mathcal{M}}e_j = 0$ at x . By multilinearity

$$\partial_\varepsilon \vec{v}(0, x) = \sum_j v_1(0, x) \wedge \dots \wedge \partial_\varepsilon v_j(0, x) \wedge \dots \wedge v_m(0, x). \quad (4.11)$$

We next compute

$$\begin{aligned} \partial_\varepsilon v_j(0, x) &= \frac{\partial}{\partial \varepsilon} \Big|_{\varepsilon=0} e_j(\Psi_\varepsilon(x)) + DN_i|_x \cdot \left(\frac{\partial}{\partial \varepsilon} \Big|_{\varepsilon=0} \left(D\Psi_\varepsilon^{-1} \Big|_{\Psi_\varepsilon(x)} \cdot e_j(\Psi_\varepsilon(x)) \right) \right) \\ &= D_Y e_j(x) + DN_i|_x \cdot [Y, e_j](x), \end{aligned} \quad (4.12)$$

where $[Y, e_j]$ is the Lie bracket. On the other hand, since $D_{\mathcal{M}}e_j(x) = 0$, we have $D_Y e_j(x) = A(e_j, Y)$ and $[Y, e_j](x) = -\nabla_{e_j} Y(x)$. Recall that $v_j(0, \cdot) = e_j + DN_i \cdot e_j$. By the usual computations in multilinear algebra, it turns out that $\zeta_i = \sum_j \det M^j$, where the entries of the $m \times m$ matrix M^j are given by:

$$\begin{aligned} M_{\alpha\beta}^j &= \langle e_\alpha + DN_i \cdot e_\alpha, e_\beta + DN_i \cdot e_\beta \rangle = \delta_{\alpha\beta} + O(|A||N|) + O(|DN|^2) \quad \text{for } \beta \neq j, \\ M_{\alpha j}^j &= \langle e_\alpha + DN_i \cdot e_\alpha, A(e_j, Y) - DN_i \cdot \nabla_{e_j} Y \rangle. \end{aligned}$$

(The entries for $\alpha \neq j$ are computed as in the proof of Theorem 3.2). Denote by $\text{Min}_{\alpha j}^j$ the $(m-1) \times (m-1)$ minor which is obtained by deleting the α row and the j column. We then easily get the following estimates:

$$|\text{Min}_{\alpha j}^j| \leq C(|DN|^2 + |A||N|) \quad \text{for } \alpha \neq j, \quad (4.13)$$

$$\text{Min}_{jj}^j = 1 + O(|DN|^2 + |A||N|). \quad (4.14)$$

Moreover, observe that

$$\begin{aligned} M_{\alpha j}^j &= -\langle DN_i \cdot e_\alpha, DN_i \cdot \nabla_{e_j} Y \rangle - \langle e_\alpha, DN_i \cdot \nabla_{e_j} Y \rangle + \langle A(e_\alpha, Y), DN_i \cdot e_j \rangle \\ &= -\langle DN_i \cdot e_\alpha, DN_i \cdot \nabla_{e_j} Y \rangle + \langle A(e_\alpha, \nabla_{e_j} Y), N_i \rangle + \langle A(e_\alpha, Y), DN_i \cdot e_j \rangle. \end{aligned} \quad (4.15)$$

We therefore conclude from (4.13), (4.14) and (4.15) that

$$\begin{aligned} \zeta_i(x) &= \sum_j \det M^j = \sum_j \sum_\alpha (-1)^{j+\alpha} M_{\alpha j}^j \text{Min}_{\alpha j}^j \\ &= \sum_j \left(-\langle DN_i \cdot e_j, DN_i \cdot \nabla_{e_j} Y \rangle + \langle A(e_j, \nabla_{e_j} Y), N_i \rangle + \langle A(e_j, Y), DN_i \cdot e_j \rangle \right) \\ &\quad + O\left(|DY|(|DN|^4 + |A|^2|N|^2) + |Y|(|A||DN|^3 + |A|^2|N||DN|)\right). \end{aligned} \quad (4.16)$$

Summing over i and integrating, we then achieve

$$I_2 = - \int_{\mathcal{M}} \sum_i DN_i : (DN_i \cdot D_{\mathcal{M}}Y) + J_2 + \text{Err}_2 + \text{Err}_3, \quad (4.17)$$

where $\text{Err}_2, \text{Err}_3$ are estimated as in (4.8), (4.9), and

$$J_2 = Q \int_{\mathcal{M}} \sum_j \left(\langle A(e_j, \nabla_{e_j} Y), \boldsymbol{\eta} \circ N \rangle + \langle A(e_j, Y), D_{e_j} \boldsymbol{\eta} \circ N \rangle \right).$$

In order to treat this last term, we consider the vector field $Z = \sum_j \langle A(e_j, Y), \boldsymbol{\eta} \circ N \rangle e_j$. Z is independent of the choice of the orthonormal frame e_j : therefore, to compute its divergence at a specific point $x \in \mathcal{M}$ we can assume $D_{\mathcal{M}} e_j = 0$. We then get

$$\text{div}_{\mathcal{M}} Z = \sum_j \left(\langle A(e_j, Y), D_{e_j} \boldsymbol{\eta} \circ N \rangle + \langle D_{e_j}^{\perp} A(e_j, Y), \boldsymbol{\eta} \circ N \rangle + \langle A(e_j, \nabla_{e_j} Y), \boldsymbol{\eta} \circ N \rangle \right),$$

where the tensor $D_X^{\perp} A(U, Y)$ is defined as

$$(D_X(A(U, Y)))^{\perp} - A(\nabla_X U, Y) - A(U, \nabla_X Y),$$

(recall that $(D_X W)^{\perp}$ denotes the normal component of $D_X W$). The Codazzi-Mainardi equations (cf. [16, Chapter 7.C, Corollary 15]) imply the symmetry of $D^{\perp} A$. Thus,

$$\sum_j \langle D_{e_j}^{\perp} A(e_j, Y), \boldsymbol{\eta} \circ N \rangle = \sum_j \langle D_Y^{\perp} A(e_j, e_j), \boldsymbol{\eta} \circ N \rangle = \langle D_Y^{\perp} H, \boldsymbol{\eta} \circ N \rangle. \quad (4.18)$$

Summarizing (and recalling that $\boldsymbol{\eta} \circ N$ is normal to \mathcal{M}),

$$\text{div}_{\mathcal{M}} Z = \sum_j \left(\langle A(e_j, Y), D_{e_j} \boldsymbol{\eta} \circ N \rangle + \langle A(e_j, \nabla_{e_j} Y), \boldsymbol{\eta} \circ N \rangle \right) + \langle D_Y H, \boldsymbol{\eta} \circ N \rangle. \quad (4.19)$$

Since Z is compactly supported in \mathcal{M} , integrating (4.19) and using the divergence theorem we conclude $0 = Q^{-1} J_2 + \int \langle D_Y H, \boldsymbol{\eta} \circ N \rangle$. We thus get

$$I_2 = - \int_{\mathcal{M}} \sum_i DN_i : (DN_i \cdot D_{\mathcal{M}} Y) - Q \int_{\mathcal{M}} \langle D_Y H, \boldsymbol{\eta} \circ N \rangle + \text{Err}_2 + \text{Err}_3.$$

Step 3. Estimate on I_3 . From the proof of Theorem 3.2, (cf. (3.11) and (3.12)) we conclude $|1 - |(DF_i)_{\sharp} \vec{e}| \leq C(|DN|^2 + |A||N|)$. To show that I_3 can be estimated with Err_2 and Err_3 observe that, by (4.16) we have

$$|\langle \partial_{\varepsilon} \vec{v}(0, x), (DF_i)_{\sharp} \vec{e} \rangle| = |\zeta_i(x)| \leq C|DN|^2|DY| + C|A||DY||N| + C|A||DN||Y|.$$

5. REPARAMETRIZING MULTIPLE VALUED GRAPHS

In this section we exploit the link between currents and multiple valued functions in the opposite direction, in order to give conditions under which Q -valued graphs can be suitably reparametrized and to establish relevant estimates on the parametrization. We fix the short-hand notation $\vec{e} = e_1 \wedge \dots \wedge e_{m+n}$, $\vec{e}_m = e_1 \wedge \dots \wedge e_m$ and $\vec{e}_n = e_{m+1} \wedge \dots \wedge e_{m+n}$, where $e_1, \dots, e_m, e_{m+1}, \dots, e_{m+n}$ is the standard basis of $\mathbb{R}^m \times \mathbb{R}^n$. We will often use the notation π_0 and π_0^{\perp} for $\mathbb{R}^m \times \{0\}$ and $\{0\} \times \mathbb{R}^n$.

Theorem 5.1 (*Q-valued parametrizations*). *Let $Q, m, n \in \mathbb{N}$ and $s < r < 1$. Then, there are constants $c_0, C > 0$ (depending on Q, m, n and $\frac{r}{s}$) with the following property. Let φ, \mathcal{M} and \mathbf{U} be as in Assumption 3.1 with $\Omega = B_s$ and let $f : B_r \rightarrow \mathcal{A}_Q(\mathbb{R}^n)$ be such that*

$$\|\varphi\|_{C^2} + \text{Lip}(f) \leq c_0 \quad \text{and} \quad \|\varphi\|_{C^0} + \|f\|_{C^0} \leq c_0 r. \quad (5.1)$$

Set $\Phi(x) := (x, \varphi(x))$. Then, there are maps F and N as in Assumption 3.1(N) such that $\mathbf{T}_F = \mathbf{G}_f \llcorner \mathbf{U}$ and

$$\text{Lip}(N) \leq C(\|D^2\varphi\|_{C^0}\|N\|_{C^0} + \|D\varphi\|_{C^0} + \text{Lip}(f)), \quad (5.2)$$

$$\frac{1}{2\sqrt{Q}}|N(\Phi(p))| \leq \mathcal{G}(f(p), Q[\varphi(p)]) \leq 2\sqrt{Q}|N(\Phi(p))| \quad \forall p \in B_s, \quad (5.3)$$

$$|\boldsymbol{\eta} \circ N(\Phi(p))| \leq C|\boldsymbol{\eta} \circ f(p) - \varphi(p)| + C\text{Lip}(f)|D\varphi(p)||N(\Phi(p))| \quad \forall p \in B_s. \quad (5.4)$$

Finally, assume $p \in B_s$ and $(p, \boldsymbol{\eta} \circ f(p)) = \xi + q$ for some $\xi \in \mathcal{M}$ and $q \perp T_\xi \mathcal{M}$. Then,

$$\mathcal{G}(N(\xi), Q[q]) \leq 2\sqrt{Q}\mathcal{G}(f(p), Q[\boldsymbol{\eta} \circ f(p)]). \quad (5.5)$$

For further reference, we state the following immediate corollary of Theorem 5.1, corresponding to the case of a linear φ .

Proposition 5.2 (*Q-valued graphical reparametrization*). *Let $Q, m, n \in \mathbb{N}$ and $s < r < 1$. There exist positive constants c, C (depending only on Q, m, n and $\frac{r}{s}$) with the following property. Let π_0 and π be m -planes with $|\pi - \pi_0| \leq c$ and $f : B_r(\pi_0) \rightarrow \mathcal{A}_Q(\pi_0^\perp)$ with $\text{Lip}(f) \leq c$ and $|f| \leq cr$. Then, there is a Lipschitz map $g : B_s(\pi) \rightarrow \mathcal{A}_Q(\pi^\perp)$ with $\mathbf{G}_g = \mathbf{G}_f \llcorner \mathbf{C}_s(\pi)$ and such that the following estimates hold on $B_s(\pi)$:*

$$\|g\|_{C^0} \leq Cr|\pi - \pi_0| + C\|f\|_{C^0}, \quad (5.6)$$

$$\text{Lip}(g) \leq C|\pi - \pi_0| + C\text{Lip}(f). \quad (5.7)$$

In fact the proof of Theorem 5.1 will give a more precise information about the map F , namely its pointwise values can be determined with a natural geometric algorithm.

Definition 5.3 (*Multiplicity in Q-valued maps*). Given a Q -valued map F , we say that a point p has *multiplicity k* in $F(x)$ if we can write $F(x) = k[p] + \sum_{i=1}^{Q-k} [p_i]$ where $p_i \neq p$ for every i , i.e. if p has multiplicity k when treating $F(x)$ as a 0-dimensional integral current.

Lemma 5.4 (*Geometric reparametrization*). *The values of F in Theorem 5.1 can be determined at any point $p \in \mathcal{M}$ as follows. Let \varkappa be the orthogonal complement of $T_p \mathcal{M}$. Then, $\text{Gr}(f) \cap (p + \varkappa)$ is nonempty, consists of at most Q points and every $q \in \text{Gr}(f) \cap (p + \varkappa)$ has in $F(p)$ the same multiplicity of $\mathbf{p}_{\pi_0^\perp}(q)$ in $f(\mathbf{p}_{\pi_0}(q))$.*

5.1. Existence of the parametrization. The next lemma is a natural outcome of the Ambrosio-Kirchheim approach to the theory of currents [2]. Following [9, Section 4.3], if T is a flat m -dimensional current in U and $h : U \rightarrow \mathbb{R}^k$ a Lipschitz map with $k \leq m$, we denote by $\langle T, h, y \rangle$ the slice of T with respect to h at the point y (well-defined for a.e. $y \in \mathbb{R}^k$). Since we deal with *normal* currents, the equivalence of the classical Federer-Fleming theory and the modern Ambrosio-Kirchheim theory (cf. [2, Theorem 11.1]) allows us to use all the results of the paper [2].

Lemma 5.5. *Consider a C^2 injective open curve $\gamma :]a, b[\rightarrow \mathbb{R}^N$, $\ell = \gamma(]a, b[)$, a regular tubular neighborhood $\mathbf{U}(\ell)$ and the map $\mathbf{q} := \gamma^{-1} \circ \mathbf{p}$, where \mathbf{p} is the associated C^1 normal projection $\mathbf{p} : \mathbf{U}(\ell) \rightarrow \ell$. Let T be an integral 1-dimensional current in $\mathbf{U}(\ell)$ with $\partial T = 0$ such that, for a.e. $p \in]a, b[$, the slice $F(p) := \langle T, \mathbf{q}, p \rangle$ is a sum of Q (not necessarily distinct) Dirac masses $\llbracket P_i \rrbracket$. If the measure $\mu(A) := \|T\|(\mathbf{q}^{-1}(A))$ is absolutely continuous, then $F \in W^{1,1}(]a, b[, \mathcal{A}_Q(\mathbb{R}^N))$ in the sense of [4, Definition 0.5] and $\mathcal{G}(F(p), F(p')) \leq C\mu([p, p'])$ for a.e. p, p' .*

Proof. Consider the metric space \mathbf{I}_0 of 0-dimensional integral currents endowed with the flat norm \mathbf{F} as defined in [2, Section 7]. By [2, Proof of Theorem 8.1] the map $p \mapsto F(p)$ is a \mathbf{I}_0 -valued function of bounded variation in the sense of [2, Definition 7.1], that is:

- there is a countable dense set $\mathcal{F} \subset \mathbf{I}_0$ such that, for every $S \in \mathcal{F}$, the map $\Phi_S(p) := \mathbf{F}(S, F(p))$ is a real-valued function of bounded variation;
- $|D\Phi_S|(A) \leq C\text{Lip}(\mathbf{q})\|T\|(\mathbf{q}^{-1}(A)) + C\|\mathbf{q}\|_{C^0}\|\partial T\|(\mathbf{q}^{-1}(A))$ for every Borel set A and a dimensional constant C .

On the other hand, $\partial T = 0$ and the measure $A \mapsto \mu(A) := \|T\|(\mathbf{q}^{-1}(A))$ is absolutely continuous with respect to the Lebesgue measure. By a simple density argument, it holds

$$|\Phi_S(p) - \Phi_S(q)| \leq C\mu([p, q]) \quad \forall S \in \mathbf{I}_0 \text{ and a.e. } p, q \in]a, b[. \quad (5.8)$$

Observe that by assumption $F(p)$ takes values in $\mathcal{A}_Q(\mathbb{R}^N)$ for a.e. p and, for $S = \sum_i \llbracket S_i \rrbracket, R = \sum_i \llbracket R_i \rrbracket \in \mathcal{A}_Q(\mathbb{R}^N)$, it is well known that

$$\mathbf{F}(S, R) = \min_{\pi \in \mathcal{P}_Q} \sum_i |S_i - R_{\sigma(i)}| \leq \mathcal{G}(S, R) \leq C\mathbf{F}(S, R).$$

Then, it follows from (5.8) that $|\mathcal{G}(S, F(p)) - \mathcal{G}(S, F(q))| \leq C\mathbf{F}(F(p), F(q)) \leq C\mu([p, q])$ for every $S \in \mathcal{A}_Q(\mathbb{R}^N)$. By [4, Definition 0.5], this concludes the proof. \square

The lemma can be used to infer, in a rather straightforward way, the existence of the parametrization F in Theorem 5.1

Proof of Theorem 5.1: Part I. After rescaling we can assume, without loss of generality, $r = 1$. This also easily shows that the constants depend only on the ratio $\frac{r}{s}$. We start with a procedure to identify the Q -valued function F . By (5.1), $\mathbf{G}_f \llcorner (B_1 \times \mathbb{R}^n)$ must be supported in a neighborhood of size $4c_0$ of $\Phi(B_1)$. Therefore, if the constant c_0 is chosen accordingly, the boundary of $T := \mathbf{G}_f \llcorner \mathbf{p}^{-1}(\mathcal{M})$ is actually supported in $\mathbf{p}^{-1}(\partial\mathcal{M})$ and the constancy theorem gives $\mathbf{p}_\# T = k \llbracket \mathcal{M} \rrbracket$ for some $k \in \mathbb{Z}$. First we show that $k = Q$. Consider the functions $\varphi_t := t\varphi$ for $t \in [0, 1]$, the manifolds $\mathcal{M}_t := \text{Gr}(\varphi_t)$ and the corresponding projections \mathbf{p}_t . It is simple to verify that the map

$$t \mapsto S_t := (\mathbf{p}_t)_\# (\mathbf{G}_f \llcorner (\mathbf{p}_t^{-1}(\mathcal{M}_t)))$$

is continuous in the space of currents. The constancy theorem gives $S_t = Q(t) \llbracket \mathcal{M}_t \rrbracket$ for some integer $Q(t)$ and since $S_0 = Q \llbracket \mathbb{R}^m \times \{0\} \rrbracket$, it follows that $S_1 = \mathbf{p}_\# T = Q \llbracket \mathcal{M} \rrbracket$.

Define for simplicity $\mathcal{M} \ni q \mapsto T_q := \langle \mathbf{G}_f, \mathbf{p}, q \rangle$. The integer rectifiable current \mathbf{G}_f is represented by the triple $(\text{Im}(G), \tau, \Theta)$ as in Proposition 1.4. The slicing theory gives then the following properties for \mathcal{H}^m -a.e. $p \in \mathcal{M}$ (see [9, 4.3.8]):

- (i) T_p consists of a finite sum of Dirac masses $\sum_{i=1}^{N_p} k_i \delta_{q_i}$;
- (ii) $q_i \in \text{Gr}(f)$ and $|k_i| = \Theta(q_i)$ for every i ;
- (iii) if \vec{v} is the continuous unitary m -vector orienting $\mathbf{p}^{-1}(p)$ compatibly with the orientation of \mathcal{M} , the sign of k_i is $\text{sgn}(\langle \vec{T}(q_i) \wedge \vec{v}(q_i), \vec{e} \rangle)$.

By the bounds on φ and f , $\vec{T}(x)$ is close to \vec{e}_m , while \vec{v} is close to \vec{e}_n . Therefore, each k_i turns out to be positive. On the other hand, since $\mathbf{p}_\# T = Q \llbracket \mathcal{M} \rrbracket$, then $\sum_i k_i = Q$. This shows that $p \mapsto F(p) := \sum k_i \llbracket q_i \rrbracket$ defines a Q -valued function.

Next we show the Lipschitz continuity of F . Fix a coordinate direction in \mathbb{R}^m , without loss of generality e_1 , and consider the map $\mathbf{U} \ni z \mapsto \Lambda(z) := P \circ \mathbf{p}(z)$, where $P : \mathbb{R}^{m+n} \rightarrow \mathbb{R}^{m-1}$ is the orthogonal projection $P(x_1, \dots, x_{m+n}) = (x_2, \dots, x_m)$. Consider the corresponding slice $\tilde{T}_{\bar{y}} := \langle T, \Lambda, \bar{y} \rangle$ for $\bar{y} \in \mathbb{R}^{m-1}$. For \mathcal{H}^{m-1} -a.e. $\bar{y} \in P(\mathcal{M})$, $\tilde{T}_{\bar{y}}$ is a rectifiable 1-dimensional current with $(\partial \tilde{T}_{\bar{y}}) \llcorner \mathbf{U} = 0$ (see [9, Section 4.3.1]). If we slice further $\tilde{T}_{\bar{y}}$ with respect to the map $\mathbf{p}_{\bar{y}} := x_1 \circ \mathbf{p}$, we conclude that for a.e. \bar{y} and a.e. $p \in \ell_{\bar{y}}$ we must have $\langle \tilde{T}_{\bar{y}}, \mathbf{p}_{\bar{y}}, p \rangle = F(p)$ (cf. [2, Lemma 5.1]). Applying the coarea formula to the rectifiable set \mathbf{G}_f shows also that, if c_0 is sufficiently small, then $\|T\|(\mathbf{p}_{\bar{y}}^{-1}(A)) \leq C|A|$, where C is a geometric constant (and $|\cdot|$ denotes the Lebesgue 1-dimensional measure); cf. [9, Theorem 4.3.8]. Define $]a, b[= \{t : (t, \bar{y}) \in B_s\}$, $\ell := \{\varphi(t, \bar{y}) : t \in]a, b[\}$ and $\gamma(t) := \varphi(t, \bar{y})$. It is easy to see that on $\text{spt}(\tilde{T}_{\bar{y}}$ the map $\mathbf{p}_{\bar{y}}$ coincides with the map \mathbf{q} of Lemma 5.5. Therefore the map $]a, b[\rightarrow F(t, \bar{y})$ is Lipschitz (up to a null-set). Arguing in the same way for each coordinate, we conclude that one can redefine F on a set of measure zero in such a way that F is Lipschitz: we will keep the notation F for such Lipschitz map.

Define next $N(x) = \sum_i \llbracket F_i(x) - x \rrbracket$. We then see that, by construction, N satisfies Assumption 3.1(N). Fix next coordinates on \mathcal{M} (for instance using Φ as chart). By Proposition 1.4 and the bounds on f and φ , we deduce that

$$\langle d\mathbf{p}, \vec{\mathbf{G}}_f \rangle \geq c > 0 \quad \text{and} \quad \langle d\mathbf{p}, \vec{\mathbf{T}}_F \rangle \geq c > 0,$$

for a suitable geometric constant c (where we use the notation $d\mathbf{p} = dp^1 \wedge \dots \wedge dp^m$ and $\mathbf{p}^1, \dots, \mathbf{p}^m$ are the components of \mathbf{p} in the particular chart chosen on \mathcal{M}). Hence, if $\mathbf{T}_F \neq \mathbf{G}_f \llcorner \mathbf{p}^{-1}(\mathcal{M})$, then necessarily $\mathbf{T}_F \llcorner d\mathbf{p} \neq \mathbf{G}_f \llcorner d\mathbf{p}$, which is a contradiction to $\langle T', \mathbf{p}, y \rangle = \langle T, \mathbf{p}, y \rangle$ for a.e. y (cf. [2, (5.7) and Theorem 5.6]).

Part II. To prove (5.2) consider first pairs of points $p, q \in \mathcal{M}$ with the following property:

- (AE) let $\sigma = \Phi([\mathbf{p}_{\pi_0}(p), \mathbf{p}_{\pi_0}(q)])$, $F|_\sigma = \sum \llbracket F_i \rrbracket$ with each F_i Lipschitz (cf. [4, Proposition 1.2]), and consider the corresponding curves $\gamma_i = F_i(\sigma)$: then, for \mathcal{H}^1 -a.e. $y \in \gamma_i$, $\vec{\gamma}_i(y)$ belongs to the tangent plane $T_y \mathbf{G}_f$.

We claim that (AE) implies:

$$|N(p) - N(q)| \leq C(\|D^2\varphi\|_{C^0} \|N\|_{C^0} + \text{Lip}(f) + \|D\varphi\|_{C^0}) |\mathbf{p}_{\pi_0}(p) - \mathbf{p}_{\pi_0}(q)|. \quad (5.9)$$

By standard measure theoretic arguments, (AE) holds for a set of pairs (p, q) of full measure in $\mathcal{M} \times \mathcal{M}$. With a simple density argument we then conclude the validity of (5.9) for every pair p, q . Denote by d the geodesic distance on \mathcal{M} . Since $|\mathbf{p}_{\pi_0}(p) - \mathbf{p}_{\pi_0}(q)| \leq d(p, q)$, we then conclude the Lipschitz estimate (5.2).

Let us turn to (5.9). We parameterize σ by arc-length $s : [0, \ell] \rightarrow \sigma$ and for every i define $n(t) := F_i(s(t)) - s(t)$. Clearly, n is Lipschitz and we claim that:

$$|n'(t)| \leq C(\|D^2\varphi\|_{C^0}\|n\|_{C^0} + \text{Lip}(f) + \|D\varphi\|_{C^0}) \quad \text{for a.e. } t.$$

Observe that $\frac{s'(t)+n'(t)}{|s'(t)+n'(t)|} = \vec{\gamma}_i(F_i(s(t)))$ which, for a.e. t , belongs to $T_{F_i(s(t))}\text{Gr}(f)$. The angle θ between $\vec{\gamma}_i(F_i(s(t)))$ and the plane $\mathbf{p}^{-1}(s(t))$ can then be estimated by

$$\left| \frac{\pi}{2} - \theta \right| \leq C(\text{Lip}(f) + \|D\varphi\|_{C^0}). \quad (5.10)$$

Let \mathbf{p}^T and \mathbf{p}^\perp be the projections to the tangent and normal planes to \mathcal{M} in $F_i(s(t))$. Then, if c_0 is chosen small enough to have $|n'(t)| \leq 1$, we get

$$\begin{aligned} |\mathbf{p}^\perp(n'(t))| &= |\mathbf{p}^\perp(n'(t) + s'(t))| = |n'(t) + s'(t)| |\mathbf{p}^\perp(\vec{\gamma}_i(F_i(s(t))))| \leq 2 |\cos \theta| \\ &\stackrel{(5.10)}{\leq} C(\text{Lip}(f) + \|D\varphi\|_{C^0}). \end{aligned} \quad (5.11)$$

In order to compute the tangential component, let ν_1, \dots, ν_n be an orthonormal frame on the normal bundle. It can be chosen so that $\|D\nu_j\|_{C^0} \leq C\|D^2\varphi\|_{C^0}$ for every j (see Lemma A.1). From $n(t) := \sum_j \lambda_j(t)\nu_j(s(t))$, with $\lambda_j(t) := n(t) \cdot \nu_j(s(t))$ Lipschitz functions, we get

$$\mathbf{p}^T(n'(t)) = \mathbf{p}^T \left(\sum \lambda_j'(t)\nu_j(s(t)) + \sum \lambda_j(t) \frac{d}{dt} \nu_j(s(t)) \right) = \sum \lambda_j(t) \mathbf{p}^T \left(\frac{d}{dt} \nu_j(s(t)) \right),$$

which implies

$$|\mathbf{p}^T(n'(t))| \leq \sum \|\lambda_j\|_{C^0} \|D\nu_j\|_{C^0} \leq C\|n\|_{C^0} \|D^2\varphi\|_{C^0}. \quad (5.12)$$

Putting together (5.12) and (5.11), we get (5.9). \square

5.2. Validity of the geometric algorithm. Before completing the proof of Theorem 5.1 we show Lemma 5.4, which indeed will be used to derive the remaining estimates in Theorem 5.1.

Proof of Lemma 5.4. . By the representation formula in Proposition 1.4, since the support of the push-forward via a Lipschitz map is the image of the map and we already proved $\mathbf{T}_F = \mathbf{G}_f \llcorner \mathbf{U}$, we then conclude that $\text{Im}(F) = \text{Gr}(f) \cap \mathbf{U}$ as sets. Thus, to complete the proof of Lemma 5.4 we just have to show the rule for determining the multiplicity of a point $q \in (p + \varkappa) \cap \text{Gr}(f)$ in $F(p)$. This rule follows easily from the area formula when $\text{Lip}(f)$, $\text{Lip}(N)$ and $\text{Lip}(\varphi)$ are smaller than a geometric constant, since under such assumption the Taylor expansions for the mass given by Theorem 3.2 and Corollary 3.3 imply the following facts:

- if y has multiplicity k in $f(x)$, then

$$k - \frac{1}{2} \leq \liminf_{r \downarrow 0} \frac{\|\mathbf{G}_f\|(\mathbf{B}_\rho((x, y)))}{\omega_m \rho^m} \leq \limsup_{r \downarrow 0} \frac{\|\mathbf{G}_f\|(\mathbf{B}_\rho((x, y)))}{\omega_m \rho^m} \leq k + \frac{1}{2};$$

- if p has multiplicity k in $F(x)$, then

$$k - \frac{1}{2} \leq \liminf_{r \downarrow 0} \frac{\|\mathbf{T}_F\|(\mathbf{B}_\rho(p))}{\omega_m \rho^m} \leq \limsup_{r \downarrow 0} \frac{\|\mathbf{T}_F\|(\mathbf{B}_\rho(p))}{\omega_m \rho^m} \leq k + \frac{1}{2}. \quad \square$$

We can now conclude the proof of Theorem 5.1.

Proof of Theorem 5.1: Part III. We first deal with (5.3) and (5.4). Observe first that, thanks to Lemma 5.4, the value of N at the point $(p, \varphi(p))$ does not change if we replace φ with its first order Taylor expansion. Moreover, upon translation we can further assume $p = 0$ and $\varphi(0) = 0$. We moreover fix the notation $\pi := \{(x, D\varphi(x) \cdot x) : x \in \pi_0\} = T_0 \text{Gr}(\varphi)$ and denote by \varkappa the orthogonal complement of π . With a slight abuse of notation, the same point $p \in \mathbb{R}^{m+n}$ is then represented by a pair $(x, y) \in \pi_0 \times \pi_0^\perp$ and a pair $(x', y') \in \pi \times \varkappa$. Concerning (5.3), since the role of the two systems can be reversed, it suffices to show only one inequality, namely

$$|f(0)| \leq 2\sqrt{Q}|N(0)|. \quad (5.13)$$

Let $f(0) = \sum_i \llbracket P_i \rrbracket$, $q_i := \mathbf{p}_\pi(P_i)$ and $N(q_i) = \sum_j \llbracket Q_{i,j} \rrbracket$. There is then a $j(i)$ such that $(q_i, Q_{i,j(i)}) \in \pi \times \varkappa$ is the same point as $(0, P_i) \in \pi_0 \times \pi_0^\perp$. Observing that $|q_i| \leq C\|D\varphi\|_0|P_i|$, we then get

$$\begin{aligned} |P_i| &\leq |q_i| + |Q_{i,j(i)}| \leq |q_i| + |N(0)| + \mathcal{G}(N(0), N(q_i)) \leq |N(0)| + (1 + \text{Lip}(N))|q_i| \\ &\leq |N(0)| + C(1 + \text{Lip}(N))\|D\varphi\|_0|P_i|. \end{aligned} \quad (5.14)$$

We use now (5.2) with φ linear: $\text{Lip}(N) \leq C(\|D\varphi\|_0 + \text{Lip}(f)) \leq Cc_0$. We thus conclude

$$|P_i| \leq |N(0)| + C(1 + c_0 C)c_0|P_i|.$$

However, the constant C in the last inequality is only geometric and does not depend on c_0 . Thus, if c_0 is chosen sufficiently small, we conclude $|P_i| \leq 2|N(0)|$. Summing upon i , we then reach $|f(0)| \leq 2Q^{\frac{1}{2}}|N(0)|$.

We now pass to (5.4), keeping the assumption $f(0) = \sum_i \llbracket P_i \rrbracket$ and writing $N(0) = F(0) = \sum_i \llbracket p_i \rrbracket$. Set $\mathbf{p}_{\pi_0}(p_i) = (x_i, 0)$ and $\mathbf{p}_{\pi_0^\perp}(p_i) = (0, y_i)$. The angle θ between p_i and $\mathbf{p}_{\pi_0^\perp}(p_i)$ is estimated by $C\|D\varphi(0)\|$, because the p_i 's are elements of \varkappa . Thus,

$$|x_i| \leq |p_i \sin \theta| \leq C\|D\varphi(0)\||N(0)| =: \rho. \quad (5.15)$$

Consider also that $\mathbf{p}_{\pi_0^\perp} : \varkappa \rightarrow \pi_0^\perp$ is a linear invertible map and in fact we can assume that the operator norm of its inverse, which we denote by L , is bounded by 2. Thus $|\boldsymbol{\eta} \circ N(0)| \leq 2|\sum_i y_i|$ and it suffices to estimate

$$\left| \sum y_i \right| \leq \left| \sum P_i \right| + C \text{Lip}(f)\rho. \quad (5.16)$$

To this aim, we notice that, if we set $h = \text{Lip}(f)\rho$, we can decompose $f(0)$ as $f(0) = \sum_j \llbracket T_j \rrbracket$ (where $T_j \in \mathcal{A}_{Q_j}$ and $Q_1 + \dots + Q_J = Q$) so that

- (i) $d(T_j) \leq 4Qh$, where $d(S) := \max_{i,j} |s_i - s_j|$ is the *diameter* of $S = \sum_i \llbracket s_i \rrbracket$ – cf. [4];
- (ii) $|z - w| > 4h$ for all $z \in T_j$ and $w \in T_i$ with $i \neq j$.

To prove this claim we order the P_i 's and partition them in subcollections T_1, \dots, T_k with the following algorithm. T_1 contains P_1 and any other point P_ℓ for which there exists a chain $P_{i(1)}, \dots, P_{i(\ell)} \in \text{spt}(T)$ of points with $i(1) = 1$, $i(\ell) = \ell$ and $|P_{i(\ell)} - P_{i(\ell-1)}| \leq 4h$. Clearly $d(T_1) \leq 4Qh$ and if $\text{spt}(T) = \text{spt}(T_1)$ we are finished. Otherwise we use the procedure above to define T_2 from $\text{spt}(T) \setminus \text{spt}(T_1)$, observing that $|q - p| > 4h$ for any pair of elements $q \in \text{spt}(T_1)$ and $p \in \text{spt}(T) \setminus \text{spt}(T_1)$.

By the choice of the constants, it then follows that the function f "separates" into J Lipschitz functions $f_j : B_\rho \rightarrow \mathcal{A}_{Q_j}(\mathbb{R}^n)$ with $f(x) = \sum_{j=1}^J \llbracket f^j(x) \rrbracket$ and $\text{Lip}(f_j) \leq \text{Lip}(f)$. Consider the corresponding graphs $\text{Gr}(f^j)$. Observe that, by the geometric algorithm, $N(0)$ contains points from each of these sets and moreover such points have, in $N(0)$, the same multiplicity that they have in f^j . This means that the points p_i such that $N(0) = \sum_i \llbracket p_i \rrbracket$ can actually be also grouped in J families $\{p_1^j, \dots, p_{Q_j}^j\}$ so that $N(0) = \sum_{j=1}^J \sum_{l=1}^{Q_j} \llbracket p_l^j \rrbracket$.

Note that, by the definition of the distance \mathcal{G} , for each $p_l^j \in \text{spt}(N(0))$ there exists a point $P_{k(j,l)} \in \text{spt}f^j(0)$ such that $|y_l^j - P_{k(j,l)}| \leq \mathcal{G}(f^j(\mathbf{p}_{\pi_0}(p_l^j)), f^j(0)) \leq \text{Lip}(f)|\mathbf{p}_{\pi_0}(p_l^j)| \leq h$. Thus

$$\begin{aligned} \left| \sum_i y_i \right| &= \left| \sum_{j=1}^J \sum_{l=1}^{Q_j} y_l^j \right| \leq \left| \sum_{j=1}^J \sum_{l=1}^{Q_j} P_l^j \right| + \sum_{j=1}^J \sum_{l=1}^{Q_j} |y_l^j - P_l^j| \\ &\leq \left| \sum_i P_i \right| + \sum_{j=1}^J \sum_{l=1}^{Q_j} (|y_l^j - P_{k(j,l)}| + |P_{k(j,l)} - P_l^j|) \leq \left| \sum_i P_i \right| + Ch. \end{aligned}$$

Finally, for what concerns (5.5), observe that, without loss of generality, we can assume $q = 0$ by simply shifting \mathcal{M} to $q + \mathcal{M}$: Lemma 5.4 implies that the map N' given by Theorem 5.1 applied to $q + \mathcal{M}$ satisfies $N'(\xi + q) = \sum_i \llbracket N_i(\xi) - q \rrbracket$ and so thus $\mathcal{G}(N'(\xi + q), Q \llbracket 0 \rrbracket) = \mathcal{G}(N(\xi), \llbracket q \rrbracket)$. Assuming $q = 0$ we have $\xi = (p, \boldsymbol{\eta} \circ f(p)) = (p, \boldsymbol{\varphi}(p))$ and thus the estimate matches the left hand side of (5.3). \square

APPENDIX A. TRIVIALIZING NORMAL BUNDLES

In this and the forthcoming papers the following procedure will be often used. Consider \mathcal{M} , $\boldsymbol{\varphi}$ and $\boldsymbol{\Phi}$ as in Assumption 3.1. We then construct a standard orthonormal frame on the normal bundle of \mathcal{M} as follows:

- (Tr1) we let e_{m+1}, \dots, e_{m+n} be the standard orthonormal base of $\{0\} \times \mathbb{R}^n$;
- (Tr2) for any $p \in \mathcal{M}$ we let $\boldsymbol{\varkappa}_p$ be the orthogonal complement of $T_p \mathcal{M}$ and denote by $\mathbf{p}_{\boldsymbol{\varkappa}_p}$ the orthogonal projection onto it;
- (Tr3) for any $i \in \{1, \dots, n\}$ and any $p \in \mathcal{M}$ we generate the frame $\nu_1(p), \dots, \nu_n(p)$ applying the Gram-Schmidt orthogonalization procedure to $\mathbf{p}_{\boldsymbol{\varkappa}_p}(e_{m+1}), \dots, \mathbf{p}_{\boldsymbol{\varkappa}_p}(e_{m+n})$.

We record then the following lemma.

Lemma A.1 (Trivialization of the normal bundle of \mathcal{M}). *If $\|D\boldsymbol{\varphi}\|_{C^0}$ is smaller than a geometric constant, then ν_1, \dots, ν_n is an orthonormal frame spanning $\boldsymbol{\varkappa}_p$ at every $p \in \mathcal{M}$. Consider ν_j as function of $x \in \Omega$ using the inverse of $\boldsymbol{\Phi}$ as chart. For every $\alpha + k \geq 0$*

there is a constant C depending on m, n, α, k such that, if $\|\varphi\|_{C^{k+1,\alpha}} \leq 1$, then $\|D\nu_j\|_{C^{k,\alpha}} \leq C\|\varphi\|_{C^{k+1,\alpha}}$.

REFERENCES

- [1] Frederick J. Almgren, Jr. *Almgren's big regularity paper*, volume 1 of *World Scientific Monograph Series in Mathematics*. World Scientific Publishing Co. Inc., River Edge, NJ, 2000. Q -valued functions minimizing Dirichlet's integral and the regularity of area-minimizing rectifiable currents up to codimension 2, With a preface by Jean E. Taylor and Vladimir Scheffer.
- [2] Luigi Ambrosio and Bernd Kirchheim. Currents in metric spaces. *Acta Math.*, 185(1):1–80, 2000.
- [3] Camillo De Lellis, Matteo Focardi, and Emanuele Spadaro. Lower semicontinuous functionals for Almgren's multiple valued functions. *Ann. Acad. Sci. Fenn. Math.*, 36(2):393–410, 2011.
- [4] Camillo De Lellis and Emanuele Spadaro. Q -valued functions revisited. *Mem. Amer. Math. Soc.*, 211(991):vi+79, 2011.
- [5] Camillo De Lellis and Emanuele Spadaro. Regularity of area-minimizing currents I: gradient L^p estimates. *Preprint*, 2013.
- [6] Camillo De Lellis and Emanuele Spadaro. Regularity of area-minimizing currents II: center manifold. *Preprint*, 2013.
- [7] Camillo De Lellis and Emanuele Spadaro. Regularity of area-minimizing currents III: blow-up. *Preprint*, 2013.
- [8] Manfredo Perdigão do Carmo. *Riemannian geometry*. Mathematics: Theory & Applications. Birkhäuser Boston Inc., Boston, MA, 1992. Translated from the second Portuguese edition by Francis Flaherty.
- [9] Herbert Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [10] Herbert Federer and Wendell H. Fleming. Normal and integral currents. *Ann. of Math. (2)*, 72:458–520, 1960.
- [11] Mariano Giaquinta, Giuseppe Modica, and Jiří Souček. *Cartesian currents in the calculus of variations. I*, volume 37 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 1998. Cartesian currents.
- [12] Mariano Giaquinta, Giuseppe Modica, and Jiří Souček. *Cartesian currents in the calculus of variations. II*, volume 38 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 1998. Variational integrals.
- [13] Leon Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis, Australian National University*. Australian National University Centre for Mathematical Analysis, Canberra, 1983.
- [14] Emanuele Spadaro. Complex varieties and higher integrability of Dir-minimizing Q -valued functions. *Manuscripta Math.*, 132(3-4):415–429, 2010.
- [15] Michael Spivak. *A comprehensive introduction to differential geometry. Vol. III*. Publish or Perish Inc., Wilmington, Del., second edition, 1979.
- [16] Michael Spivak. *A comprehensive introduction to differential geometry. Vol. IV*. Publish or Perish Inc., Wilmington, Del., second edition, 1979.