

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

Spatially discrete reaction-diffusion equations
with discontinuous hysteresis

by

Pavel Gurevich and Sergey Tikhomirov

Preprint no.: 25

2015



Spatially discrete reaction-diffusion equations with discontinuous hysteresis

Pavel Gurevich*, Sergey Tikhomirov†

April 15, 2015

Abstract

We deal with a class of lattice dynamical systems, namely hysteretic reaction-diffusion equations that are continuous in time and discrete in space. Our primary goal is to analyze a new mechanism that leads to appearance of a spatio-temporal pattern called *rattling*. The rattling is characterized by a specific behavior of the solution profile: while evolving in time, it oscillates up- and downwards and "switches" hysteresis at nodes satisfying a certain rule. In the one-dimensional case, this switching rule makes the profile shape two hills propagating outwards the origin with decreasing velocity. In a prototype case, we prove the existence of rattling and identify the propagation velocity.

Contents

1	Introduction	3
1.1	Background	3
1.2	Structure of the paper	8
2	Setting of the problem	11
3	Main result	14
4	Auxiliary Statements	14
4.1	Properties of the discrete Green function $y_n(t)$	15
4.2	Equivalence of some equations	16
4.3	Error estimates for Riemann sums	17

*Free University of Berlin, Peoples' Friendship University of Russia, email: gurevich@math.fu-berlin.de

†Max Planck Institute for Mathematics in the Science, Chebyshev Laboratory, Saint-Petersburg State Univeristy; email: sergey.tikhomirov@mis.mpg.de

5	Asymptotics for $u_n(t_n)$	18
5.1	Preliminaries	18
5.2	Candidates for switching moments t_n	20
5.2.1	Formulation of a theorem	20
5.2.2	Proof of Theorem 5.4	21
5.3	Proof of Proposition 5.3	24
5.4	Proof of Proposition 5.6	26
5.4.1	Proof of Proposition 5.6: Preliminaries	26
5.4.2	Proof of Proposition 5.6: Part 1	27
5.4.3	Proof of Proposition 5.6: Part 2	27
5.4.4	Proof of Proposition 5.6: Part 3	28
6	Asymptotics for $\nabla u_n(t_n)$	30
6.1	Preliminaries	31
6.2	Leading order terms	31
6.3	Remainders and proof of Theorem 6.2	32
7	Estimates of $u_n(t), u_{n+1}(t), \dots$ for $t \in (t_{n-1}, t_n)$	33
7.1	Uniqueness of a switching moment	33
7.2	Proof of Proposition 7.2: Part 1	35
7.2.1	Leading order terms	35
7.2.2	Remainders	36
7.3	Proof of Proposition 7.2: Part 2	37
7.3.1	Preliminaries	37
7.3.2	Leading order terms	38
7.3.3	Remainders	40
8	Main result: proof of Theorem 3.1	41
A	Existence and uniqueness: proof of Theorem 2.3	41
B	Equivalence of three equations: proof of Proposition 4.2	43
C	Error estimates for Riemann sums: proof of Propositions 4.3 and 4.4	44
C.1	Proof of item 2 of Proposition 4.3	44
C.2	Proof of Proposition 4.4	47
D	Requirements on $n_0(E)$	49
D.1	Constants not depending on a or E	49
D.2	Constants depending on a but not depending on E	50
D.3	Constants depending on E	54
D.4	Requirements on $n_0(E)$	54

1 Introduction

1.1 Background

Hysteresis, or, more generally, bistability, refers to a class of nonlinear phenomena which are observed in numerous real-world systems. It arises in description of ferromagnetic materials, shape-memory alloys, elasto-plastic bodies, as well as many biological, economical, and social models, see [8, 18–21, 25]. The primary goal of the present paper is to analyze a new mechanism (which we call *rattling*) for pattern formation in spatially discrete systems of reaction-diffusion equations (lattice dynamical systems) with hysteresis. The phenomenon occurs in any space dimension, including dimension one, and persists even for scalar equations. As it is explained below, our results are relevant not only for lattice dynamical systems, but also for continuous systems with hysteresis. On the other hand, they link pattern formation mechanisms in hysteretic and bistable slow-fast systems.

Let us begin with the prototype spatially continuous problem

$$\begin{cases} v_\tau = v_{xx} + \mathcal{H}(v), & x \in (-1, 1), \tau > 0, \\ v(x, 0) = \varphi(x), & x \in (-1, 1), \end{cases} \quad (1.1)$$

supplemented with, e.g., Neumann boundary conditions. Here $\mathcal{H}(\cdot)$ is the simplest *hysteresis operator*, namely, the *non-ideal relay* or *bistable switch*, see Fig. 1.1.a and the (slightly modified) rigorous definition in Section 2. Hysteresis is defined by two thresholds $\alpha < \beta$




Figure 1.1: a) Hysteresis with thresholds $\alpha < \beta$. b) Hysteresis with thresholds $\alpha = -\infty$ and $\beta = 0$.

and two values $h_1, -h_2 \in \mathbb{R}$ (in what follows, we are interested in the case $h_1 > 0 \geq -h_2$). Given a continuous *input* function $w(\tau)$, its *output* $\mathcal{H}(w)(\tau)$ remains constant unless the input achieves the lower threshold α or the upper threshold β . In the former case, the output either *switches* to h_1 if it was equal to $-h_2$ “just before” or otherwise remains h_1 . Analogously, in the latter case, the output either *switches* to $-h_2$ if it was equal to h_1 “just before” or otherwise remains $-h_2$. Since the function $v(x, \tau)$ in (1.1) depends not only on τ , but also on the spatial variable x , one defines $\mathcal{H}(v) = \mathcal{H}(v(x, \cdot))(\tau)$ “pointwise”, i.e., for each fixed x . Thus, the hysteresis operator \mathcal{H} becomes *spatially distributed*.

Problem (1.1) is the simplest model of a reaction-diffusion process in which a diffusive substance with density $v(x, \tau)$ interacts in a hysteretic way with a non-diffusive substance that affects the diffusive one via the reaction term taking values h_1 or $-h_2$. The first model

of such a type was suggested by Hoppensteadt and Jäger [14]. It consisted of two reaction-diffusion equations and one ordinary differential equation and described the concentric rings pattern that occurs in a colony of bacteria (*Salmonella typhimurium*) on a Petri plate, see Fig. 1.2. Numerical simulations in [14, 15] yielded a pattern that was consistent with




Figure 1.2: Density of bacteria on a Petri plate after growth has stopped as a result of a numerical simulation via spatial discretization.

experiments, however the rigorous mathematical description of the model was lacking. To begin with, the well-posedness was an open question, due to the discontinuous nature of the hysteresis operator. First analytical results were obtained in [3, 24] (see also [2, 17, 25] and a recent survey [26]), where existence of solutions for multi-valued hysteresis was proved. Formal asymptotic expansions of solutions were recently obtained in a special case in [16]. Questions about the uniqueness of solutions and their continuous dependence on the initial data as well as a thorough analysis of pattern formation still remained open.

In [11, 12], we formulated the so-called *transversality* condition for the initial data $\varphi(x)$ in (1.1) that guaranteed existence, uniqueness, and continuous dependence of solutions on initial data for scalar equations with hysteresis. In [13], this condition was generalized to systems, and in [9] to the case $x \in \mathbb{R}^2$. For problem (1.1), the transversality loosely speaking means that if $\varphi(x_0) = \alpha$ or $\varphi(x_0) = \beta$ for some $x_0 \in (-1, 1)$, then $\varphi'(x_0) \neq 0$. Due to [11–13], the solution either exists and is unique for all $\tau \in [0, \infty)$, or there is $T > 0$ such that the solution exists and is unique for $\tau \in [0, T]$ and $v(x, T)$ is not transverse. The approach of [11–13] was based on treating the problem with transverse initial data as a special free boundary problem. The study of regularity of the emerging free boundary was initiated in [5, 6], based on methods developed for obstacle-type problems (see, e.g., [7, 22, 23] and the references therein).

The key question which we address in this paper is how the solution may behave after it becomes nontransverse. To answer this question, we consider the nontransverse initial data. First, set $\beta = 0$ (without loss of generality) and consider an initial function $\varphi(x) = -cx^2 + o(x^2)$ in a neighborhood $\mathcal{B}(0)$ of $x = 0$. By taking a smaller neighborhood if needed, we have $\varphi(x) < 0$ for $x \in \mathcal{B}(0) \setminus \{0\}$. We define the hysteresis at the initial moment in this neighborhood as follows: $\mathcal{H}(\varphi(0)) = -h_2$ and $\mathcal{H}(\varphi(x)) = h_1$ for $x \neq 0$. Now we “regularize” the parabolic equation in $\mathcal{B}(0)$ by discretizing the spatial variable: for any $\varepsilon > 0$, setting

$v_n(\tau; \varepsilon) := v(\varepsilon n, \tau)$, we replace the continuous model (1.1) in $\mathcal{B}(0)$ by the discrete one

$$\begin{cases} \frac{dv_n}{d\tau} = \frac{\Delta v_n}{\varepsilon^2} + \mathcal{H}(v_n), & \tau > 0, \quad n = -N_\varepsilon, \dots, N_\varepsilon, \\ v_n(0) = -c(\varepsilon n)^2 + o(\varepsilon^2 n^2), & n = -N_\varepsilon, \dots, N_\varepsilon, \end{cases} \quad (1.2)$$

where $\Delta v_n := v_{n+1} - 2v_n + v_{n-1}$ and $N_\varepsilon \rightarrow \infty$ as $\varepsilon \rightarrow 0$. Since we are interested in small ε and in the behavior near the threshold $\beta = 0$ (i.e., in a small neighborhood $\mathcal{B}(0)$), we consider the next approximation by omitting $o(\varepsilon^2 n^2)$ in the initial data, replacing N_ε by ∞ , and formally setting $\alpha := -\infty$. Thus, (1.2) assumes the form

$$\begin{cases} \frac{dv_n}{d\tau} = \frac{\Delta v_n}{\varepsilon^2} + \mathcal{H}(v_n), & \tau > 0, \quad n \in \mathbb{Z}, \\ v_n(0) = -c(\varepsilon n)^2, & n \in \mathbb{Z}, \end{cases} \quad (1.3)$$

and the hysteresis operator is now represented by Fig. 1.1.b (see the rigorous definition in Section 2).

A nontrivial dynamics occurs in the case $h_1 > 2c > 0 \geq -h_2$. To indicate the difficulty, note that, due to the initial configuration of hysteresis, we have $\frac{dv_0}{d\tau}(0; \varepsilon) = -h_2 - 2c < 0$, but $\frac{\partial v_n}{\partial \tau}(0; \varepsilon) = h_1 - 2c > 0$ for $n \in \mathbb{Z} \setminus \{0\}$. Thus, for small $\tau > 0$, $v_0(\tau; \varepsilon)$ decreases, while all the other nodes $v_n(\tau; \varepsilon)$, $n \in \mathbb{Z} \setminus \{0\}$, increase. It is not clear at all, which node achieves the threshold $\beta = 0$ and switches first and hence what a further dynamics is.

In fact, numerical analysis does not reveal any general rule that could describe the behavior of $v_n(\tau; \varepsilon)$ for small τ . However, it reveals the formation of quite a specific spatio-temporal pattern for large τ , see Fig. 1.3. If $h_2 = 0$, then each node eventually achieves the threshold




Figure 1.3: Upper graphs represent spatial profiles of the hysteresis $\mathcal{H}(v_n)$ and lower graphs the spatial profiles of the solution v_n . a) Nontransverse initial data. b) Spatial profiles at a moment $\tau > 0$ for $h_2 = 0$. c) Spatial profiles at a moment $\tau > 0$ for $h_2 = h_1 > 0$.

$\beta = 0$ and thus $\mathcal{H}(v_n)$ eventually switches from h_1 to $h_2 = 0$ for each $n \in \mathbb{Z}$. If $h_2 > 0$, then some nodes achieve the threshold and some do not. If we denote by $N_1(j)$ and $N_2(j)$ the number of nodes in the set $\{v_0, v_{\pm 1}, \dots, v_{\pm j}\}$ that switch and do not switch, respectively, on the time interval $[0, \infty)$, then numerics suggests that

$$\lim_{j \rightarrow \infty} \frac{N_2(j)}{N_1(j)} = \frac{h_2}{h_1}. \quad (1.4)$$

Moreover, if $h_2/h_1 = p_2/p_1$, where p_1 and p_2 are co-prime integers, then, for any j large enough, the set $\{v_{j+1}, \dots, v_{j+p_1+p_2}\}$ contains exactly p_1 nodes that switch and p_2 nodes that do not switch on the time interval $[0, \infty)$.

The next numerical observation is as follows. Let $\tau_n = \tau_n(\varepsilon)$ be the switching moment of the node $v_n(\tau; \varepsilon)$ if this node switches on the time interval $[0, \infty)$ and $\tau_n := \infty$ otherwise. Then, for any fixed $h_2 \geq 0$, the τ_n 's that are finite satisfy, as $n \rightarrow \infty$,

$$\tau_n = a(\varepsilon n)^2 + \begin{cases} \varepsilon^2 O(\sqrt{n}) & \text{if } h_2 = 0, \\ \varepsilon^2 O(n) & \text{if } h_2 > 0, \end{cases} \quad (1.5)$$

where $a > 0$ depends on h_1/c but *does not depend* on h_2 or ε and $O(\cdot)$ does not depend on ε .

Remark 1.1. In Section 1.2, we will show that ε in (1.3) can be scaled out, see scaling (1.8). In particular, all the numerical observations concerning the dynamics of v_n have been done for $\varepsilon = 1$ and then transferred to an arbitrary ε according to the scaling in (1.8).

Consider the function

$$H(x, \tau; \varepsilon) := \mathcal{H}(v_n(\cdot; \varepsilon))(\tau), \quad x \in [\varepsilon n - \varepsilon/2, \varepsilon n + \varepsilon/2], \quad n \in \mathbb{Z},$$

which is supposed to approximate the hysteresis $\mathcal{H}(v(x, \cdot))(\tau)$ in (1.1). Assuming the dynamics (1.4) and (1.5) and taking into account Remark 1.1, we see that $H(x, \tau; \varepsilon)$ has no pointwise limit as $\varepsilon \rightarrow 0$, but converges in a certain weak sense to the function $H(x, \tau)$ given by $H(x, \tau) = 0$ for $\tau > ax^2$ and $H(x, \tau) = h_1$ for $\tau < ax^2$. We emphasize that $H(x, \tau)$ does not depend on h_2 (because a does not). On the other hand, if $h_2 > 0$, the hysteresis operator $\mathcal{H}(v(x, \cdot))(\tau)$ in (1.1) cannot take value 0 by definition, which clarifies the essential difficulty with the well-posedness of the original problem (1.1) in the nontransverse case. To overcome the non-wellposedness, one needs to allow the intermediate value 0 for the hysteresis operator.

Such a re-definition of hysteresis is consistent with the behavior of $v(x, \tau)$ (also observed numerically) in the following sense. For a fixed $\varepsilon > 0$, the spatial profile of $v_n(\cdot; \varepsilon)(\tau)$ forms two humps propagating away from the origin according to (1.5). The cavity between the humps has a bounded steepness characterized by the relations

$$|v_{k+1}(\tau; \varepsilon) - v_k(\tau; \varepsilon)| \leq b\varepsilon^2, \quad |k| \leq n, \quad \tau \geq \tau_n, \quad n = 0, 1, 2, \dots, \quad (1.6)$$

where $b > 0$ does not depend on k , n , and ε . As time goes on, the profile executes downwards and upwards motions, always remaining beneath the threshold $\beta = 0$ and hitting this threshold at specific nodes characterized by (1.4). We call such a behavior of v_n and $\mathcal{H}(v_n)$ *rattling*. Furthermore, numerics indicates that, as $\varepsilon \rightarrow 0$, the function

$$V(x, \tau; \varepsilon) := v_n(\tau; \varepsilon), \quad x \in [\varepsilon n - \varepsilon/2, \varepsilon n + \varepsilon/2], \quad n \in \mathbb{Z},$$

approximates a smooth function $V(x, \tau)$, which satisfies $V(x, \tau) = 0$ for $\tau > ax^2$ due to (1.5) and (1.6). In other words, $V(x, \tau)$ sticks to the threshold line $\beta = 0$ on the expanding interval $x \in (-\sqrt{\tau/a}, \sqrt{\tau/a})$.

We recall paper [3], in which Alt proved the existence of a function $V(x, \tau)$ that satisfies the equation in (1.1) on the set $A = \{(x, \tau) : V(x, \tau) \neq \alpha, \beta\}$ and the homogeneous heat equation on the set $B = \{(x, \tau) : V(x, \tau) = \alpha \text{ or } \beta\}$. Thus, our heuristic argument provides a qualitative description of the sets A and B and justifies the completion of hysteresis by the zero value via the thermodynamical limit. To make this argument mathematically rigorous, we should first rigorously describe the rattling phenomenon in the discrete system (1.3). This is the central topic of the present paper, in which we concentrate on the case $h_2 = 0$ and develop general tools for treating discrete reaction-diffusion equations with discontinuous hysteresis. The application of these tools to the case $h_2 > 0$ will be a subject of a forthcoming paper. We expect that these tools will be applicable whenever h_2/h_1 is rational.

Before we proceed with the description of our tools and of the structure of the paper, let us make two more comments. First, the rattling phenomenon also occurs in multidimensional domains. For example, Fig. 1.4 illustrates the switching pattern for a two-dimensional analog of (1.3), where we have implemented spatial discretizations on the square and triangular lattices, respectively. Moreover, numerical analysis of the Hoppensteadt–Jäger system




Figure 1.4: A snapshot for a time moment $\tau > 0$ of a two-dimensional spatial profile of hysteresis taking values $h_1 > 2c > 0$ and $h_2 = h_1 > 0$. The nontransverse initial data is given by $\varphi(x) = -c(x_1^2 + x_2^2)$. Grey (black) squares or hexagons correspond to the nodes that have (not) switched on the time interval $[0, \tau]$. a) Discretization on the square lattice. b) Discretization on the triangular lattice.

indicates that the solution remains transverse as long as the central disc in Fig. 1.2 gets formed, but the formation of all the rings occurs via rattling.

Second, the rattling phenomenon is not a pure consequence of a discontinuous nature of hysteresis, but rather a consequence of bistability in a system. In particular, it persists in bistable slow-fast reaction-diffusion systems. The simplest example is the system of two equations

$$v_\tau = v_{xx} + w, \quad \delta w_\tau = f(v, w), \quad (1.7)$$

where $\delta > 0$ is a small parameter and the nullcline of $f(v, w)$ is Z - or S -shaped. Formally, system (1.7) can be treated as another regularization of system (1.1). In the case where the nullcline of $f(v, w)$ is S -shaped, one should replace h_1 and $-h_2$ in the definition of hysteresis $\mathcal{H}(v)$ by appropriate functions $H_1(v)$ and $H_2(v)$, see Fig. 1.5.

As $\delta \rightarrow 0$, the spatial profiles of v and w in (1.7) behave similarly to $V(x, \tau; \varepsilon)$ and $H(x, \tau; \varepsilon)$, respectively, as $\varepsilon \rightarrow 0$, see Fig. 1.6, with the exception that the profile of w




Figure 1.5: a) The nullcline of the S -shaped nonlinearity $f(v, w)$. b) Hysteresis with non-constant branches $H_1(v)$ and $H_2(v)$.

remains continuous and forms steep transition layers between mildly sloping steps of width tending to 0 as $\delta \rightarrow 0$. Interestingly, the time-scale separation parameter δ in (1.7) yields




Figure 1.6: Upper and lower graphs are spatial profiles of the solution $w(x, \tau)$ and $v(x, \tau)$, respectively, for problem (1.7) with initial data $v|_{\tau=0} = -cx^2 + o(x^2)$, $w|_{\tau=0} = h_1$.

the same effect as the grid-size parameter ε in (1.3). As far as we know, such a rattling phenomenon for slow-fast systems has not been explained in the literature, either.

1.2 Structure of the paper

Now we come back to the main topic of this paper, namely, discrete system (1.3). As it was mentioned in Remark 1.1, ε in (1.3) can be scaled out. Indeed, setting

$$t := \varepsilon^{-2}\tau, \quad u_n(t) := \varepsilon^{-2}v_n(\tau; \varepsilon) \quad (1.8)$$

and using the equalities (recall that $\alpha = -\infty$ and $\beta = 0$)

$$\mathcal{H}(v_n)(\tau) = \mathcal{H}(\varepsilon^2 u_n(\varepsilon^{-2}\cdot))(\tau) = \mathcal{H}(u_n(\varepsilon^{-2}\cdot))(\tau) = \mathcal{H}(u_n)(\varepsilon^{-2}\tau) = \mathcal{H}(u_n)(t),$$

we can rewrite (1.3) as follows:

$$\begin{cases} \dot{u}_n = \Delta u_n + \mathcal{H}(u_n), & t > 0, n \in \mathbb{Z}, \\ u_n(0) = -cn^2, & n \in \mathbb{Z}, \end{cases} \quad (1.9)$$

where $\dot{} = d/dt$. Problem (1.9) does not involve ε , which justifies the fact that $u_n(t)$ in (1.8) does not depend on ε . Note that c in (1.9) could be also scaled out if we replaced $u_n(t)$ by $c\tilde{u}_n(t)$ and h_1 and $-h_2$ by \tilde{h}_1 and $-\tilde{h}_2$, respectively. However, we prefer not to do this, in order to keep track of what exactly is influenced in our intermediate calculations by the ‘‘tangency’’ constant c .

From now on, we concentrate on the case $h_2 = 0$. Due to (1.5) and (1.8), the asymptotics for the switching moment t_n of the node $u_n(t)$ is expected to be

$$t_n = an^2 + q_n, \quad |q_n| \leq E\sqrt{n}, \quad (1.10)$$

where $E > 0$ does not depend on $n \in \mathbb{Z}$.

Our *main result* is formulated as follows (see Theorem 3.1). Let $h_1 > 2c > 0$ and $h_2 = 0$. Assume that

$$\begin{aligned} & \text{finitely many nodes } u_0(t), u_1(t), \dots, u_{n_0}(t) \\ & \text{switch at time moments } t_n \text{ satisfying (1.10),} \end{aligned} \quad (1.11)$$

where the constants $a = a(h_1/c) > 0$ and $n_0 = n_0(E) = n_0(E, h_1, c)$ will be explicitly specified in the main text. Then each node $u_n(t)$, $n \in \mathbb{Z}$, switches; moreover, the switching occurs at a time moment t_n satisfying (1.10).

Since we will provide an explicit formula for the solution $u_n(t)$, the fulfillment of finitely many assumptions (1.11) can be verified numerically with an arbitrary accuracy for any given values of h_1 and c (see Fig. 1.7).




Figure 1.7: Values of q_n/\sqrt{n} for $c = 1/2$ and a) $h_1 = 1.5$, b) $h_1 = 2.0$.

The paper is organized as follows. In Section 2, we give definitions for the hysteresis operator and for the solution of problem (1.9). Next, we formulate the existence and uniqueness theorem (Theorem 2.3), which includes a representation of the solution $u_n(t)$ via the so-called *discrete Green function* $y_n(t)$. In particular, Theorem 2.3 implies that $u_n(t) = u_{-n}(t)$, $n \in \mathbb{Z}$. This theorem is proved in Appendix A.

In Section 3, we formulate our main result (Theorem 3.1).

Section 4 contains information about three main ingredients for the proof of the main result.

1. The first ingredient is asymptotic formulas for the Green function $y_n(t)$ and for its time derivatives, which were derived in [10].
2. The second ingredient is three equations for finding the constant a from (1.10). The equivalence of these equations as well as the existence and uniqueness of their root $a > 0$ are proved in Appendix B.
3. The third ingredient is the approximation of some singular integrals by Riemann sums and corresponding error estimates, which are proved in Appendix C.

Sections 5, 6, and 7 are three key steps in the proof of our main result. The scheme of the proof is inductive. Assume we have proved that t_0, t_1, \dots, t_{n-1} satisfy (1.10) for some fixed $n \geq n_0 + 1$. We *fix* the hysteresis configuration, i.e., set $H_n := \mathcal{H}(u_n)(t_{n-1})$ and consider the solution $v_n(t)$ of the problem

$$\begin{cases} \dot{v}_n = \Delta v_n + H_n, & t > t_{n-1}, \quad n \in \mathbb{Z}, \\ v_n(t_{n-1}) = u(t_{n-1}), & n \in \mathbb{Z} \end{cases}$$

(we abuse the notation by using the same letter v as in Section 1.1). Obviously, $v_n(t) = u_n(t)$ as long as the nodes $v_n(t), v_{n+1}(t), v_{n+2}(t), \dots$ remain below the threshold $\beta = 0$.

The main theorem of Section 5 (Theorem 5.4) claims that the equation $v_n(an^2 + q_n) = 0$ has a root q_n satisfying (1.10). To prove this, we use an explicit representation of $v_n(t)$ via the convolution of H_n with the Green function $y_n(t)$ (see (5.20)). Then we use asymptotic formulas for $y_n(t)$ (recall the first ingredient from Section 4) and replace the convolution by a singular integral (recall the third ingredient from Section 4). As a result, we obtain a leading order term of order n^2 , which depends only on a and h_1/c , and a remainder of order \sqrt{n} , which also depends on q_0, q_1, \dots, q_{n-1} (that are known due to the inductive hypothesis) and on the unknown q_n . It appears that the coefficient at n^2 vanishes due to the choice of a (recall the second ingredient from Section 4). The hard part is to show that the remainder vanishes for some q_n satisfying (1.10). This is done by an application of Brouwer's fixed-point theorem.

The time moment $t_n := an^2 + q_n$ given by Theorem 5.4 is a *candidate* for being the switching moment of $u_n(t)$. In order that it be the switching moment, we have to prove that neither of the nodes $v_{n+1}(t), v_{n+2}(t), \dots$ achieves the threshold $\beta = 0$ on the time interval $(t_{n-1}, t_n]$, while $v_n(t)$ achieves it at the moment t_n for the first time. This is done in Sections 6 and 7.

In Section 6, we prove that $v_{n+1}(t_n) < 0$ (Theorem 6.2). To do so, we estimate the gradient $\nabla v_n(t_n) := v_{n+1}(t_n) - v_n(t_n)$ by using the representation of $\nabla v_n(t)$ via the gradient $\nabla y_n(t)$ of the Green function, applying asymptotic formulas for $\nabla y_n(t)$ (recall the first ingredient from Section 4) and again replacing the corresponding convolution by an integral (recall the

third ingredient from Section 4). It appears that the leading order term of order n vanishes due to the second ingredient from Section 4. Thus, we have to calculate the next term in the asymptotics, which turns out to be $-3h_1/4 < 0$. Hence, $v_{n+1}(t_n) = \nabla v_n(t_n) \leq -3h_1/8 < 0$.

In Section 7, we first show that $v_n(t)$ does not achieve the threshold $\beta = 0$ for $t \in (t_{n-1}, t_n)$ (Theorem 7.1). To do so, we divide the interval (t_{n-1}, t_n) into two parts. We prove that the function $\dot{v}_n(t)$ is so small on the first interval that it cannot overcome the distance exceeding $-3h_1/8$ (the value coming from Theorem 6.2 with $n + 1$ replaced by n). Then we prove that $\ddot{v}_n(t)$ is nonnegative on the second interval. Hence, the equation $v_n(t) = 0$ has a unique root, which must be t_n . In particular, $v_n(t) < 0$ for $t \in (t_{n-1}, t_n)$. Finally, we show that $\nabla v_j(t) < 0$ for all $t \in (t_{n-1}, t_n]$ and $j \geq n$, which implies that the nodes $v_{n+1}(t), v_{n+2}(t), \dots$ remain negative for $t \in (t_{n-1}, t_n]$ (Theorem 7.3).

In Section 8, we combine the results from Sections 5, 6, and 7 and rigorously implement the inductive scheme, which completes the proof of the main result, namely, Theorem 3.1.

The crucial role in our main result (Theorem 3.1) is played by the number $n_0 = n_0(E)$, which determines the number of switchings one has to check “by hand” (see (1.11)). The number $n_0(E)$ is determined explicitly in the form of 12 inequalities that must hold for $n \geq n_0(E)$. Each inequality is referred to as a **requirement** and is introduced in the text where it is used for the first time. These 12 requirements contain constants that are also introduced in the text where they are needed for the first time. For reader’s convenience, we have collected all the constants in Appendices D.1–D.3 and the 12 requirements determining $n_0(E)$ in Appendix D.4.

The graphs in Fig. 1.8 represent the values of a , E , and $n_0(E)$ that fulfill assumptions (1.11) for $c = 1/2$ and $h_1 = 1.1, 1.2, 1.3, \dots, 2.5$.




Figure 1.8: Dependence on h_1 of the values of a , E , and $n_0(E)$ that fulfill assumptions (1.11) for $c = 1/2$. a) The values of a are found explicitly for all $h_1 > 1$ (see Section 4.2 below). b), c) The values of E and $n_0(E)$ are calculated numerically for $h_1 = 1.1, 1.2, 1.3, \dots, 2.5$.

2 Setting of the problem

For a sequence $\{v_n\}_{n \in \mathbb{Z}}$ of real numbers, we use the notation

$$\nabla v_n := v_{n+1} - v_n, \quad \Delta v_n := \nabla v_n - \nabla v_{n-1} = v_{n-1} - 2v_n + v_{n+1}.$$

Let $\{u_n(t)\}_{n \in \mathbb{Z}}$ be a sequence of real-valued functions defined for $t \geq 0$. We study the problem

$$\dot{u}_n = \Delta u_n + \mathcal{H}(u_n), \quad t > 0, \quad n \in \mathbb{Z}, \quad (2.1)$$

$$u_n(0) = -cn^2, \quad n \in \mathbb{Z}, \quad (2.2)$$

where $c > 0$ and $\mathcal{H}(w)(t)$, $t \geq 0$, is the hysteresis operator defined for functions $w \in C[0, \infty)$ such that $v(0) \leq 0$ by

$$\mathcal{H}(w)(t) := \begin{cases} h_1 & \text{if } w(s) < 0 \text{ for all } s \in [0, t], \\ 0 & \text{if } w(s) = 0 \text{ for some } s \in [0, t], \end{cases} \quad (2.3)$$

where $h_1 > 0$ is fixed. In other words, the output of hysteresis is h_1 unless the input achieves the zero threshold; at this moment, the hysteresis *switches* and since then the output of hysteresis remains 0. In the context of problem (2.1)–(2.3), we will use the phrasing “a node $u_n(t)$ switches” or just “a node n switches” whenever $u_n(t)$ achieves the value zero for the first time.

Remark 2.1. In the terminology of, e.g., [18,25], the hysteresis operator (2.3) is a non-ideal relay with the thresholds $-\infty$ and 0; see Fig. 1.1.b.

From now on, we assume throughout that the following condition holds.

Condition 2.1. $h_1 > 2c > 0$.

We note that the function $\mathcal{H}(v)(t)$ may have discontinuity (actually, at most one) even if $v \in C^\infty[0, \infty)$. Therefore, one cannot expect that a solution of problem (2.1)–(2.3) is continuously differentiable on $[0, \infty)$. Thus, we define a solution as follows.

Definition 2.1. We say that a sequence $\{u_n(t)\}_{n \in \mathbb{Z}}$ is a *solution of problem (2.1)–(2.3) on the time interval $(0, T)$* , $T > 0$, if

1. $u_n \in C[0, T]$ for all $n \in \mathbb{Z}$,
2. for each $t \in [0, T]$, there are constants $A, \alpha \geq 0$ such that $\sup_{s \in [0, t]} |u_n(s)| \leq Ae^{\alpha|n|}$ for all $n \in \mathbb{Z}$,
3. there is a finite sequence $0 = \tau_0 < \tau_1 < \dots < \tau_J = T$, $J \geq 1$, such that $u_n \in C^1(\tau_j, \tau_{j+1})$ for all $n \in \mathbb{Z}$ and $j = 0, \dots, J - 1$,
4. the equations in (2.1) hold in (τ_j, τ_{j+1}) for all $n \in \mathbb{Z}$ and $j = 0, \dots, J - 1$,
5. $u_n(0) = -cn^2$ for all $n \in \mathbb{Z}$.

We say that a sequence $\{u_n(t)\}_{n \in \mathbb{Z}}$ is a *solution of problem (2.1)–(2.3) on the time interval $(0, \infty)$* if it is a solution on $(0, T)$ for all $T > 0$.

Remark 2.2. If $\{u_n(t)\}_{n \in \mathbb{Z}}$ is a solution, then, as we have mentioned above, the function $\mathcal{H}(u_n)(t)$ has at most one discontinuity point for each fixed $n \in \mathbb{Z}$. Hence, the equations in (2.1) imply that each function $\dot{u}_n(t)$ has at most one discontinuity point on $[0, \infty)$.

Before we treat existence and uniqueness of a solution, let us introduce one of our main tools, namely, the so-called *discrete Green function*

$$y_n(t) := \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1 - e^{-2t(1-\cos\theta)}}{2(1-\cos\theta)} e^{in\theta} d\theta, \quad t \geq 0. \quad (2.4)$$

One can directly check that $y_n \in C^\infty[0, \infty)$ and $y_n(t)$ solves the problem

$$\begin{cases} \dot{y}_0 = \Delta y_0 + 1, & t > 0, \\ \dot{y}_n = \Delta y_n, & t > 0, n \neq 0, \\ y_n(0) = 0, & n \in \mathbb{Z}. \end{cases} \quad (2.5)$$

Below, we will use the fact that

$$\dot{y}_{n+1}(t) < \dot{y}_n(t), \quad t > 0, n = 0, 1, 2, \dots, \quad (2.6)$$

which follows from the formula $\dot{y}_n(t) = e^{-2t} I_n(2t)$, where $I_n(s)$ is the modified Bessel function of the first kind (see [1, Sec. 9.6.19]), and from, e.g., [4]. We will also use the estimate, which follows from the series representation of the modified Bessel function [1, Sec. 9.6.10]:

$$0 \leq \dot{y}_n(t) = e^{-2t} I_n(2t) = e^{-2t} t^{|n|} \sum_{m=0}^{\infty} \frac{t^{2m}}{m!(m+|n|)!} \leq \frac{e^{t^2-2t}|n|!}{|n|!}, \quad n \in \mathbb{Z}. \quad (2.7)$$

The following existence and uniqueness result is proved in Appendix A.

Theorem 2.3. 1. Problem (2.1)–(2.3) has a unique solution $\{u_n(t)\}_{n \in \mathbb{Z}}$ on the time interval $(0, \infty)$.

2. Let t_n be the switching moment of the node $u_n(t)$ if this node switches on the time interval $[0, \infty)$ and $t_n := \infty$ otherwise. Then

$$t_n \geq \frac{cn^2}{h_1 - 2c}, \quad n \in \mathbb{Z}. \quad (2.8)$$

3. Let $S(t)$ be the set of nodes that switch on the time interval $[0, t]$, i.e.,

$$S(t) := \{k \in \mathbb{Z} : \mathcal{H}(u_k)(t) = 0\}, \quad (2.9)$$

and let $|S(t)|$ be the number of elements in $S(t)$. Then $S(t)$ is finite for each $t > 0$, symmetric with respect to the origin, $|S(t)| \rightarrow \infty$ as $t \rightarrow \infty$, and

$$u_n(t) = -cn^2 + (h_1 - 2c)t - h_1 \sum_{k \in S(t)} y_{n-k}(t - t_k), \quad t \in [0, \infty), \quad (2.10)$$

where we put $y_{n-k}(t - t_k) = 0$ for $t < t_k$,

4. for each $n \in \mathbb{N}$, we have $t_{-n} = t_n$ and $u_{-n}(t) \equiv u_n(t)$.

3 Main result

We recall that Condition 2.1 is assumed to hold throughout. Below in the text we define $a > 0$ (see Lemma 4.2), $E_0 > 0$ (see (5.13)) and an increasing function $n_0 : (E_0, \infty) \rightarrow \mathbb{N}$ (see Requirements 1–12 in Section D.4).

Definition 3.1. We say that a number $E \geq E_0$ is *admissible* if the following holds:

1. each node u_k , $k = 0, \pm 1, \dots, \pm n_0 = \pm n_0(E)$ switches at a moment t_k satisfying

$$t_k = ak^2 + q_k, \quad |q_k| \leq E\sqrt{n_0}, \quad (3.1)$$

while neither of the nodes $u_{\pm(n_0+1)}, u_{\pm(n_0+2)}, \dots$ switches on the time interval $[0, t_{n_0}]$;

2. at the switching moment t_{n_0} , we have

$$u_{n_0+1}(t_{n_0}) = \nabla u_{n_0}(t_{n_0}) \leq -\frac{3h_1}{8}. \quad (3.2)$$

The main result of this paper is as follows. If finitely many nodes $k = 0, \dots, n_0(E)$ switch at time moments t_k satisfying (3.1), then *all* the nodes $n \in \mathbb{Z}$ will switch and their switching moments will be of order an^2 . On Fig. 1.8.b, one can see the values of admissible E , which we found numerically for $c = 1/2$ and $h_1 = 1.1, 1.2, 1.3, \dots, 2.5$. Figures 1.8.a and 1.8.c depict corresponding values of a and $n_0(E)$, respectively.

The rigorous formulation of our main result is as follows.

Theorem 3.1. *Assume that $E \geq E_0$ is an admissible number, and let $n_0 = n_0(E)$. Then for all $n \geq n_0 + 1$:*

1. *Each of the nodes u_n switches at a moment t_n satisfying*

$$t_n = an^2 + q_n, \quad |q_n| \leq E\sqrt{n}, \quad (3.3)$$

$$t_k < t_{n_0} < t_{n_0+1} < \dots, \quad k = 0, 1, \dots, n_0 - 1,$$

2. *There exists $A_\nabla > 0$ depending on h_1, c, E , but not on n , such that*

$$\left| \nabla u_n(t_n) + \frac{3h_1}{4} \right| \leq A_\nabla n^{-1/2}, \quad \nabla u_n(t_n) \leq -\frac{3h_1}{8}.$$

4 Auxiliary Statements

In this section, we formulate several auxiliary statements. Each of them is a key ingredient in the proof of our main result, i.e., Theorem 3.1.

In Section 4.1 (Proposition 4.1), we establish asymptotic formulas for the discrete Green function $y_n(t)$ given by (2.4). It is essential that the leading order terms in the asymptotics depend only on n/\sqrt{t} , while the remainders are estimated uniformly with respect to n .

In Section 4.2, we consider three expressions containing integrals (4.17) of leading order terms in the asymptotics of $y_n(t)$, $\nabla y_n(t)$, and $\dot{y}_n(t)$, respectively. These three expressions will enter the leading order terms in asymptotic formulas for $u_n(t_n)$, $\nabla u_n(t_n)$, and $\dot{u}_n(t_n)$. In Proposition 4.2, we show that these terms vanish for the same value of a , thus determining the “propagation rate” an^2 in the switching moment asymptotics for t_n in (3.1) and (3.3).

In Section 4.3, we elaborate on properties of integrals (4.17) from Section 4.2. In the proof of our main result, these integrals will play the role of approximation of some Riemann sums. Note that the corresponding integrands are not smooth functions, but have singularities of order $(1-x)^{1/2}$ or $(1-x)^{-1/2}$ at $x=1$. In Propositions 4.3 and 4.4, we provide error estimates for approximation of such integrals by their Riemann sums.

4.1 Properties of the discrete Green function $y_n(t)$

Consider the functions $h, f, g, \tilde{f} : \mathbb{R}^+ \rightarrow \mathbb{R}$ given by

$$h(x) := \frac{1}{2\sqrt{\pi}} e^{-x^2/4}, \quad f(x) := 2x \int_x^\infty y^{-2} h(y) dy, \quad g(x) := f'(x), \quad \tilde{f}(x) := -\frac{h'''(x)}{6x}. \quad (4.1)$$

Note that these functions belong to $C^\infty[0, \infty)$ and decay to zero as $x \rightarrow \infty$, together with all their derivatives, faster than exponentially. Moreover,

$$h(x) = g'(x) = f''(x), \quad 2h(x) + xg(x) - f(x) = 0. \quad (4.2)$$

Consider the functions $r_0, \tilde{r}_1, r_1, r_2, w_0, w_1 : (\mathbb{N} \cup \{0\}) \times \mathbb{R}_+ \rightarrow \mathbb{R}$ satisfying the following relations for $n = 0, 1, 2, \dots$ and $t > 0$:

$$y_n(t) = \sqrt{t} f\left(\frac{n}{\sqrt{t}}\right) + r_0(n, t), \quad (4.3)$$

$$y_n(t) = \sqrt{t} f\left(\frac{n}{\sqrt{t}}\right) + \frac{1}{\sqrt{t}} \tilde{f}\left(\frac{n}{\sqrt{t}}\right) + \tilde{r}_1(n, t), \quad (4.4)$$

$$\dot{y}_n(t) = \frac{1}{\sqrt{t}} h\left(\frac{n}{\sqrt{t}}\right) + r_1(n, t), \quad (4.5)$$

$$\ddot{y}_n(t) = \frac{1}{t\sqrt{t}} h''\left(\frac{n}{\sqrt{t}}\right) + r_2(n, t), \quad (4.6)$$

$$\nabla y_n(t) = g\left(\frac{n}{\sqrt{t}}\right) + \frac{1}{2\sqrt{t}} h\left(\frac{n}{\sqrt{t}}\right) + w_0(n, t), \quad (4.7)$$

$$\nabla \dot{y}_n(t) = \frac{1}{t} h'\left(\frac{n}{\sqrt{t}}\right) + w_1(n, t). \quad (4.8)$$

We fix throughout the paper

$$\tau_0 > 0. \quad (4.9)$$

The following estimates are proved in [10].

Proposition 4.1. *There exist constants $A_0, A_1, A_2, \tilde{A}_1, B_0, B_1, A_2^*, B_2^* > 0$ (depending on τ_0) such that, for all $t \geq \tau_0$, $n = 0, 1, 2, \dots$, and $i = 0, 1, 2$, the following inequalities hold:*

$$|r_i(n, t)| \leq A_i \frac{1}{t^i \sqrt{t}}, \quad |\tilde{r}_1(n, t)| \leq \tilde{A}_1 \frac{1}{t \sqrt{t}}, \quad (4.10)$$

$$|w_0(n, t)| \leq B_0 \frac{1}{t}, \quad |w_1(n, t)| \leq B_1 \frac{1}{t \sqrt{t}}, \quad (4.11)$$

$$|\ddot{y}_n(t)| \leq A_2^* \frac{1}{t \sqrt{t}}, \quad |\nabla \ddot{y}_n(t)| \leq B_2^* \frac{1}{t^2}. \quad (4.12)$$

4.2 Equivalence of some equations

Consider the functions

$$F(a, x) := \sqrt{a(1-x^2)} f\left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-x}{1+x}}\right), \quad (4.13)$$

$$G(a, x) := g\left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-x}{1+x}}\right), \quad (4.14)$$

$$H(a, x) := \frac{1}{\sqrt{a(1-x^2)}} h\left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-x}{1+x}}\right), \quad (4.15)$$

$$H_1(a, x) := \frac{1}{a(1-x^2)} h'\left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-x}{1+x}}\right), \quad (4.16)$$

where $a > 0$, $x \in (-1, 1)$ and f, g, h are given by (4.1). Set

$$I_F(a) := \int_{-1}^1 F(a, x) dx, \quad I_G(a) := \int_{-1}^1 G(a, x) dx, \quad I_H(a) := \int_{-1}^1 H(a, x) dx. \quad (4.17)$$

The following proposition is proved in Appendix B.

Proposition 4.2. *Each of the three equations*

$$-c + (h_1 - 2c)a - h_1 I_F(a) = 0, \quad (4.18)$$

$$-2c - h_1 I_G(a) = 0, \quad (4.19)$$

$$(h_1 - 2c) - h_1 I_H(a) = 0 \quad (4.20)$$

has a unique root on the interval $(0, \infty)$. Moreover, all these equations have the same root.

In what follows, we fix a given by Proposition 4.2 and write $F(x)$, $G(x)$, $H(x)$, $H_1(x)$, I_F , I_G , I_H , omitting the dependence on a .

4.3 Error estimates for Riemann sums

Let $N \in \mathbb{N}$, and let $z_0 = -1$ or $z_0 = 0$. The following propositions are proved in Appendix C.

Proposition 4.3. *Assume that a function $F_1(x)$ can be represented as*

$$F_1(x) = c_1(1-x)^{1/2} + c_2(1-x)^{3/2} + \tilde{F}_1(x),$$

where $c_1, c_2 \in \mathbb{R}$ and $\tilde{F}_1 \in C^2[z_0, 1]$. Denote by

$$R_n := \int_{z_0}^1 F_1(x) dx - \left(\frac{1}{2n} F_1(z_0) + \frac{1}{2n} F_1(1) + \sum_{k=z_0n+1}^{n-1} \frac{1}{n} F_1\left(\frac{k}{n}\right) \right)$$

the error estimate of the Riemann sum of the integral $\int_{z_0}^1 F_1(x) dx$.

1. There exists $L_1 = L_1(F_1, N) > 0$ such that

$$|R_n| \leq L_1 \frac{1}{n^{3/2}}, \quad n \geq N.$$

2. If, additionally, $\tilde{F}_1(x) = c_3(1-x)^{5/2} + c_4(1-x)^{7/2} + \bar{F}_1(x)$, where $c_3, c_4 \in \mathbb{R}$ and $\bar{F}_1 \in C^4[z_0, 1]$, then there exists $\bar{L}_1 = \bar{L}_1(F_1, N) > 0$ such that

$$|(n+1)^2 R_{n+1} - n^2 R_n| \leq \bar{L}_1 \frac{1}{n^{1/2}}.$$

Proposition 4.4. *Assume that a function $F_2(x)$ can be represented as*

$$F_2(x) = c_1(1-x)^{-1/2} + c_2(1-x)^{1/2} + \tilde{F}_2(x), \quad (4.21)$$

where $c_1 > 0$, $c_2 \in \mathbb{R}$, and $\tilde{F}_2 \in C^1[z_0, 1]$. Then there exists $L_2 = L_2(F_2, N) > 0$, $L_2^* = L_2^*(F_2, N) > c_1$ and $l_2 = l_2(F_2, N)$ such that

$$L_2^* \frac{1}{n^{1/2}} - l_2 \frac{1}{n} \leq \int_{z_0}^1 F_2(x) dx - \sum_{k=z_0n}^{n-1} \frac{1}{n} F_2\left(\frac{k}{n}\right) \leq L_2 \frac{1}{n^{1/2}}, \quad n \geq N. \quad (4.22)$$

In particular, $\left| \sum_{k=z_0n}^{n-1} \frac{1}{n} F_2\left(\frac{k}{n}\right) \right|$ is bounded.

The following lemma is straightforward.

Lemma 4.5. *Assume that $F_3 \in C^0[-1, 1)$ and $|(1-x)^{3/2} F_3(x)|$ is bounded on $[-1, 1)$. Then there exists $L_3 = L_3(F_3, N) > 0$ such that*

$$\left| \sum_{|k| \leq n-1} \frac{1}{n} F_3\left(\frac{k}{n}\right) \right| \leq L_3 n^{1/2}, \quad n \geq N.$$

5 Asymptotics for $u_n(t_n)$

5.1 Preliminaries

Set

$$h_a(x) := h\left(\frac{x}{\sqrt{a}}\right) + h\left(\frac{1}{x\sqrt{a}}\right), \quad D_a := \inf_{x \in (0,1]} h_a(x) > h(a^{-1/2}), \quad (5.1)$$

$$p := \sup_{x \in (0,1]} \frac{h'_a(x)x}{h_a(x)}, \quad \Phi_p(x) := \frac{h_a(x)}{x^p}, \quad x > 0, \quad (5.2)$$

where h is defined in (4.1).

Lemma 5.1. 1. $0 < p \leq 2e^{-1} < 1$.

2. The function $\Phi_p(x)$ is nonincreasing on $(0, 1)$.

Proof. 1. Let $\mu := 1/(4a)$. Then

$$\begin{aligned} 2\sqrt{\pi}h_a(x) &= e^{-\mu x^2} \left(1 + e^{\mu x^2 - \mu x^{-2}}\right) > e^{-\mu x^2}, \\ 2\sqrt{\pi}h'_a(x)x &= e^{-\mu x^2} 2\mu \left(-x^2 + x^{-2}e^{\mu x^2 - \mu x^{-2}}\right) \leq e^{-\mu x^2} 2\mu(x^{-2} - x^2)e^{-\mu(x^{-2} - x^2)} \leq 2e^{-1}e^{-\mu x^2}. \end{aligned}$$

Hence, $h'_a(x)x < 2e^{-1}h_a(x)$ and $p \leq 2e^{-1}$. The inequality $p > 0$ is obvious.

2. Relations (5.2) imply the following for $x \in (0, 1]$:

$$\Phi'_p(x) = \frac{h'_a(x)x^p - h_a(x)px^{p-1}}{x^{2p}} = \frac{1}{x^{p+1}}(h'_a(x)x - ph_a(x)) \leq 0.$$

Hence, Φ_p is nonincreasing on $(0, 1)$. □

For any $N \in \mathbb{N}$, set

$$D_{p1} := 2^{\frac{1-p}{2}} - 1, \quad D_{p2} := N \left(\left(1 + \frac{1}{N}\right)^{\frac{1+p}{2}} - 1 \right). \quad (5.3)$$

Lemma 5.2. For any $N \in \mathbb{N}$ we have

$$\left(1 + \frac{1}{k}\right)^{\frac{1-p}{2}} \geq 1 + D_{p1} \frac{1}{k}, \quad k \geq 1, \quad (5.4)$$

$$\left(1 + \frac{1}{n}\right)^{\frac{1+p}{2}} \geq 1 + D_{p2} \frac{1}{n}, \quad n \geq N, \quad (5.5)$$

$$D_{p1} < D_{p2}, \quad D_{p1} < \frac{1}{2}. \quad (5.6)$$

Proof. Fix $\alpha \in (0, 1)$. Note that

$$(1+x)^{1-\alpha} \leq 1+x(1-\alpha), \quad x \in (0, 1]. \quad (5.7)$$

Consider the function $P_\alpha(x) := \frac{(1+x)^\alpha - 1}{x}$. Inequality (5.7) implies that

$$P'_\alpha(x) = \frac{\alpha(1+x)^{\alpha-1}x - ((1+x)^\alpha - 1)}{x^2} \leq 0.$$

Hence, $P_\alpha(x)$ is nonincreasing and, for any $k_1 \geq k_2$, the following holds:

$$P_\alpha(1/k_1) \geq P_\alpha(1/k_2). \quad (5.8)$$

Inequalities (5.4) and (5.5) are straightforward consequences of (5.8) for $\alpha = (1-p)/2$ and $\alpha = (1+p)/2$, respectively. Obviously, $D_{p_2} \geq 2^{(1+p)/2} - 1 > D_{p_1}$ and $D_{p_1} < \sqrt{2} - 1 < 1/2$. \square

Set

$$\varkappa := D_{p_2} - D_{p_1} - 2(D_{p_1} + D_{p_2})D_{p_1}\frac{1}{N}, \quad (5.9)$$

where D_{p_1}, D_{p_2} are given by (5.3). Lemma 5.1 and relations (5.3) imply that $\varkappa > 0$ for all sufficiently large N . In what follows, we fix

$$N \in \mathbb{N} \quad \text{such that} \quad \varkappa > 0. \quad (5.10)$$

Assume that $n_0 = n_0(E)$ satisfies the following.

Requirement 1. $n_0 \geq N$.

We remind that the complete list of requirements determining $n_0(E)$ is given in Section D.4.

Set

$$C_n := -cn^2 + (h_1 - 2c)an^2 - h_1 \sum_{|k| \leq n-1} y_{n-k}(a(n^2 - k^2)). \quad (5.11)$$

In other words, the values C_n are obtained by formally substituting $t = an^2$, $t_k = ak^2$, and $S(t) = \{-(n-1), \dots, n-1\}$ in (2.10).

In Section 5.3 below, we will prove the following result.

Proposition 5.3. *There exist $K, K' > 0$ such that for $n \geq N$ the following inequalities hold:*

$$|C_{n+1} - C_n| \leq K \frac{1}{\sqrt{n}}, \quad |C_n| \leq K' \sqrt{n}. \quad (5.12)$$

Fix K and K' from Proposition 5.3 and set

$$E_0 := \frac{K + \varkappa K'}{(h_1 - 2c)\varkappa}. \quad (5.13)$$

Note that $E_0 > 0$ due to (5.10). For each $E > E_0$, set

$$a_n^{\min} := a - \frac{2E\sqrt{n}}{2n-1}, \quad a_n^{\max} := a + \frac{2E\sqrt{n}}{2n-1}. \quad (5.14)$$

We assume that $n_0 = n_0(E)$ satisfies the following requirement.

Requirement 2. $a_n^{\min} \geq \tau_0/(2n-1)$ for all $n \geq n_0$, where τ_0 was fixed in (4.9).

Note that Requirement 2 implies that

$$a(n^2 - k^2) - 2E\sqrt{n} \geq \tau_0, \quad n \geq n_0, \quad |k| \leq n-1. \quad (5.15)$$

Set

$$\delta_n := \frac{2A_1}{aD_a} \frac{1}{n} + \frac{2A_2^* E a^{1/2}}{D_a (a_n^{\min})^{3/2}} \frac{1}{\sqrt{n}}. \quad (5.16)$$

Obviously,

$$a_n^{\min} \rightarrow a, \quad \delta_n \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

and

$$a_{n+1}^{\min} > a_n^{\min}, \quad \delta_{n+1} < \delta_n, \quad n \geq n_0.$$

We assume that $n_0 = n_0(E)$ satisfies the following requirement.

Requirement 3. $\delta_n + \delta_{n+1} \leq 2D_{p1}$ for all $n \geq n_0$.

Below we will use the following constants S_α , T_α , and R . For $\alpha > 0$, let S_α be the smallest number satisfying the inequalities

$$\sum_{|k| \leq n-1} \frac{(n-k)^\alpha}{(n^2 - k^2)^{\alpha+1/2}} \leq S_\alpha \frac{1}{\sqrt{n}}, \quad n \geq N. \quad (5.17)$$

For $\alpha > 1$, let T_α be the smallest number satisfying the inequalities

$$\sum_{|k| \leq n-1} \frac{1}{(n^2 - k^2)^\alpha} \leq T_\alpha \frac{1}{n^\alpha}, \quad n \geq N. \quad (5.18)$$

Let R be the smallest number satisfying the inequalities

$$\sum_{|k| \leq n-1} \frac{1}{(n^2 - k^2)^{1/2}} \leq R (= \pi), \quad n \geq N. \quad (5.19)$$

5.2 Candidates for switching moments t_n

5.2.1 Formulation of a theorem

In this section, we will prove the following result.

Theorem 5.4. *Let the assumptions of Theorem 3.1 hold. Then there exists a sequence t_k , $k \in \mathbb{Z}$, such that $t_0, t_{\pm 1}, \dots, t_{\pm n_0}$ are given by (3.1),*

$$t_n = an^2 + q_n, \quad |q_n| \leq E\sqrt{n}, \quad t_{-n} = t_n \quad \text{for } n \geq n_0 + 1,$$

$$t_k < t_{n_0} < t_{n_0+1} < \dots, \quad k = 0, 1, \dots, n_0 - 1,$$

and the functions

$$v_n(t) := -cn^2 + (h_1 - 2c)t - h_1 \sum_{|k| \leq n-1} y_{n-k}(t - t_k), \quad n \geq n_0 + 1, \quad (5.20)$$

satisfy

$$v_n(t_n) = 0, \quad n \geq n_0 + 1. \quad (5.21)$$

Remark 5.5. The sequence t_k in Theorem 5.4 is a sequence of candidates for *switching* moments in the following sense. Assume that, for some $n \geq n_0 + 1$, we know the following (this is what we will in particular prove in Sections 6 and 7 below):

1. the nodes u_0, \dots, u_{n-1} switch at time moments t_0, \dots, t_{n-1} , respectively,
2. the nodes $u_n, u_{n+1}, u_{n+2}, \dots$ do not switch on the time interval $[0, t_n]$.

Then $v_n(t)$ coincides with the solution $u_n(t)$ of problem (2.1)–(2.3) on the time interval $[0, t_n]$ and equality (5.21) implies that t_n is the switching moment of $u_n(t)$.

5.2.2 Proof of Theorem 5.4

First, we substitute (5.20) into (5.21), replace t_k and t_n by $ak^2 + q_k$ and $an^2 + q_n$, respectively, and expand y_{n-k} into the Taylor series around $a(n^2 - k^2)$. This yields

$$\begin{aligned} & -cn^2 + (h_1 - 2c)(an^2 + q_n) - h_1 \sum_{|k| \leq n-1} y_{n-k}(a(n^2 - k^2)) \\ & - h_1 \sum_{|k| \leq n-1} \dot{y}_{n-k}(a(n^2 - k^2))(q_n - q_k) \\ & - h_1 \sum_{|k| \leq n-1} \frac{1}{2} \ddot{y}_{n-k}(a(n^2 - k^2) + \xi_{n,k})(q_n - q_k)^2 = 0, \quad n \geq n_0 + 1, \end{aligned} \quad (5.22)$$

where $|\xi_{n,k}| \leq |q_n - q_k|$. We introduce the notation

$$\alpha_{n,k} := \dot{y}_{n-k}(a(n^2 - k^2)), \quad \beta_{n,k}(q_n) := \frac{1}{2} \ddot{y}_{n-k}(a(n^2 - k^2) + \xi_{n,k})(q_n - q_k), \quad (5.23)$$

where we omit an explicit indication of the dependence of $\beta_{n,k}$ on q_k with $|k| \leq n - 1$. Further, set

$$J_{n,k}(q_n) := \alpha_{n,k} + \alpha_{n,-k} + \beta_{n,k}(q_n) + \beta_{n,-k}(q_n), \quad k = 1, 2, \dots, n-1, \quad (5.24)$$

$$J_{n,0}(q_n) := \alpha_{n,0} + \beta_{n,0}(q_n), \quad (5.25)$$

$$J_n(q_n) := \sum_{k=0}^{n-1} J_{n,k}, \quad D_n(q_n) := h_1 - 2c - h_1 J_n(q_n). \quad (5.26)$$

Using this notation and recalling the definition of the constants C_n in (5.11), we rewrite (5.22) as follows (it will also be convenient to replace n by $n + 1$):

$$C_{n+1} + D_{n+1}(q_{n+1})q_{n+1} + h_1 \sum_{k=0}^n J_{n+1,k}(q_{n+1})q_k = 0, \quad n \geq n_0. \quad (5.27)$$

Thus, it remains to find a sequence q_k , $k \in \mathbb{Z}$, such that $|q_k| \leq E\sqrt{n_0}$ for $k = 0, \pm 1, \dots, \pm n_0$, $|q_{n+1}| \leq E\sqrt{n+1}$, $q_{-(n+1)} = q_{n+1}$ for $n = n_0, n_0 + 1, \dots$, and the equalities (5.27) hold.

First, we note that $q_0, \dots, q_{\pm n_0}$ are already prescribed by the assumption of the theorem. Moreover, (5.27) holds with $n + 1$ replaced by n_0 :

$$C_{n_0} + D_{n_0}(q_{n_0})q_{n_0} + h_1 \sum_{k=0}^{n_0-1} J_{n_0,k}(q_{n_0})q_k = 0. \quad (5.28)$$

Indeed, Requirement 2 implies that $t_{n_0} > t_k$, $k = 0, \dots, n_0 - 1$. Therefore,

$$S(t) = \{-(n_0 - 1), \dots, n_0 - 1\}, \quad t \in [t_{n_0-1}, t_{n_0}),$$

in (2.10) and

$$u_{n_0}(t_{n_0}) = -cn_0^2 + (h_1 - 2c)t_{n_0} - h_1 \sum_{|k| \leq n_0-1} y_{n_0-k}(t_{n_0} - t_k). \quad (5.29)$$

Hence, (5.28) is obtained in the same way as (5.27) from (5.20) and (5.21).

Now we proceed by induction. Fix $n \geq n_0$. Suppose, we have constructed the desired sequence q_0, \dots, q_n . Let us find q_{n+1} satisfying $|q_{n+1}| \leq E\sqrt{n+1}$ and equation (5.27). We rewrite equation (5.27) in the form

$$q_{n+1} = \mathbf{F}(q_{n+1}), \quad \mathbf{F}(q_{n+1}) := -\frac{C_{n+1}}{D_{n+1}(q_{n+1})} - h_1 \sum_{k=0}^n \frac{J_{n+1,k}(q_{n+1})}{D_{n+1}(q_{n+1})} \cdot q_k. \quad (5.30)$$

To prove Theorem 5.4, it now suffices to show that if $q_k \in [-E\sqrt{n}, E\sqrt{n}]$ for $k = 0, \pm 1, \dots, \pm n$, then \mathbf{F} has a fixed point on the interval $[-E\sqrt{n+1}, E\sqrt{n+1}]$.

To do so, we need to show that \mathbf{F} maps the interval $[-E\sqrt{n+1}, E\sqrt{n+1}]$ into itself. Let us indicate the main difficulty on this way. We will see in Sections 5.3 and 5.4 that $C_n \sim \sqrt{n}$, $D_n \sim 1/\sqrt{n}$, and $J_{n+1,k}(q_{n+1}) \sim 1/\sqrt{n^2 - k^2}$, provided that $|q_{n+1}| \leq E\sqrt{n+1}$. Therefore, the straightforward attempt to estimate $|\mathbf{F}(q_{n+1})|$ would yield

$$|\mathbf{F}(q_{n+1})| \leq \left| \frac{C_{n+1}}{D_{n+1}(q_{n+1})} \right| + E\sqrt{n} \cdot h_1 \sum_{k=0}^n \left| \frac{J_{n+1,k}(q_{n+1})}{D_{n+1}(q_{n+1})} \right|, \quad (5.31)$$

and we would obtain nothing better than $|\mathbf{F}(q_{n+1})| \leq \text{const} \cdot n$.

To overcome this difficulty, we will use the following trick. Note that, by the induction hypothesis, (5.27) holds with $n + 1$ replaced by n . Therefore, we can multiply (5.27) by

$1 + \varkappa/n$ with an appropriate $\varkappa > 0$ and subtract (5.27) with $n + 1$ replaced by n . As a result, we will obtain the equation

$$q_{n+1} = \tilde{\mathbf{F}}(q_{n+1}), \quad \tilde{\mathbf{F}}(q_{n+1}) = -\frac{\tilde{C}_{n+1}}{\tilde{D}_{n+1}(q_{n+1})} - h_1 \sum_{k=0}^n \frac{\tilde{J}_{n+1,k}(q_{n+1})}{\tilde{D}_{n+1}(q_{n+1})} \cdot q_k, \quad (5.32)$$

which is equivalent to (5.30). The advantage of this new representation will be that we will obtain $\tilde{C}_n \sim 1/\sqrt{n}$ and $\tilde{D}_n \sim 1/\sqrt{n}$. Hence, the first term in the formula for $\tilde{\mathbf{F}}$ can be estimated by a constant $\alpha_1 > 0$. Furthermore, we will show that the expression $h_1 \sum_{k=0}^n \left| \frac{\tilde{J}_{n+1,k}(q_{n+1})}{\tilde{D}_{n+1}(q_{n+1})} \right|$ is estimated by $1 - \alpha_2/\sqrt{n}$ with $\alpha_2 > 0$. Therefore, (5.32) will yield

$$|\tilde{\mathbf{F}}(q_{n+1})| \leq \alpha_1 + E\sqrt{n} - E\alpha_2 < E\sqrt{n+1}, \quad (5.33)$$

provided that $E \geq E_0 = \alpha_1/\alpha_2$. In particular, it will turn out that the appropriate \varkappa is given by (5.9) and E_0 by (5.13). Interestingly, $\varkappa = 0$ would not be sufficient for this scheme as it would then follow that $\alpha_2 = 0$.

To make the above argument rigorous, we need the following proposition, in which we do not explicitly indicate the dependence of the functions on q_{n+1} .

Proposition 5.6. *Let the assumptions of Theorem 3.1 hold. Then for any $n \geq n_0$ and $q_0, q_{\pm 1}, \dots, q_{\pm(n+1)} \in [-E\sqrt{n+1}, E\sqrt{n+1}]$, the following holds with \varkappa given by (5.9):*

1. $J_{n,k} \geq 0, k = 0, 1, \dots, n-1,$
2. $J_{n,k} - \left(1 + \frac{\varkappa}{n}\right) J_{n+1,k} \geq 0, k = 0, 1, \dots, n-1,$
3. $D_n - h_1 \left(1 + \frac{\varkappa}{n}\right) J_{n+1,n} \geq 0.$

Now, assuming that Proposition 5.6 is true, we complete the proof of Theorem 5.4. After that, in Section 5.4, we prove Proposition 5.6.

Substituting n instead of $n + 1$ into (5.27) and using (5.28) for $n = n_0$ or the induction hypothesis for $n > n_0$ (and omitting the dependence of $J_{n,k}$ on q_n), we obtain the equation

$$C_n + D_n q_n + h_1 \sum_{k=0}^{n-1} J_{n,k} q_k = 0.$$

Multiplying (5.27) by $1 + \varkappa/n$ and subtracting the latter expression we have

$$\begin{aligned} 0 &= (C_{n+1} - C_n) + \frac{\varkappa}{n} C_{n+1} + \left(1 + \frac{\varkappa}{n}\right) D_{n+1} q_{n+1} \\ &\quad + \left(\left(1 + \frac{\varkappa}{n}\right) h_1 J_{n+1,n} - D_n \right) q_n + h_1 \sum_{k=0}^{n-1} \left(\left(1 + \frac{\varkappa}{n}\right) J_{n+1,k} - J_{n,k} \right) q_k. \end{aligned}$$

Equivalently (cf. (5.32)), $q_{n+1} = \tilde{\mathbf{F}}(q_{n+1})$, where $\tilde{\mathbf{F}}(q_{n+1})$ satisfies

$$\begin{aligned} \left(1 + \frac{\varkappa}{n}\right) D_{n+1} \tilde{\mathbf{F}}(q_{n+1}) &= -\left((C_{n+1} - C_n) + \frac{\varkappa}{n} C_{n+1}\right) \\ &\quad + \left(D_n - \left(1 + \frac{\varkappa}{n}\right) h_1 J_{n+1,n}\right) q_n + h_1 \sum_{k=0}^{n-1} \left(J_{n,k} - \left(1 + \frac{\varkappa}{n}\right) J_{n+1,k}\right) q_k. \end{aligned}$$

According to Proposition 5.6, all the coefficients at q_k , q_n , q_{n+1} are positive. The inductive hypothesis $|q_k| \leq E\sqrt{n}$, $k = 0, \dots, n$, Proposition 5.3, and the inequality $E \geq E_0$ imply that

$$\begin{aligned} \left(1 + \frac{\varkappa}{n}\right) D_{n+1} |\tilde{\mathbf{F}}(q_{n+1})| &\leq \left| (C_{n+1} - C_n) + \frac{\varkappa}{n} C_{n+1} \right| \\ &\quad + \left(D_n - \left(1 + \frac{\varkappa}{n}\right) h_1 J_{n+1,n} \right) E\sqrt{n} + h_1 \sum_{k=0}^{n-1} \left(J_{n,k} - \left(1 + \frac{\varkappa}{n}\right) J_{n+1,k} \right) E\sqrt{n} \\ &\leq \frac{K + \varkappa K'}{\sqrt{n}} + E\sqrt{n} \left(\left(D_n + h_1 \sum_{k=0}^{n-1} J_{n,k} \right) - \left(1 + \frac{\varkappa}{n}\right) h_1 \left(J_{n+1,n} + \sum_{k=0}^{n-1} J_{n+1,k} \right) \right). \end{aligned}$$

Combining this with (5.26) yields (cf. (5.33))

$$\begin{aligned} \left(1 + \frac{\varkappa}{n}\right) D_{n+1} |\tilde{\mathbf{F}}(q_{n+1})| &\leq (K + \varkappa K') \frac{1}{\sqrt{n}} + E\sqrt{n} \left((h_1 - 2c) - \left(1 + \frac{\varkappa}{n}\right) ((h_1 - 2c) - D_{n+1}) \right) \\ &= (K + \varkappa K') \frac{1}{\sqrt{n}} + E\sqrt{n} D_{n+1} \left(1 + \frac{\varkappa}{n}\right) - E\sqrt{n} (h_1 - 2c) \frac{\varkappa}{n}. \end{aligned}$$

The latter estimate and the inequality $E \geq E_0$, where E_0 is given by (5.13), imply

$$\left(1 + \frac{\varkappa}{n}\right) D_{n+1} |\tilde{\mathbf{F}}(q_{n+1})| \leq E\sqrt{n} D_{n+1} \left(1 + \frac{\varkappa}{n}\right).$$

Hence, $|\tilde{\mathbf{F}}(q_{n+1})| \leq E\sqrt{n} < E\sqrt{n+1}$. Therefore, $\tilde{\mathbf{F}}$ maps $[-E\sqrt{n+1}, E\sqrt{n+1}]$ into itself.

Furthermore, the function $\tilde{\mathbf{F}}$ is continuous, because the functions

$$\beta_{m,k}(q_m) = \begin{cases} \frac{y_{m-k}(a(m^2 - k^2) + q_m - q_k) - y_{m-k}(a(m^2 - k^2))}{q_m - q_k} - \alpha_{m,k}, & q_m \neq q_k, \\ 0, & q_m = q_k, \end{cases}$$

are continuous with respect to $q_m \in \mathbb{R}$ for $k = 0, \dots, m-1$ and $D_{n+1}(q_{n+1}) > 0$ by Proposition 5.6. Hence, by Brouwer's fixed-point theorem, the map $\tilde{\mathbf{F}}$ has a fixed point on the interval $[-E\sqrt{n+1}, E\sqrt{n+1}]$. The latter implies Theorem 5.4.

It remains to prove Propositions 5.3 and 5.6, which we do in the next two sections.

5.3 Proof of Proposition 5.3

Taking into account equalities (4.4) and (4.18) we write (5.11) as follows:

$$C_n = C_n^{(1)} + C_n^{(2)} + C_n^{(3)},$$

where

$$\begin{aligned}
C_n^{(1)} &:= h_1 \left(n^2 I_F - \sum_{|k| \leq n-1} \sqrt{a(n^2 - k^2)} f \left(\frac{n-k}{\sqrt{a(n^2 - k^2)}} \right) \right), \\
C_n^{(2)} &:= -h_1 \sum_{|k| \leq n-1} \frac{1}{\sqrt{a(n^2 - k^2)}} \tilde{f} \left(\frac{n-k}{\sqrt{a(n^2 - k^2)}} \right), \\
C_n^{(3)} &:= -h_1 \sum_{|k| \leq n-1} \tilde{r}_1(n-k, a(n^2 - k^2)).
\end{aligned} \tag{5.34}$$

Proof of inequality (5.12). Below we separately estimate $C_n^{(1)}$ and $C_n^{(2)} + C_n^{(3)}$.

Step 1. Applying Proposition 4.3 (item 1) to the function $F(x)$ (see (4.13)), we conclude that

$$|C_n^{(1)}| \leq h_1 L_1(F, N) \sqrt{n} = K'_1 \sqrt{n}, \quad K'_1 := h_1 L_1(F, N). \tag{5.35}$$

Step 2. Due to (4.3), (4.4), and (4.10),

$$|C_n^{(2)} + C_n^{(3)}| = \left| h_1 \sum_{|k| \leq n-1} r_0(n-k, a(n^2 - k^2)) \right| \leq h_1 A_0 \sum_{|k| \leq n-1} \frac{1}{\sqrt{a(n^2 - k^2)}} \leq l_1,$$

where, by (5.19),

$$l_1 := h_1 A_0 \frac{1}{a^{1/2}} R. \tag{5.36}$$

Taking

$$K' := K'_1 + l_1 \frac{1}{\sqrt{N}}, \tag{5.37}$$

we obtain inequality (5.12). \square

Proof of inequality (5.12). Below we separately estimate $C_{n+1}^{(1)} - C_n^{(1)}$, $C_{n+1}^{(2)} - C_n^{(2)}$, and $C_{n+1}^{(3)} - C_n^{(3)}$ in Steps 1, 2, and 3 respectively.

Step 1. Applying Proposition 4.3 (item 2) to the function $F(x)$, we conclude that

$$|C_{n+1}^{(1)} - C_n^{(1)}| \leq K_1 \frac{1}{\sqrt{n}}, \quad K_1 := h_1 \bar{L}_1(F, N). \tag{5.38}$$

Step 2. Set

$$\tilde{F}(x) := \frac{1}{\sqrt{a(1-x^2)}} \tilde{f} \left(\frac{1-x}{\sqrt{a(1-x^2)}} \right), \quad I_{\tilde{F}} := \int_{-1}^1 \tilde{F}(x) dx. \tag{5.39}$$

Note that $\lim_{x \rightarrow 1} \tilde{F}(x)(1-x)^{1/2} < 0$. Therefore, by Proposition 4.4 applied to the function $-\tilde{F}(x)$, for some constants $L_{\tilde{f}}, L_{\tilde{f}}^*, l_{\tilde{f}} \geq 0$, we have

$$L_{\tilde{f}}^* \frac{1}{\sqrt{n}} - l_{\tilde{f}} \frac{1}{n} \leq -I_{\tilde{F}} - \frac{1}{h_1} C_n^{(2)} \leq L_{\tilde{f}} \frac{1}{\sqrt{n}}. \tag{5.40}$$

Hence,

$$|C_{n+1}^{(2)} - C_n^{(2)}| \leq K_2 \frac{1}{\sqrt{n}}, \quad (5.41)$$

where

$$K_2 := h_1 \left(L_{\tilde{f}} - L_{\tilde{f}}^* + l_{\tilde{f}} \frac{1}{\sqrt{N}} \right). \quad (5.42)$$

Step 3. Inequalities (4.10) and (5.18) imply that

$$|C_n^{(3)}| \leq h_1 \tilde{A}_1 \sum_{|k| < n} \frac{1}{(a(n^2 - k^2))^{3/2}} \leq h_1 \tilde{A}_1 \frac{1}{a^{3/2}} T_{3/2} \frac{1}{n^{3/2}}.$$

Hence,

$$|C_{n+1}^{(3)} - C_n^{(3)}| \leq K_3 \frac{1}{n^{3/2}}, \quad K_3 := 2h_1 \tilde{A}_1 \frac{1}{a^{3/2}} T_{3/2} \quad (5.43)$$

Summarising (5.38), (5.41), and (5.43) yields

$$|C_{n+1} - C_n| \leq K \frac{1}{\sqrt{n}}, \quad K := K_1 + K_2 + K_3 \frac{1}{N}. \quad (5.44)$$

□

5.4 Proof of Proposition 5.6

5.4.1 Proof of Proposition 5.6: Preliminaries

For $k = 1, \dots, n-1$, consider the following representation of $J_{n,k}$ (see (5.24) and (4.5)):

$$J_{n,k} := J_{n,k}^{\text{main}} + (w_{n,k} + w_{n,-k}) + (\beta_{n,k} + \beta_{n,-k}),$$

where

$$J_{n,k}^{\text{main}} := \gamma_{n,k} + \gamma_{n,-k},$$

$$\gamma_{n,\pm k} := \frac{1}{\sqrt{a(n^2 - k^2)}} h \left(\sqrt{\frac{n \mp k}{a(n \pm k)}} \right), \quad w_{n,\pm k} := r_1(n \mp k, (a(n^2 - k^2))),$$

and

$$J_{n,0} = J_{n,0}^{\text{main}} + w_{n,0} + \beta_{n,0},$$

where

$$J_{n,0}^{\text{main}} := \gamma_{n,0} := \frac{1}{\sqrt{an^2}} h(1/\sqrt{a}), \quad w_{n,0} := r_1(n, an^2).$$

The general idea is to prove each assertion of Proposition 5.6 for $J_{n,k}^{\text{main}}$ first, and then consider $J_{n,k}$ as a small perturbation of $J_{n,k}^{\text{main}}$. We formulate the fact that it is a small perturbation as a lemma.

Lemma 5.7. *Let δ_n be given by (5.16). Then*

$$\frac{|J_{n,k} - J_{n,k}^{\text{main}}|}{J_{n,k}^{\text{main}}} \leq \delta_n \frac{n}{n+k} \frac{1}{n-k}, \quad k \geq 0. \quad (5.45)$$

Proof. Fix $k \in \{1, \dots, n-1\}$ (the case $k = 0$ can be treated similarly). It follows from (4.10) and (5.15) that

$$|w_{n,k} + w_{n,-k}| \leq 2A_1 \frac{1}{(a(n^2 - k^2))^{3/2}}. \quad (5.46)$$

Definition (5.23) of $\beta_{n,k}$, (5.15), (4.12) and (5.14) imply that

$$\begin{aligned} |\beta_{n,k} + \beta_{n,-k}| &\leq \max_{|\xi_{n,k}| \leq |q_n - q_k|} |\ddot{y}_{n-k}(a(n^2 - k^2) + \xi_{n,k})| \cdot |q_n - q_k| \\ &\leq \frac{2A_2^* E \sqrt{n}}{(a(n^2 - k^2) - 2E\sqrt{n})^{3/2}} \leq \frac{2A_2^* E \sqrt{n}}{(a_n^{\min}(n^2 - k^2))^{3/2}}. \end{aligned} \quad (5.47)$$

Note that, by (5.1),

$$J_{n,k}^{\text{main}} \geq \frac{D_a}{(a(n^2 - k^2))^{1/2}}. \quad (5.48)$$

Relations (5.46)–(5.48) imply

$$\begin{aligned} \frac{|w_{n,k} + w_{n,-k}|}{J_{n,k}^{\text{main}}} &\leq \frac{2A_1 \frac{1}{(a(n^2 - k^2))^{3/2}}}{\frac{D_a}{(a(n^2 - k^2))^{1/2}}} = \frac{2A_1}{aD_a} \frac{1}{n+k} \frac{1}{n-k}, \\ \frac{|\beta_{n,k} + \beta_{n,-k}|}{J_{n,k}^{\text{main}}} &\leq \frac{\frac{2A_2^* E \sqrt{n}}{(a_n^{\min}(n^2 - k^2))^{3/2}}}{\frac{D_a}{(a(n^2 - k^2))^{1/2}}} = \frac{2A_2^* E a^{1/2}}{D_a (a_n^{\min})^{3/2}} \frac{\sqrt{n}}{n+k} \frac{1}{n-k}. \end{aligned}$$

Hence, inequality (5.45) holds. \square

5.4.2 Proof of Proposition 5.6: Part 1

Since $J_{n,k}^{\text{main}} > 0$, it suffices to show that the right-hand side in (5.45) is less than or equal to 1. The latter is true because $\delta_n \leq 1$ due to (5.6) and Requirement 3.

5.4.3 Proof of Proposition 5.6: Part 2

Fix $k \in \{1, \dots, n-1\}$ (the case $k = 0$ can be treated similarly). Note that

$$\begin{aligned} J_{n,k}^{\text{main}} &= \frac{1}{\sqrt{a(n^2 - k^2)}} \left(\sqrt{\frac{n-k}{n+k}} \right)^p \Phi_p \left(\sqrt{\frac{n-k}{n+k}} \right) \\ &= \frac{1}{\sqrt{a(n-k)^{(1-p)/2} (n+k)^{(1+p)/2}}} \Phi_p \left(\sqrt{\frac{n-k}{n+k}} \right), \end{aligned}$$

where $\Phi_p(x)$, $x \in (0, 1]$, is given by (5.2). Taking into account that $\Phi_p(x)$ is nonincreasing (see Lemma 5.1, item 2) and using (5.4) and (5.5), we have

$$\begin{aligned} \frac{J_{n,k}^{\text{main}}}{J_{n+1,k}^{\text{main}}} &\geq \frac{(n+1-k)^{(1-p)/2}(n+1+k)^{(1+p)/2}}{(n-k)^{(1-p)/2}(n+k)^{(1+p)/2}} = \left(1 + \frac{1}{n-k}\right)^{(1-p)/2} \left(1 + \frac{1}{n+k}\right)^{(1+p)/2} \\ &\geq \left(1 + D_{p1}\frac{1}{n-k}\right) \left(1 + D_{p2}\frac{1}{n+k}\right) \geq 1 + D_{p1}\frac{1}{n-k} + D_{p2}\frac{1}{n+k}. \end{aligned} \quad (5.49)$$

Combining (5.49) with (5.45) yields

$$\begin{aligned} \frac{J_{n,k}}{J_{n+1,k}} &= \frac{J_{n,k}^{\text{main}}}{J_{n+1,k}^{\text{main}}} \frac{1 + \frac{J_{n,k} - J_{n,k}^{\text{main}}}{J_{n,k}^{\text{main}}}}{1 + \frac{J_{n+1,k} - J_{n+1,k}^{\text{main}}}{J_{n+1,k}^{\text{main}}}} \geq \frac{J_{n,k}^{\text{main}}}{J_{n+1,k}^{\text{main}}} \frac{1 - \frac{\delta_n}{n-k} \frac{n}{n+k}}{1 + \frac{\delta_{n+1}}{n+1-k} \frac{n+1}{n+1+k}} \\ &\geq \frac{J_{n,k}^{\text{main}}}{J_{n+1,k}^{\text{main}}} \left(1 - \frac{\delta_n}{n-k} \frac{n}{n+k}\right) \left(1 + \frac{\delta_{n+1}}{n-k} \frac{n}{n+k}\right)^{-1} \\ &\geq \left(1 + D_{p1}\frac{1}{n-k} + D_{p2}\frac{1}{n+k}\right) \left(1 - (\delta_n + \delta_{n+1})\frac{n}{n+k} \frac{1}{n-k}\right). \end{aligned}$$

It is easy to show that Requirement 3 implies that the minimum of the last expression is achieved for $k = 0$. Hence, using Requirement 3 again, we obtain

$$\begin{aligned} \frac{J_{n,k}}{J_{n+1,k}} &\geq \left(1 + (D_{p1} + D_{p2})\frac{1}{n}\right) \left(1 - (\delta_n + \delta_{n+1})\frac{1}{n}\right) \\ &\geq 1 + \left(D_{p1} + D_{p2} - (\delta_n + \delta_{n+1}) - (D_{p1} + D_{p2})(\delta_n + \delta_{n+1})\frac{1}{N}\right) \frac{1}{n} \\ &\geq 1 + \frac{\varkappa}{n}, \end{aligned}$$

where \varkappa is given by (5.9).

5.4.4 Proof of Proposition 5.6: Part 3

Using (5.26) and the notation in the beginning of Section 5.4.1, we write

$$D_n - h_1 J_{n+1,n} \left(1 + \frac{\varkappa}{n}\right) = (h_1 - 2c) - h_1 J_n - h_1 J_{n+1,n} \left(1 + \frac{\varkappa}{n}\right) = \Sigma_{1,n} - (\Sigma_{2,n} + \Sigma_{3,n} + \Sigma_{4,n}), \quad (5.50)$$

where

$$\begin{aligned}\Sigma_{1,n} &:= (h_1 - 2c) - h_1 \sum_{k=0}^{n-1} J_{n,k}^{\text{main}}, \\ \Sigma_{2,n} &:= h_1 \gamma_{n+1,n} \left(1 + \frac{\varkappa}{n}\right), \\ \Sigma_{3,n} &:= h_1 \sum_{k=0}^{n-1} (J_{n,k} - J_{n,k}^{\text{main}}), \\ \Sigma_{4,n} &:= h_1 \left((J_{n+1,n} - J_{n+1,n}^{\text{main}}) + \gamma_{n+1,-n} \right) \left(1 + \frac{\varkappa}{n}\right).\end{aligned}$$

In steps 1–4 below, we will estimate $\Sigma_{1,n}, \dots, \Sigma_{4,n}$. We will see that $\Sigma_{1,n}$ and $\Sigma_{2,n}$ are “large” with respect to $\Sigma_{3,n}$ and $\Sigma_{4,n}$, which motivates the splitting in (5.50).

Set

$$\tilde{H}_1(x) := H(x) + H(-x), \quad x \in [0, 1),$$

where $H(x)$ is given by (4.15). Then

$$\begin{aligned}J_{n,k}^{\text{main}} &= \frac{1}{n} \tilde{H}_1\left(\frac{k}{n}\right), \quad k = 1, \dots, n-1, \\ J_{n,0}^{\text{main}} &= \frac{1}{2n} \tilde{H}_1(0).\end{aligned}$$

Step 1. Applying Proposition 4.4 to the function $H(x)$ given by (4.15) and using (4.17), we conclude that

$$\Sigma_{1,n} \geq h_1 \left(C_H \frac{1}{\sqrt{n}} - l_H \frac{1}{n} \right),$$

where

$$C_H := L_2^*(H, N), \quad l_H := l_2(H, N). \quad (5.51)$$

Step 2. Since the function $h(x)$ is decreasing,

$$\Sigma_{2,n} \leq h_1 \frac{1}{\sqrt{2a}} h(0) \frac{1}{\sqrt{n}} \left(1 + \frac{\varkappa}{n}\right).$$

Step 3. Due to (5.45), we have

$$|\Sigma_{3,n}| \leq h_1 \delta_n \sum_{k=0}^{n-1} \frac{n}{n^2 - k^2} J_{n,k}^{\text{main}}.$$

Applying Lemma 4.5 to the function $\frac{1}{1-x^2} H(x)$, we conclude that

$$|\Sigma_{3,n}| \leq h_1 \delta_n C_{H2} \frac{1}{\sqrt{n}}, \quad C_{H2} := L_3 \left(\frac{1}{1-x^2} H \right). \quad (5.52)$$

Step 4. Using (5.45), we obtain

$$|J_{n+1,n} - J_{n+1,n}^{\text{main}}| \leq \delta_{n+1} \frac{n+1}{2n+1} J_{n+1,n}^{\text{main}} \leq \delta_{n+1} \frac{n+1}{2n+1} \left(\frac{1}{\sqrt{2a}} h(0) \frac{1}{\sqrt{n}} + \gamma_{n+1,-n} \right).$$

Note that

$$\gamma_{n+1,-n} \leq \frac{1}{\sqrt{2a}} h \left(\sqrt{\frac{2n+1}{a}} \right) \frac{1}{\sqrt{n}}.$$

Summarising the last two inequalities, we have

$$\Sigma_{4,n} \leq h_1 \frac{1}{\sqrt{2a}} \left(1 + \frac{\varkappa}{n} \right) \left(h(0) \frac{n+1}{2n+1} \delta_{n+1} + (1 + \delta_{n+1}) h \left(\sqrt{\frac{2n+1}{a}} \right) \right) \frac{1}{\sqrt{n}}.$$

Steps 1–4 yield Proposition 5.6 (part 2), provided that $n_0 = n_0(E)$ satisfies the following (we multiply the original expressions by \sqrt{n} and divide by h_1).

Requirement 4. For $n \geq n_0$, the following holds:

$$\begin{aligned} C_H - \frac{1}{\sqrt{2a}} h(0) &\geq l_H \frac{1}{\sqrt{n}} + \frac{\varkappa}{\sqrt{2a}} h(0) \frac{1}{n} + C_{H2} \delta_n \\ &\quad + \frac{1}{\sqrt{2a}} \left(1 + \frac{\varkappa}{n} \right) \left(h(0) \frac{n+1}{2n+1} \delta_{n+1} + (1 + \delta_{n+1}) h \left(\sqrt{\frac{2n+1}{a}} \right) \right). \end{aligned}$$

Note that, according to (5.51) and Proposition 4.4, the left-hand side of this expression is positive.

6 Asymptotics for $\nabla u_n(t_n)$

We consider the sequence t_k ($k \in \mathbb{Z}$) given by Theorem 5.4 and the quantities

$$\nabla v_n(t_n) = -2cn - c - h_1 \sum_{|k| \leq n-1} \nabla y_{n-k}(t_n - t_k), \quad n \geq n_0 + 1, \quad (6.1)$$

where $v_n(t)$ is given by (5.20).

Remark 6.1. Under the assumptions of Remark 5.5, we have

$$\nabla v_n(t_n) = \nabla u_n(t_n).$$

In this section, we will prove that $\nabla v_n(t_n) < 0$.

Theorem 6.2. *There exists $A_\nabla > 0$ depending on h_1, c, E such that, for all $n \geq n_0 + 1$, we have*

$$\left| \nabla v_n(t_n) + \frac{3h_1}{4} \right| \leq A_\nabla n^{-1/2}, \quad \nabla v_n(t_n) \leq -\frac{3h_1}{8}.$$

6.1 Preliminaries

Set

$$x_{n,k} := \frac{k}{n}. \quad (6.2)$$

Consider a constant K_{h_1} such that

$$\left| \sum_{|k| \leq n-1} \frac{1}{n} \frac{1}{a(1-x_{n,k}^2)} h' \left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-x_{n,k}}{1+x_{n,k}}} \right) \right| \leq K_{h_1}, \quad n \geq N. \quad (6.3)$$

Such a constant exists because the left-hand side in (6.3) is the Riemann sum of the finite integral (note that $h'(0) = 0$).

6.2 Leading order terms

Substituting t_k given by Theorem 5.4 into (6.1), we have

$$\nabla v_n(t_n) = -2cn - c - h_1 \sum_{|k| \leq n-1} \nabla y_{n-k}(a(n^2 - k^2) + q_n - q_k). \quad (6.4)$$

Due to the Taylor expansion,

$$\sum_{|k| \leq n-1} \nabla y_{n-k}(a(n^2 - k^2) + q_n - q_k) = \sum_{|k| \leq n-1} \nabla y_{n-k}(a(n^2 - k^2)) + \Sigma_{1,n}, \quad (6.5)$$

where

$$\Sigma_{1,n} := \sum_{|k| \leq n-1} \nabla \dot{y}_{n-k}(a(n^2 - k^2))(q_n - q_k) + \frac{1}{2} \sum_{|k| \leq n-1} \nabla \ddot{y}_{n-k}(a(n^2 - k^2) + \xi_{n,k})(q_n - q_k)^2 \quad (6.6)$$

with $|\xi_{n,k}| \leq |q_n - q_k| \leq 2En^{1/2}$.

Using (4.7) and the functions $G(x) = G(a, x)$ and $H(x) = H(a, x)$ given by (4.14) and (4.15), we represent the sum in (6.5) as follows:

$$\sum_{|k| \leq n-1} \nabla y_{n-k}(a(n^2 - k^2)) = n \sum_{|k| \leq n-1} \frac{1}{n} G(x_{n,k}) + \frac{1}{2} \sum_{|k| \leq n-1} \frac{1}{n} H(x_{n,k}) + \Sigma_{2,n}, \quad (6.7)$$

where

$$\Sigma_{2,n} := \sum_{|k| \leq n-1} w_0(n - k, a(n^2 - k^2)). \quad (6.8)$$

Now we replace the respective sums in (6.7) by the integrals. Set

$$\Sigma_{g,n} := \sum_{|k| \leq n-1} \frac{1}{n} G(x_{n,k}) - I_G + \frac{G(1)}{2n} = \sum_{|k| \leq n-1} \frac{1}{n} G(x_{n,k}) - I_G - \frac{1}{4n}, \quad (6.9)$$

$$\Sigma_{h,n} := \sum_{|k| \leq n-1} \frac{1}{n} H(x_{n,k}) - I_H, \quad (6.10)$$

where $I_G = I_G(a)$ and $I_H = I_H(a)$ are given by (4.17). Then (6.7) takes the form

$$\sum_{|k| \leq n-1} \nabla y_{n-k}(a(n^2 - k^2)) = I_G n + \frac{I_H}{2} + \frac{1}{4} + n\Sigma_{g,n} + \frac{\Sigma_{h,n}}{2} + \Sigma_{2,n}. \quad (6.11)$$

Combining (6.4), (6.5), and (6.11) and using Lemma 4.2, we obtain

$$\begin{aligned} \nabla v_n(t_n) &= (-2c - h_1 I_G)n + \left(-c - \frac{h_1 I_H}{2} - \frac{h_1}{4}\right) - h_1 \left(\Sigma_{1,n} + \Sigma_{2,n} + n\Sigma_{g,n} + \frac{\Sigma_{h,n}}{2}\right) \\ &= -\frac{3h_1}{4} - h_1 \left(\Sigma_{1,n} + \Sigma_{2,n} + n\Sigma_{g,n} + \frac{\Sigma_{h,n}}{2}\right). \end{aligned} \quad (6.12)$$

6.3 Remainders and proof of Theorem 6.2

It remains to estimate $\Sigma_{1,n}$, $\Sigma_{2,n}$, $\Sigma_{g,n}$, and $\Sigma_{h,n}$ in (6.12).

Lemma 6.3. $|\Sigma_{1,n}| \leq 2EK_{h_1}n^{-1/2} + \left(\frac{2EB_1T_{3/2}}{a^{3/2}} + 2E^2B_2^* \frac{T_2}{(a_n^{\min})^2}\right)n^{-1}$.

Proof. Using (6.6), we write $\Sigma_{1,n} = \Sigma'_{1,n} + \Sigma''_{1,n}$, where

$$\begin{aligned} \Sigma'_{1,n} &:= \sum_{|k| \leq n-1} \nabla \dot{y}_{n-k}(a(n^2 - k^2))(q_n - q_k), \\ \Sigma''_{1,n} &:= \frac{1}{2} \sum_{|k| \leq n-1} \nabla \dot{y}_{n-k}(a(n^2 - k^2) + \xi_{n,k})(q_n - q_k)^2. \end{aligned}$$

Using (4.8), (4.11), (6.3), (5.18), and the inequality $|q_n - q_k| \leq 2En^{1/2}$, we have

$$\begin{aligned} |\Sigma'_{1,n}| &\leq 2En^{-1/2} \left| \sum_{|k| \leq n-1} \frac{1}{n} \frac{1}{a(1 - x_{n,k}^2)} h' \left(\frac{1}{\sqrt{a}} \sqrt{\frac{1 - x_{n,k}}{1 + x_{n,k}}} \right) \right| + \sum_{|k| \leq n-1} \frac{2EB_1n^{1/2}}{(a(n^2 - k^2))^{3/2}} \\ &\leq 2EK_{h_1}n^{-1/2} + \frac{2EB_1T_{3/2}}{a^{3/2}}n^{-1}. \end{aligned}$$

Further, using (4.12) and the inequalities $|q_n - q_k| \leq 2En^{1/2}$ and $a(n^2 - k^2) + \xi_{n,k} \geq a_n^{\min}(n^2 - k^2)$, we have

$$|\Sigma''_{1,n}| \leq \frac{1}{2} \sum_{|k| \leq n-1} \frac{4E^2B_2^*n}{(a_n^{\min}(n^2 - k^2))^2} \leq \frac{2E^2B_2^*T_2}{(a_n^{\min})^2}n^{-1}. \quad (6.13)$$

□

Lemma 6.4. $|\Sigma_{2,n}| \leq \sum_{|k| \leq n-1} \frac{B_0}{a(n^2 - k^2)}$.

Proof. Lemma 6.4 is a straightforward consequence of (6.8) and (4.11). \square

Remark 6.5. For $n \geq 4$, we have

$$\begin{aligned} \sum_{|k| \leq n-1} \frac{1}{n^2 - k^2} &= n^{-2} + 2 \sum_{k=1}^{n-1} \frac{1}{(n-k)(n+k)} \\ &\leq n^{-2} + 2n^{-1} \sum_{k=1}^{n-1} \frac{1}{k} \leq n^{-1}(2 \ln(n-1) + 2 + n^{-1}) \rightarrow 0. \end{aligned}$$

Proposition 4.3 (item 1) implies the following.

Lemma 6.6. *There exists $K_g > 0$ such that for $n \geq N$ the following holds: $|\Sigma_{g,n}| \leq K_g n^{-3/2}$.*

Proposition 4.4 implies the following.

Lemma 6.7. *There exists $K_h > 0$ such that for $n \geq N$ the following holds: $|\Sigma_{h,n}| \leq K_h n^{-1/2}$.*

Now Theorem 6.2 follows from (6.12) and Lemmas 6.3–6.7, provided that the following requirement is satisfied.

Requirement 5. *For $n \geq n_0$, the following holds:*

$$2EK_{h1}n^{-1/2} + \left(\frac{2EB_1T_{3/2}}{a^{3/2}} + \frac{2E^2B_2^*T_2}{(a_n^{\min})^2} \right) n^{-1} + \sum_{|k| \leq n-1} \frac{B_0}{a(n^2 - k^2)} + \left(K_g + \frac{K_h}{2} \right) n^{-1/2} \leq \frac{3}{8}. \quad (6.14)$$

7 Estimates of $u_n(t), u_{n+1}(t), \dots$ for $t \in (t_{n-1}, t_n)$

7.1 Uniqueness of a switching moment

As before, we consider the sequence t_k and the functions v_n given by Theorem 5.4. In this section, we will prove the following result.

Theorem 7.1. *For all $n \geq n_0 + 1$, we have*

$$v_n(t) < 0, \quad t \in [t_{n-1}, t_n). \quad (7.1)$$

Fix $\theta_0 > 0$, satisfying

$$\theta_0 K_h < \frac{1}{4}, \quad (7.2)$$

where K_h is given by Lemma 6.7.

The proof of Theorem 7.1 is based on the following proposition.

Proposition 7.2. *Let the assumptions of Theorem 3.1 hold. Then, for all $n \geq n_0 + 1$, we have*

1. $\dot{v}_n(t) \leq \frac{3h_1 K_h}{2}(n-1)^{-1/2}$ for all $t \in [t_{n-1}, t_{n-1} + \theta_0(n-1)^{1/2}]$,
2. $\ddot{v}_n(t) \geq 0$ for all $t \in [t_{n-1} + \theta_0(n-1)^{1/2}, t_n]$.

We first assume that Proposition 7.2 is true and prove Theorem 7.1. The proof of Proposition 7.2 is given in Sections 7.2 and 7.3 below.

Proof of Theorem 7.1. By (3.2) (for $n = n_0 + 1$) and Theorem 6.2 (for $n \geq n_0 + 2$), we have $v_n(t_{n-1}) \leq -3h_1/8$. Therefore, Proposition 7.2 (part 1) and inequality (7.2) imply that, for $t \in [t_{n-1}, t_{n-1} + \theta_0(n-1)^{1/2}]$,

$$v_n(t) = v_n(t_{n-1}) + \int_{t_{n-1}}^t \dot{v}_n(s) ds \leq -\frac{3h_1}{8} + \theta_0(n-1)^{1/2} \frac{3h_1 K_h}{2}(n-1)^{-1/2} < 0. \quad (7.3)$$

Now, for $t \in [t_{n-1} + \theta_0(n-1)^{1/2}, t_n]$, Proposition 7.2 (part 2) guarantees that $v_n(t)$ can vanish no more than once, and thus, by Theorem 5.4, this happens no earlier than at $t = t_n$. \square

As a corollary of Theorem 7.1, we obtain the following result.

Theorem 7.3. *For all $n \geq n_0 + 1$, we have*

$$v_j(t) < 0, \quad t \in [t_{n-1}, t_n], \quad j \geq n.$$

Proof. First, we show that

$$\nabla v_j(t_n) < 0, \quad j \geq n. \quad (7.4)$$

To do so, we estimate $\Delta v_j(t)$ for $t \in (t_{n-1}, t_n]$ and $j \geq n$. Using (5.20), (2.5), and the fact that $\dot{y}_n(t) \geq 0$, we have

$$\Delta v_j(t) = -2c - h_1 \sum_{|k| \leq n-1} \Delta y_{j-k}(t - t_k) = -2c - h_1 \sum_{|k| \leq n-1} \dot{y}_{j-k}(t - t_k) \leq -2c.$$

In particular, $\Delta v_j(t_n) < 0$. Together with the relations $v_n(t_n) = 0$ (Theorem 5.4) and $v_{n+1}(t_n) < 0$ (Theorem 6.2), this yields (7.4).

On the other hand, (2.6) implies that $\nabla y_{j-k}(t - t_k)$ decreases and thus

$$\nabla v_j(t) = -c(2j+1) - h_1 \sum_{|k| \leq n-1} \nabla y_{j-k}(t - t_k)$$

increases. Together with (7.4), this yields $\nabla v_j(t) < 0$ for all $t \in (t_{n-1}, t_n]$ and $j \geq n$. Since, additionally, $v_n(t) < 0$ for all $t \in [t_{n-1}, t_n]$ (Theorem 7.1), the desired result follows. \square

Remark 7.4. Under the assumptions of Remark 5.5, Theorem 7.3 implies that t_n is indeed the switching moment of the node u_n .

Remark 7.5. Theorem 7.1 implies that the equation in (5.30) has a *unique* root on the interval $[-E\sqrt{n+1}, E\sqrt{n+1}]$

In the rest part of this section, we will prove Lemma 7.1

7.2 Proof of Proposition 7.2: Part 1

7.2.1 Leading order terms

We take $\theta \in [0, \theta_0(n-1)^{1/2}]$ and set $t = t_{n-1} + \theta \in [t_{n-1}, t_{n-1} + \theta_0(n-1)^{1/2}]$.

First, we represent $\dot{v}_n(t_{n-1} + \theta)$, using (5.20) and the relation $t_{-(n-1)} = t_{n-1}$, as follows:

$$\begin{aligned} \dot{v}_n(t_{n-1} + \theta) &= h_1 - 2c - h_1 \sum_{|k| \leq n-2} \dot{y}_{n-1-k}(t_{n-1} + \theta - t_k) \\ &\quad - h_1 \sum_{|k| \leq n-2} \nabla \dot{y}_{n-1-k}(t_{n-1} + \theta - t_k) - h_1(\dot{y}_1(\theta) + \dot{y}_{2n-1}(\theta)). \end{aligned} \quad (7.5)$$

Set

$$m := n - 1.$$

Since $\dot{y}_1(\theta) \geq 0$ and $\dot{y}_{2n-1}(\theta) \geq 0$ for all $\theta \geq 0$, we obtain

$$\dot{v}_n(t_m + \theta) \leq h_1 - 2c - h_1 \sum_{|k| \leq m-1} \dot{y}_{m-k}(t_m + \theta - t_k) - h_1 \Sigma_{3,m}, \quad (7.6)$$

where

$$\Sigma_{3,m} := \sum_{|k| \leq m-1} \nabla \dot{y}_{m-k}(t_m + \theta - t_k). \quad (7.7)$$

Further, to apply the Taylor expansion in (7.6), we note that

$$t_m - t_k + \theta = a(m^2 - k^2) + (q_m - q_k + \theta).$$

Therefore, using (4.5), the function $H(x) = H(a, x)$ given by (4.15), and (6.10), we obtain

$$\begin{aligned} \sum_{|k| \leq m-1} \dot{y}_{m-k}(t_m + \theta - t_k) &= \sum_{|k| \leq m-1} \dot{y}_{m-k}(a(m^2 - k^2)) + \Sigma_{4,m} \\ &= \sum_{|k| \leq m-1} \frac{1}{m} H(x_{m,k}) + \Sigma_{5,m} + \Sigma_{4,m} = I_H + \Sigma_{h,m} + \Sigma_{5,m} + \Sigma_{4,m}, \end{aligned} \quad (7.8)$$

where

$$\Sigma_{4,m} := \sum_{|k| \leq m-1} \ddot{y}_{m-k}(a(m^2 - k^2) + \xi_{m,k})(q_m - q_k + \theta), \quad (7.9)$$

$$\Sigma_{5,m} := \sum_{|k| \leq m-1} r_1(m - k, a(m^2 - k^2)), \quad (7.10)$$

$x_{m,k} = m/k$, and $-2Em^{1/2} \leq \xi_{m,k} \leq 2Em^{1/2} + \theta_0 m^{1/2}$.

Combining (7.6), (7.8), and Lemmas 4.2 and 6.7, we have

$$\dot{v}_n(t_m + \theta) \leq h_1 (K_h m^{-1/2} + |\Sigma_{3,m}| + |\Sigma_{4,m}| + |\Sigma_{5,m}|). \quad (7.11)$$

7.2.2 Remainders

It remains to estimate $\Sigma_{3,m}$, $\Sigma_{4,m}$, and $\Sigma_{5,m}$ in (7.11).

Lemma 7.6. $|\Sigma_{3,m}| \leq K_{h1}m^{-1} + \left(\frac{B_1 T_{3/2}}{a^{3/2}} + (2E + \theta_0)B_2^* \frac{T_2}{(a_m^{\min})^2} \right) m^{-3/2}$.

Proof. Using (7.7), we write $\Sigma_{3,m} = \Sigma'_{3,m} + \Sigma''_{3,m}$, where

$$\begin{aligned}\Sigma'_{3,m} &:= \sum_{|k| \leq m-1} \nabla \dot{y}_{m-k}(a(m^2 - k^2)), \\ \Sigma''_{3,m} &:= \sum_{|k| \leq m-1} \nabla \ddot{y}_{m-k}(a(m^2 - k^2) + \xi_{m,k})(q_m - q_k + \theta).\end{aligned}$$

Using (4.8), (4.11), (6.3), and (5.18), we have

$$\begin{aligned}|\Sigma'_{3,m}| &\leq m^{-1} \left| \sum_{|k| \leq m-1} \frac{1}{m} \frac{1}{a(1 - x_{m,k}^2)} h' \left(\frac{1}{\sqrt{a}} \sqrt{\frac{1 - x_{m,k}}{1 + x_{m,k}}} \right) \right| \\ &\quad + \sum_{|k| \leq m-1} \frac{B_1}{(a(m^2 - k^2))^{3/2}} \leq K_{h1}m^{-1} + B_1 \frac{T_{3/2}}{a^{3/2}} m^{-3/2}.\end{aligned}$$

Further, using (4.12), (5.18), and the inequalities $|q_m - q_k + \theta| \leq (2E + \theta_0)m^{1/2}$ and $a(m^2 - k^2) + \xi_{m,k} \geq a_m^{\min}(m^2 - k^2)$, we have

$$|\Sigma''_{3,m}| \leq \sum_{|k| \leq m-1} \frac{(2E + \theta_0)m^{1/2}B_2^*}{(a_m^{\min}(m^2 - k^2))^2} \leq (2E + \theta_0)B_2^* \frac{T_2}{(a_m^{\min})^2} m^{-3/2}.$$

□

Lemma 7.7. $|\Sigma_{4,m}| \leq \frac{(2E + \theta_0)A_2^*T_{3/2}}{(a_m^{\min})^{3/2}} m^{-1}$.

Proof. Using (7.9), (4.12), (5.18), and the inequalities $|q_m - q_k + \theta| \leq (2E + \theta_0)m^{1/2}$ and $a(m^2 - k^2) + \xi_{m,k} \geq a_m^{\min}(m^2 - k^2)$, we have

$$|\Sigma_{4,m}| \leq \sum_{|k| \leq m-1} \frac{(2E + \theta_0)A_2^*m^{1/2}}{(a_m^{\min}(m^2 - k^2))^{3/2}} \leq \frac{(2E + \theta_0)A_2^*T_{3/2}}{(a_m^{\min})^{3/2}} m^{-1}.$$

□

Lemma 7.8. $|\Sigma_{5,m}| \leq \frac{A_1 T_{3/2}}{a^{3/2}} m^{-3/2}$.

Proof. The assertion is a straightforward consequence of (7.10), (4.10), and (5.18),

□

Now inequality (7.11) together with Lemmas 7.6–7.8 yield part 1 in Proposition 7.2, provided that the following requirement is satisfied

Requirement 6. For $n \geq n_0$, the following holds:

$$\left(K_{h1} + \frac{(2E + \theta_0)A_2^*T_{3/2}}{(a_m^{\min})^{3/2}} \right) m^{-1/2} + \left(\frac{B_1T_{3/2}}{a^{3/2}} + \frac{(2E + \theta_0)B_2^*T_2}{(a_m^{\min})^2} + \frac{A_1T_{3/2}}{a^{3/2}} \right) m^{-1} \leq \frac{K_h}{2}. \quad (7.12)$$

7.3 Proof of Proposition 7.2: Part 2

7.3.1 Preliminaries

We introduce the function

$$\Psi(x) := h'' \left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-x}{1+x}} \right) + h'' \left(\frac{1}{\sqrt{a}} \sqrt{\frac{1+x}{1-x}} \right), \quad x \in (0, 1). \quad (7.13)$$

Note that

$$\lim_{x \rightarrow 1} \Psi(x) = h''(0) = -\frac{1}{4\sqrt{\pi}}. \quad (7.14)$$

Fix

$$\eta \in \left(0, \frac{1}{4\sqrt{\pi}} \right). \quad (7.15)$$

We will need the following lemma.

Lemma 7.9. There exist $x_0 \in (0, 1)$ and $\varepsilon_0 > 0$ such that

$$h'' \left(\frac{1-x}{\sqrt{a(1-x^2)} + \varepsilon_1} \right) + h'' \left(\frac{1+x}{\sqrt{a(1-x^2)} + \varepsilon_1} \right) \leq -\eta \quad (7.16)$$

for all $x \in [x_0, (m-1)/m]$ and ε_1 with

$$-\min \left(\varepsilon_0, \frac{a}{2m} \left(2 - \frac{1}{m} \right) \right) \leq \varepsilon_1 \leq \varepsilon_0. \quad (7.17)$$

Proof. Choose $\eta_1 \in (\eta, 1/(4\sqrt{\pi}))$. Equation (7.14) implies that there is $x_0 \in [0, 1)$ such that

$$\Psi(x) \leq -\eta_1, \quad x \in [x_0, 1). \quad (7.18)$$

It is not difficult to check that there exists $\varepsilon_0 > 0$ such that

$$\left| h'' \left(\frac{1-x}{\sqrt{a(1-x^2)} + \varepsilon} \right) - h'' \left(\frac{1-x}{\sqrt{a(1-x^2)}} \right) \right| \leq \frac{\eta_1 - \eta}{2} \quad (7.19)$$

for all $x \in (-1, 1)$ and ε satisfying $|\varepsilon| \leq \varepsilon_0$ and $\varepsilon \geq -a(1-x^2)/2$.

Formulas (7.18) and (7.19) imply Lemma 7.9. \square

We fix x_0 and ε_0 from Lemma 7.9 and b such that

$$b > 2a. \quad (7.20)$$

We introduce numbers R_1 and R_2 satisfying

$$\sum_{|k| < x_0 m} \frac{1}{(m^2 - k^2)^{3/2}} \leq R_1 m^{-2}, \quad m \geq N, \quad (7.21)$$

$$\sum_{k \in [x_0 m, m-1]} \frac{1}{(a_m^{\max}(m^2 - k^2) + bm)^{3/2}} \geq R_2 m^{-3/2}, \quad m \geq \max\left(N, \frac{1}{1 - x_0}\right). \quad (7.22)$$

Set

$$B_{h_2} := \sup_{x \in [0, \infty)} |h''(x)| = -h''(0), \quad B_{h_4} := \sup_{x \in [0, \infty)} |h''''(x)| = h''''(0). \quad (7.23)$$

7.3.2 Leading order terms

As before, we assume that $m = n - 1$. We take $\theta \in [\theta_0 m^{1/2}, bm]$ and set $t = t_m + \theta$. Then $t \in [t_m + \theta_0 m^{1/2}, t_m + bm]$, and the latter interval contains $[t_m + \theta_0 m^{1/2}, t_n]$, provided that the following is satisfied.

Requirement 7. For $n \geq n_0$, the following holds (see (7.20)):

$$a(n-1)^2 - E(n-1)^{1/2} + b(n-1) \geq an^2 + En^{1/2}. \quad (7.24)$$

First, we represent $\ddot{v}_n(t_{n-1} + \theta)$, using (5.20), as follows (cf. (7.5)):

$$\ddot{v}_n(t_m + \theta) = -h_1(I_{1,m} + I_{2,m} + \Sigma_{6,m}), \quad (7.25)$$

where

$$I_{1,m} := \sum_{|k| \leq m-1} \ddot{y}_{m-k}(t_m + \theta - t_k), \quad (7.26)$$

$$I_{2,m} := \ddot{y}_1(\theta) + \ddot{y}_{2m+1}(\theta), \quad (7.27)$$

$$\Sigma_{6,m} := \sum_{|k| \leq m-1} \nabla \ddot{y}_{m-k}(t_m + \theta - t_k). \quad (7.28)$$

Using (4.6), we represent $I_{1,m}$ as follows:

$$I_{1,m} = \sum_{|k| \leq m-1} \frac{1}{(a(m^2 - k^2) + q_m - q_k + \theta)^{3/2}} h'' \left(\frac{m-k}{\sqrt{a(m^2 - k^2) + q_m - q_k + \theta}} \right) + \Sigma_{7,m}, \quad (7.29)$$

where

$$\Sigma_{7,m} := \sum_{|k| \leq m-1} r_2(m-k, a(m^2 - k^2)) + q_m - q_k + \theta. \quad (7.30)$$

Now we split the sum in (7.29) into two sums in which the summation is taken over $|k| < x_0 m$ and $|k| \in [x_0 m, m-1]$, respectively. Let us estimate the first sum, using (7.21), (7.23), and the inequality $a(m^2 - k^2) + q_m - q_k \geq a_m^{\min}(m^2 - k^2)$:

$$\begin{aligned} & \left| \sum_{|k| < x_0 m} \frac{1}{(a(m^2 - k^2) + q_m - q_k + \theta)^{3/2}} h'' \left(\frac{m - k}{\sqrt{a(m^2 - k^2) + q_m - q_k + \theta}} \right) \right| \\ & \leq \sum_{|k| < x_0 m} \frac{B_{h_2}}{(a_m^{\min}(m^2 - k^2))^{3/2}} \leq \frac{B_{h_2} R_1}{(a_m^{\min})^{3/2}} m^{-2}. \end{aligned} \quad (7.31)$$

To estimate the second sum (which we have to do if $m - 1 \geq x_0 m$, i.e., $m \geq 1/(1 - x_0)$), we set

$$\varepsilon_{m,k} := \frac{q_m - q_k + \theta}{m^2}.$$

Below we assume that the following holds.

Requirement 8. For $n \geq \max\left(n_0(E), \frac{1}{1 - x_0}\right)$, the following holds:

$$- \min\left(\varepsilon_0, \frac{a}{2m} \left(2 - \frac{1}{m}\right)\right) \leq \frac{(\theta_0 - 2E)m^{1/2}}{m^2}.$$

Requirement 9. For $n \geq \max\left(n_0(E), \frac{1}{1 - x_0}\right)$, the following holds:

$$\frac{2Em^{1/2} + bm}{m^2} \leq \varepsilon_0.$$

Then

$$- \min\left(\varepsilon_0, \frac{a}{2m} \left(2 - \frac{1}{m}\right)\right) \leq \varepsilon_{m,k} \leq \varepsilon_0 \quad (7.32)$$

and

$$a(m^2 - k^2) + q_m - q_k \leq a_m^{\max}(m^2 - k^2). \quad (7.33)$$

Using (7.32) and (7.33) as well as (7.16) and (7.22), we have

$$\begin{aligned} & \sum_{|k| \in [x_0 m, m-1]} \frac{1}{(a(m^2 - k^2) + q_m - q_k + \theta)^{3/2}} h'' \left(\frac{m - k}{\sqrt{a(m^2 - k^2) + q_m - q_k + \theta}} \right) \\ & = \sum_{k \in [x_0 m, m-1]} \frac{1}{(a(m^2 - k^2) + q_m - q_k + \theta)^{3/2}} \\ & \quad \cdot \left[h'' \left(\frac{1 - x_{m,k}}{\sqrt{a(1 - x_{m,k}^2) + \varepsilon_{m,k}}} \right) + h'' \left(\frac{1 + x_{m,k}}{\sqrt{a(1 - x_{m,k}^2) + \varepsilon_{m,k}}} \right) \right] \\ & \leq -\eta \sum_{k \in [x_0 m, m-1]} \frac{1}{(a_m^{\max}(m^2 - k^2) + bm)^{3/2}} \leq -\eta R_2 m^{-3/2}, \end{aligned} \quad (7.34)$$

where $x_{m,k}$ is given by (6.2).

Thus, using (7.29), (7.31), and (7.34), we obtain the following estimate for $I_{1,m}$ in (7.26):

$$I_{1,m} \leq -\eta R_2 m^{-3/2} + \frac{B_{h2} R_1}{(a_m^{\min})^{3/2}} m^{-2} + \Sigma_{7,m}. \quad (7.35)$$

In what follows, we assume that the following is satisfied.

Requirement 10. For $n \geq n_0$, the following holds:

$$\tau_0 \leq \theta_0 n^{1/2}.$$

Now we represent $I_{2,m}$ in (7.27), using (4.6) and the equalities $h''(0) = -1/(4\sqrt{\pi})$ and $h'''(0) = 0$, as follows:

$$\begin{aligned} I_{2,m} &= \frac{1}{\theta^{3/2}} h'' \left(\frac{1}{\sqrt{\theta}} \right) + \frac{1}{\theta^{3/2}} h'' \left(\frac{2m+1}{\sqrt{\theta}} \right) + r_2(1, \theta) + r_2(2m+1, \theta) \\ &= \frac{1}{\theta^{3/2}} \left(-\frac{1}{4\sqrt{\pi}} + \frac{1}{2\theta} h''''(\xi) + h'' \left(\frac{2m+1}{\sqrt{\theta}} \right) + \theta^{3/2} (r_2(1, \theta) + r_2(2m+1, \theta)) \right), \end{aligned}$$

where $\xi \in [0, \theta^{-1/2}]$.

Below we assume that the following holds.

Requirement 11. For $n \geq n_0$ and $m = n - 1$, the following holds:

$$\frac{B_{h4}}{2\theta_0} m^{-1/2} + \sup_{x \geq \frac{2m+1}{\sqrt{bm}}} h''(x) + \frac{2A_2}{\theta_0^{1/2}} m^{-1/2} \leq \frac{1}{4\sqrt{\pi}}.$$

Hence,

$$I_{2,m} \leq 0. \quad (7.36)$$

Combining (7.25), (7.35), and (7.36), we see that, to prove Proposition 7.2 (part 2), it suffices to show that

$$-\eta R_2 m^{-3/2} + \frac{B_{h2} R_1}{(a_m^{\min})^{3/2}} m^{-2} + |\Sigma_{7,m}| + |\Sigma_{6,m}| \leq 0. \quad (7.37)$$

7.3.3 Remainders

Let us prove (7.37). To do so, we need to estimate $\Sigma_{6,m}$ and $\Sigma_{7,m}$.

Lemma 7.10. $|\Sigma_{6,m}| \leq \frac{B_2^* T_2}{(a_m^{\min})^2} m^{-2}$.

Proof. Using (7.28), (4.12), (5.18), and the inequality $a(m^2 - k^2) + q_m - q_k \geq a_m^{\min}(m^2 - k^2)$, we have

$$|\Sigma_{6,m}| \leq \sum_{|k| \leq m-1} \frac{B_2^*}{(a_m^{\min}(m^2 - k^2))^2} \leq \frac{B_2^* T_2}{(a_m^{\min})^2} m^{-2}.$$

Lemma 7.11. $|\Sigma_{7,m}| \leq \frac{A_2 T_{5/2}}{(a_m^{\min})^{5/2}} m^{-2}$.

Proof. Using (7.30), (4.10), (5.18), and the inequality $a(m^2 - k^2) + q_m - q_k \geq a_m^{\min}(m^2 - k^2)$, we have

$$|\Sigma_{7,m}| \leq \sum_{|k| \leq m-1} \frac{A_2}{(a_m^{\min}(m^2 - k^2) + \theta_0 m^{1/2})^{5/2}} \leq \frac{A_2 T_{5/2}}{(a_m^{\min})^{5/2}} m^{-5/2}.$$

Using Lemmas 7.10 and 7.11, we see that (7.37) holds, provided that the following is satisfied.

Requirement 12. For $n \geq n_0$ and $m = n - 1$, the following holds:

$$\left(\frac{B_{h_2} R_1}{(a_m^{\min})^{3/2}} + \frac{B_2^* T_2}{(a_m^{\min})^2} \right) m^{-1/2} + \frac{A_2 T_{5/2}}{(a_m^{\min})^{5/2}} m^{-1} \leq \eta R_2. \quad (7.38)$$

8 Main result: proof of Theorem 3.1

For $n = n_0 + 1$, Theorems 5.4 and 7.3 imply that the node $u_n(t)$ achieves the threshold 0 at a time moment $t_n = an^2 + q_n$, where $q_n \in [-E\sqrt{n}, 0]$ with the same E as in (3.1). Moreover, neither of the nodes u_n, u_{n+1}, \dots switches on the interval $[t_{n-1}, t_n)$, and thus $u_n(t)$ switches exactly at the moment t_n . Furthermore, by Theorem 6.2, the required estimates for $\nabla u_n(t_n)$ hold. Thus, the assertion of Theorem 3.1 holds for $n = n_0 + 1$.

In particular, we see that items 1 and 2 in Definition 3.1 hold with n_0 replaced by $n_0 + 1$ and with the same E as before. Hence, we can repeat the above argument to obtain the assertion of Theorem 3.1 for $n = n_0 + 2$, and so on, by induction, for any $n \geq n_0 + 1$.

A Existence and uniqueness: proof of Theorem 2.3

Step 1. Using (2.5), we see that the functions

$$z_n^{(1)}(t) := -cn^2 + (h_1 - 2c)t - h_1 y_n(t), \quad n \in \mathbb{Z}, \quad (A.1)$$

satisfy the initial condition (2.2) and the equation in (2.1) as long as $z_n^{(1)}(t) < 0$ for all $n \in \mathbb{Z} \setminus \{0\}$. By comparing $z_n^{(1)}(t)$ with the solution

$$z_n^+(t) = -cn^2 + (h_1 - 2c)t \quad (A.2)$$

of problem (2.1)–(2.3) with $\mathcal{H}(u_n)$ replaced by h_1 for all $n \in \mathbb{Z}$, it is not difficult to see that

$$z_n^{(1)}(t) \leq z_n^+(t), \quad t \geq 0, \quad n \in \mathbb{Z}. \quad (A.3)$$

Therefore, the time moment t_n at which $z_n(t)$ vanishes for the first time is not less than the moment $t_n^+ = cn^2/(h_1 - 2c)$ at which $z_n^+(t)$ vanishes.

In particular, $z_n^{(1)}(t) < 0$ for all $n \in \mathbb{Z} \setminus \{0\}$ and $t \in [0, c/(h_1 - 2c))$. Let $\tau_1 := \sup\{t > 0 : z_n^{(1)}(t) < 0 \forall n \in \mathbb{Z} \setminus \{0\}\}$. It follows from (A.1), (4.3), and (4.10) that τ_1 is finite. As we have seen, $\tau_1 \geq c/(h_1 - 2c) > 0$. Furthermore, $z_n^{(1)}(t)$ satisfy the growth condition from item 2 of Definition 2.1 for $t \in [0, \tau_1]$. This follows from (A.1) and the fact that $|y_n(t)|$ given by (2.4) are bounded on any finite time interval, uniformly with respect to $n \in \mathbb{Z}$. Thus, $u_n(t) := z_n^{(1)}(t)$ is a solution of problem (2.1)–(2.3) on the time interval $(0, \tau_1)$.

Let us prove that the solution $u_n(t)$ is unique on $(0, \tau_1)$. Assume we have another solution $\tilde{u}_n(t)$ on a time interval $(0, \tilde{\tau}_1)$, where $\tilde{\tau}_1 \leq \tau_1$ is such that $\tilde{u}_n(t) < 0$ for all $t \in (0, \tilde{\tau}_1)$ and $n \in \mathbb{Z} \setminus \{0\}$. Then the difference $w_n(t) := u_n(t) - \tilde{u}_n(t)$ must satisfy the homogeneous diffusion equation on the time interval $(0, \tilde{\tau}_1)$ with the zero initial data

$$\begin{aligned} \dot{w}_n(t) &= \Delta w_n(t), \quad t \in (0, \tilde{\tau}_1), \quad n \in \mathbb{Z}, \\ w_n(0) &= 0, \quad n \in \mathbb{Z}. \end{aligned}$$

If we looked for solutions that are square summable with respect to $n \in \mathbb{Z}$, then the application of the discrete Fourier transform would immediately imply that all $w_n(t) \equiv 0$. However, we are interested in solutions that may have exponential growth with respect to $n \in \mathbb{N}$ (see item 2 in Definition 2.1). Therefore, we will argue as follows. For each $N \in \mathbb{N}$, we introduce the functions

$$\zeta_n(t) = \zeta_n^N(t) := w_n(t) \text{ for } |n| \leq N, \quad \zeta_n(t) = \zeta_n^N(t) := 0 \text{ for } |n| \geq N + 1. \quad (\text{A.4})$$

They satisfy the relations

$$\begin{aligned} \dot{\zeta}_n(t) &= \Delta \zeta_n(t) + G_n^N(t), \quad t \in (0, \tilde{\tau}_1), \quad n \in \mathbb{Z}, \\ \zeta_n(0) &= 0, \quad n \in \mathbb{Z}, \end{aligned} \quad (\text{A.5})$$

where $G_n^N(t) = 0$ for $|n| \leq N - 1$ and $|n| \geq N + 2$, $G_{\pm N}^N(t) = w_{\pm(N+1)}(t)$, and $G_{\pm(N+1)}^N(t) = -w_{\pm N}(t)$. Since no more than finitely many elements in the sequences $\{\zeta_n(t)\}_{n \in \mathbb{Z}}$ and $\{G_n^N(t)\}_{n \in \mathbb{Z}}$ are nonzero, we can apply the discrete Fourier transform to (A.5) and obtain

$$\zeta_n^N(t) = \sum_{|k|=N}^{N+1} \int_0^t \dot{y}_{n-k}(t-s) G_k^N(s) ds, \quad t \in [0, \tilde{\tau}_1], \quad n \in \mathbb{Z}. \quad (\text{A.6})$$

Now let us fix $n \geq 0$ and $t \in [0, \tilde{\tau}_1]$. By assumption, there exist $A, \alpha \geq 0$ such that

$$\sup_{s \in [0, t]} |w_k(s)| \leq A e^{\alpha|k|}, \quad k \in \mathbb{Z}. \quad (\text{A.7})$$

Combining (A.4), (A.6), (2.7), (A.7) and choosing $N \geq n$, we have

$$|w_n(t)| = |\zeta_n^N(t)| \leq t \sum_{|k|=N}^{N+1} \sup_{s \in [0, t]} (|\dot{y}_{n-k}(s)| \cdot |G_k^N(s)|) \leq c \frac{t^N e^{\alpha N}}{(N-n)!} \rightarrow 0 \quad \text{as } N \rightarrow \infty,$$

where $c = c(n, t) \geq 0$ does not depend on N . Therefore, $w_{-n}(t) \equiv w_n(t) \equiv 0$. This proves that $u_n(t)$ is a unique solution of problem (2.1)–(2.3) on the time interval $(0, \tau_1)$.

Step 2. Set $S(\tau_1) := \{0\} \cup \{n \in \mathbb{Z} : z_n^{(1)}(\tau_1) = 0\}$, cf. (2.9). Due to (A.2) and (A.3), the set $S(\tau_1)$ is finite. Due to (A.1) and the symmetry $y_n(t) \equiv y_{-n}(t)$, the set $S(\tau_1)$ is symmetric with respect to the origin. Note that $t_0 = 0$ is the switching moment of the node 0, while $t_n = \tau_1$ are the switching moments of the nodes $n \in S(\tau_1) \setminus \{0\}$.

Using (2.5) and assuming $y_n(t) := 0$ for $t < 0$, we see that the functions

$$\begin{aligned} z_n^{(2)}(t) &= -cn^2 + (h_1 - 2c)t - h_1 \left(y_n(t) + \sum_{k \in S(\tau_1) \setminus \{0\}} y_{n-k}(t - \tau_1) \right) \\ &= -cn^2 + (h_1 - 2c)t - h_1 \sum_{k \in S(\tau_1)} y_{n-k}(t - t_k), \quad t \in [0, \infty), \end{aligned} \quad (\text{A.8})$$

satisfy the equations in (2.1) as long as $z_n^{(2)}(t) < 0$ for all $n \in \mathbb{Z} \setminus S(\tau_1)$, i.e., as long as $S(t) = S(\tau_1)$. Obviously, $z_n^{(2)}(t)$ also satisfy the initial condition (2.2).

As in Step 1, we see that the time moment t_n at which $z_n(t)$, $n \in \mathbb{Z} \setminus S(\tau_1)$, vanishes for the first time is not less than $cn^2/(h_1 - 2c)$. Hence, there is a positive time interval (of length bigger than τ_1) on which $z_n^{(2)}(t) < 0$ for all $n \in \mathbb{Z} \setminus S(\tau_1)$.

Let $\tau_2 := \sup\{t > 0 : z_n^{(2)}(t) < 0 \forall n \in \mathbb{Z} \setminus S(\tau_1)\}$. It follows from (A.1), (4.3), and (4.10) that τ_2 is finite. We have proved that $\tau_2 > \tau_1$. Furthermore, $z_n^{(2)}(t)$ satisfy the growth condition from item 2 of Definition 2.1 for $t \in [0, \tau_2]$. Thus, $u_n(t) := z_n^{(2)}(t)$ is a solution of problem (2.1)–(2.3) on the time interval $(0, \tau_2)$. Note that $u_n(t) = z_n^{(1)}(t)$ for $t \in [0, \tau_1]$. The uniqueness of $u_n(t)$ on the interval (τ_1, τ_2) can be proved similarly to Step 1.

Continuing these steps, we obtain the desired infinite sequence $\{\tau_j\}_{j \geq 0}$ (of switching moments) from Definition 2.1. On each step, we compare $u_n(t)$ with $z_n^+(t)$ given by (A.2) and conclude that the switching moments satisfy $t_n \geq cn^2/(h_1 - 2c)$. Hence, $\tau_j \rightarrow \infty$ as $j \rightarrow \infty$.

B Equivalence of three equations: proof of Proposition 4.2

We have to prove that

1. equation (4.20) has a unique root,
2. equations (4.18) and (4.20) have the same roots,
3. equations (4.19) and (4.20) have the same roots.

We will prove in detail items 1 and 2. Item 3 can be proved similarly to item 2.

Making the change of variables $y = \frac{1}{\sqrt{a}}\sqrt{\frac{1-x}{1+x}}$ in (4.15) and (4.13), we have

$$I_H(a) = \int_0^{+\infty} \frac{2}{1+ay^2} h(y) dy, \quad I_F(a) = \int_0^{+\infty} \frac{8a^2 y^2}{(1+ay^2)^3} f(y) dy.$$

Now we see that $I_H(a)$ decreases from 1 to 0 as a increases from 0 to $+\infty$. Hence, for any $0 < c < h_1/2$, equation (4.20) has a unique root $a > 0$. Item 1 is proved.

Let us prove item 2. Integrating by parts and using equations (4.1) and (4.2), we calculate

$$\begin{aligned} I_F(a) &= \int_0^{+\infty} \left(\frac{8a^2 y^2}{(1+ay^2)^3} - \frac{a}{1+ay^2} \right) f(y) dy + \int_0^{+\infty} \frac{a}{1+ay^2} f(y) dy \\ &= \frac{ay(-1+ay^2)}{(1+ay^2)^2} g(y) \Big|_0^{+\infty} - \int_0^{+\infty} \frac{ay(-1+ay^2)}{(1+ay^2)^2} g(y) dy + \int_0^{+\infty} \frac{a}{1+ay^2} f(y) dy \\ &= \int_0^{+\infty} \left(-\frac{ay(-1+ay^2)}{(1+ay^2)^2} + y \frac{a}{1+ay^2} \right) g(y) dy + \int_0^{+\infty} \frac{a}{1+ay^2} (f(y) - yg(y)) dy \\ &= -\frac{1}{1+ay^2} g(y) \Big|_0^{+\infty} + \int_0^{+\infty} \frac{1}{1+ay^2} h(y) dy + \int_0^{+\infty} \frac{2a}{1+ay^2} h(y) dy \\ &= -\frac{1}{2} + \int_0^{+\infty} \frac{2a+1}{1+ay^2} h(y) dy = \frac{2a+1}{2} I_H(a) - \frac{1}{2}. \end{aligned}$$

It is easy to conclude from the last identity that equations (4.18) and (4.20) have the same roots. Item 2 is proved.

C Error estimates for Riemann sums: proof of Propositions 4.3 and 4.4

Here we provide the proofs of item 2 of Proposition 4.3 and Proposition 4.4 for the case $z_0 = -1$. Item 1 of Proposition 4.3 and the case $z_0 = 0$ can be proved similarly.

C.1 Proof of item 2 of Proposition 4.3

Consider the function $\Phi(x) := F_1(x) + F_1(-x)$. Then

$$n^2 R_n = n^2 I_{F_1} - \left(\frac{n}{2} \Phi(0) + \frac{n}{2} \Phi(1) + \sum_{k=1}^{n-1} n \Phi\left(\frac{k}{n}\right) \right),$$

where $I_{F_1} := \int_{-1}^1 F_1(x) dx$. Therefore,

$$\begin{aligned} &(n+1)^2 R_{n+1} - n^2 R_n \\ &= (2n+1) I_{F_1} - \left(\frac{1}{2} \Phi(0) + \frac{1}{2} \Phi(1) + \sum_{k=1}^n (n+1) \Phi\left(\frac{k}{n+1}\right) - \sum_{k=1}^{n-1} n \Phi\left(\frac{k}{n}\right) \right) \\ &= \Sigma_{1,n} + \Sigma_{2,n}, \end{aligned} \tag{C.1}$$

where

$$\begin{aligned}\Sigma_{1,n} &:= (n+1)I_{F_1} - \left(\frac{1}{2}\Phi(0) + \frac{1}{2}\Phi(1) + \sum_{k=1}^n \Phi\left(\frac{k}{n+1}\right) \right), \\ \Sigma_{2,n} &:= n(I_{F_1} - \Phi(0) - \Sigma_{\Phi,n}), \\ \Sigma_{\Phi,n} &:= \sum_{k=1}^n \left(\Phi\left(\frac{k}{n+1}\right) - \Phi\left(\frac{k-1}{n}\right) \right).\end{aligned}$$

Note that $\Sigma_{1,n} = (n+1)R_{n+1}$. Hence, Proposition 4.3 (item 1) implies that, for some $L_1 > 0$, the following holds:

$$|\Sigma_{1,n}| \leq L_1 \frac{1}{\sqrt{n+1}}. \quad (\text{C.2})$$

Estimates of $\Sigma_{2,n}$ are more subtle. First, note that $\frac{k}{n+1} - \frac{k-1}{n} = \frac{n+1-k}{n(n+1)}$ and, hence,

$$\begin{aligned}\Phi\left(\frac{k}{n+1}\right) - \Phi\left(\frac{k-1}{n}\right) &= \frac{n+1-k}{n(n+1)}\Phi'\left(\frac{k}{n+1}\right) \\ &\quad - \frac{1}{2}\left(\frac{n+1-k}{n(n+1)}\right)^2\Phi''\left(\frac{k}{n+1}\right) + \frac{1}{6}\left(\frac{n+1-k}{n(n+1)}\right)^3\Phi'''(\xi_k),\end{aligned}$$

where $\xi_k \in \left[\frac{k-1}{n}, \frac{k}{n+1}\right]$. Therefore,

$$\begin{aligned}\Sigma_{\Phi,n} &= \frac{n+1}{n} \sum_{k=1}^n \frac{1}{n+1} \frac{n+1-k}{n+1} \Phi'\left(\frac{k}{n+1}\right) \\ &\quad - \frac{1}{2} \frac{n+1}{n^2} \sum_{k=1}^n \frac{1}{n+1} \left(\frac{n+1-k}{n+1}\right)^2 \Phi''\left(\frac{k}{n+1}\right) \\ &\quad + \frac{1}{6} \frac{n+1}{n^3} \sum_{k=1}^n \frac{1}{n+1} \left(\frac{n+1-k}{n+1}\right)^3 \Phi'''(\xi_k).\end{aligned} \quad (\text{C.3})$$

Set

$$\begin{aligned}\Phi_1(x) &:= (1-x)\Phi'(x), \quad \Phi_2(x) := (1-x)^2\Phi''(x), \quad \Phi_3(x) := (1-x)^3\Phi'''(x), \\ I_{\Phi_1} &:= \int_0^1 \Phi_1(x)dx, \quad I_{\Phi_2} := \int_0^1 \Phi_2(x)dx,\end{aligned} \quad (\text{C.4})$$

$$\Sigma_{\Phi_1,n} := I_{\Phi_1} - \left(\sum_{k=1}^n \frac{1}{n+1} \frac{n+1-k}{n+1} \Phi'\left(\frac{k}{n+1}\right) + \frac{1}{2(n+1)}\Phi'(0) \right), \quad (\text{C.5})$$

$$\Sigma_{\Phi_2,n} := I_{\Phi_2} - \left(\sum_{k=1}^n \frac{1}{n+1} \left(\frac{n+1-k}{n+1}\right)^2 \Phi''\left(\frac{k}{n+1}\right) + \frac{1}{2(n+1)}\Phi''(0) \right), \quad (\text{C.6})$$

$$\Sigma_{\Phi_3,n} := \sum_{k=1}^n \frac{1}{n+1} \left(\frac{n+1-k}{n+1}\right)^3 \Phi'''(\xi_k). \quad (\text{C.7})$$

Since $\xi_k \leq \frac{k}{n+1}$, we have

$$\left| \left(\frac{n+1-k}{n+1} \right)^3 \Phi'''(\xi_k) \right| \leq |\Phi_3(\xi_k)|. \quad (\text{C.8})$$

Proposition 4.3 (item 1) and (C.8) imply that, for

$$K_{\Phi_1} := L_1(\Phi_1, N), \quad K_{\Phi_2} := L_1(\Phi_2, N), \quad K_{\Phi_3} := \sup_{x \in (0,1)} |\Phi_3(x)|,$$

we have

$$|\Sigma_{\Phi_1, n}| \leq K_{\Phi_1} \frac{1}{(n+1)^{3/2}}, \quad |\Sigma_{\Phi_2, n}| \leq K_{\Phi_2} \frac{1}{(n+1)^{3/2}}, \quad |\Sigma_{\Phi_3, n}| \leq K_{\Phi_3}. \quad (\text{C.9})$$

In this notation, (C.3) takes the form

$$\begin{aligned} \Sigma_{\Phi, n} &= \frac{n+1}{n} \left(I_{\Phi_1} - \frac{1}{2(n+1)} \Phi'(0) - \Sigma_{\Phi_1, n} \right) \\ &\quad - \frac{1}{2} \frac{n+1}{n^2} \left(I_{\Phi_2} - \frac{1}{2(n+1)} \Phi''(0) - \Sigma_{\Phi_2, n} \right) + \frac{1}{6} \frac{n+1}{n^3} \Sigma_{\Phi_3, n}. \end{aligned}$$

Finally, taking into account (C.2), we conclude that

$$\begin{aligned} \Sigma_{2, n} &= nI_{F_1} - (n+1)I_{\Phi_1} + \frac{1}{2} \frac{n+1}{n} I_{\Phi_2} - \left(n\Phi(0) - \frac{1}{2} \Phi'(0) + \frac{1}{4n} \Phi''(0) \right) \\ &\quad + \left((n+1)\Sigma_{\Phi_1, n} - \frac{1}{2} \frac{n+1}{n} \Sigma_{\Phi_2, n} - \frac{1}{6} \frac{n+1}{n^2} \Sigma_{\Phi_3, n} \right). \end{aligned}$$

Using the relations $I_{F_1} = I_{\Phi_1} + \Phi(0)$ and $I_{\Phi_2} = 2I_{\Phi_1} - \Phi'(0)$, we obtain

$$\begin{aligned} \Sigma_{2, n} &= nI_{\Phi_1} + n\Phi(0) - (n+1)I_{\Phi_1} + \frac{n+1}{n} I_{\Phi_1} - \frac{1}{2} \frac{n+1}{n} \Phi'(0) - \left(n\Phi(0) - \frac{1}{2} \Phi'(0) + \frac{1}{4n} \Phi''(0) \right) \\ &\quad + \left((n+1)\Sigma_{\Phi_1, n} - \frac{1}{2} \frac{n+1}{n} \Sigma_{\Phi_2, n} - \frac{1}{6} \frac{n+1}{n^2} \Sigma_{\Phi_3, n} \right) \\ &= \frac{1}{n} I_{\Phi_1} - \frac{1}{4n} \Phi''(0) - \frac{1}{2n} \Phi'(0) + \left((n+1)\Sigma_{\Phi_1, n} - \frac{1}{2} \frac{n+1}{n} \Sigma_{\Phi_2, n} - \frac{1}{6} \frac{n+1}{n^2} \Sigma_{\Phi_3, n} \right). \end{aligned}$$

The latter equality and estimates (C.9) yield

$$\begin{aligned} |\Sigma_{2, n}| &\leq \frac{1}{n} \left(|I_{\Phi_1}| + \frac{|\Phi''(0)|}{4} + \frac{|\Phi'(0)|}{2} + \frac{K_{\Phi_3}}{3} \right) + K_{\Phi_1} \frac{1}{n^{1/2}} + \frac{K_{\Phi_2}}{2} \frac{1}{n^{3/2}} \\ &\leq \frac{1}{n^{1/2}} \left(K_{\Phi_1} + \left(|I_{\Phi_1}| + \frac{|\Phi''(0)|}{4} + \frac{|\Phi'(0)|}{2} + \frac{K_{\Phi_3}}{3} \right) \frac{1}{N^{1/2}} + \frac{K_{\Phi_2}}{2} \frac{1}{N} \right). \end{aligned}$$

Combining this with (C.1) and (C.2), we have

$$|(n+1)^2 R_{n+1} - n^2 R_n| \leq \bar{L}_1 \frac{1}{\sqrt{n}},$$

where

$$\bar{L}_1 := L_1 + K_{\Phi_1} + \left(|I_{\Phi_1}| + \frac{|\Phi''(0)|}{4} + \frac{|\Phi'(0)|}{2} + \frac{K_{\Phi_3}}{3} \right) \frac{1}{N^{1/2}} + K_{\Phi_2} \frac{1}{N}.$$

C.2 Proof of Proposition 4.4

Fix $c_1^{\max} > c_1$ and $c_1^{\min} \in (\frac{2}{3\sqrt{2}-2}c_1, c_1)$. Let us choose $\delta \in (0, 1)$ such that, for $x \in [1 - \delta, 1)$,

$$\frac{c_1^{\min}}{(1-x)^{1/2}} \leq F_2(x) \leq \frac{c_1^{\max}}{(1-x)^{1/2}}, \quad (\text{C.10})$$

$$\frac{1}{2} \frac{c_1^{\min}}{(1-x)^{3/2}} \leq F_2'(x) \leq \frac{1}{2} \frac{c_1^{\max}}{(1-x)^{3/2}}. \quad (\text{C.11})$$

Using the representation (4.21) and (C.10), we choose $C = C(F) > 0$ such that

$$\int_{-1}^{\beta} |F_2'(x)| dx \leq F(\beta) + C, \quad \beta \in [1 - \delta, 1). \quad (\text{C.12})$$

Finally, we assume that (increasing L_2 and l_2 if necessarily)

$$n \geq \frac{2}{\delta}. \quad (\text{C.13})$$

Integrating by parts, we have

$$\begin{aligned} \int_{-1}^1 F_2(x) dx &= \sum_{k=-n}^{n-1} \int_{k/n}^{(k+1)/n} F_2(x) dx \\ &= \sum_{k=-n}^{n-1} \left(\left(x - \frac{k+1}{n} \right) F_2(x) \Big|_{k/n}^{(k+1)/n} - \int_{k/n}^{(k+1)/n} \left(x - \frac{k+1}{n} \right) F_2'(x) dx \right) \\ &= \sum_{k=-n}^{n-1} \frac{1}{n} F_2 \left(\frac{k}{n} \right) + \Sigma_{1,n}, \end{aligned}$$

where

$$\Sigma_{1,n} := \sum_{k=-n}^{n-1} \int_{k/n}^{(k+1)/n} \left(\frac{k+1}{n} - x \right) F_2'(x) dx.$$

Inequalities (C.10)–(C.13) imply that

$$\begin{aligned}
|\Sigma_{1,n}| &= \left| \int_{1-1/n}^1 (1-x)F_2'(x) + \sum_{k=-n}^{n-2} \int_{k/n}^{(k+1)/n} \left(\frac{k+1}{n} - x \right) F_2'(x) dx \right| \\
&\leq \int_{1-1/n}^1 \frac{1}{2} \frac{c_1^{\max}}{(1-x)^{1/2}} + \sum_{k=-n}^{n-2} \frac{1}{n} \int_{k/n}^{(k+1)/n} |F_2'(x)| dx \\
&\leq \frac{c_1^{\max}}{\sqrt{n}} + \frac{1}{n} \int_{-1}^{1-1/n} |F_2'(x)| dx.
\end{aligned} \tag{C.14}$$

Since $1 - 1/n > 1 - \delta$ due to (C.13), we conclude from (C.10), (C.12) and (C.14) that

$$|\Sigma_{1,n}| \leq \frac{c_1^{\max}}{\sqrt{n}} + \frac{1}{n} \left(F_2 \left(1 - \frac{1}{n} \right) + C \right) \leq 2c_1^{\max} \frac{1}{\sqrt{n}} + C \frac{1}{n}.$$

Taking $L_2 := 2c_1^{\max} + \frac{C}{\sqrt{N}}$, we obtain the second inequality in (4.22).

Set $k_0 := [(1 - \delta)n]$, where $[\cdot]$ is the integer part of a real number. We represent $\Sigma_{1,n}$ as follows:

$$\Sigma_{1,n} = \Sigma_{2,n} + \Sigma_{3,n}, \tag{C.15}$$

where

$$\begin{aligned}
\Sigma_{2,n} &:= \left(\sum_{k=-n}^{k_0-1} \int_{k/n}^{(k+1)/n} \left(\frac{k+1}{n} - x \right) F_2'(x) dx + \int_{k_0}^{1-\delta} \left(\frac{k_0+1}{n} - x \right) F_2'(x) dx \right), \\
\Sigma_{3,n} &:= \left(\int_{1-\delta}^{(k_0+1)/n} \left(\frac{k_0+1}{n} - x \right) F_2'(x) dx + \sum_{k=k_0+1}^{n-1} \int_{k/n}^{(k+1)/n} \left(\frac{k+1}{n} - x \right) F_2'(x) dx \right).
\end{aligned} \tag{C.16}$$

Below we estimate $\Sigma_{2,n}$ and $\Sigma_{3,n}$ separately.

Inequality (C.12) implies that

$$\Sigma_{2,n} \geq -\frac{1}{n} \left(\int_{-1}^{1-\delta} |F_2'(x)| dx \right) \geq -\frac{1}{n} (F_2(1 - \delta) + C). \tag{C.17}$$

Due to (C.11) all the terms in expression $\Sigma_{3,n}$ are positive. Inequality (C.13) implies that the last sum in formula (C.16) contains at least two terms. Hence, inequality (C.11) implies that

$$\Sigma_{3,n} \geq \int_{1-2/n}^{1-1/n} \left(1 - \frac{1}{n} - x \right) F_2'(x) dx + \int_{1-1/n}^1 (1-x)F_2'(x) dx \tag{C.18}$$

$$\geq \int_{1-2/n}^{1-1/n} \frac{1}{2} \left(1 - \frac{1}{n} - x \right) \frac{c_1^{\min}}{(1-x)^{3/2}} dx + \int_{1-1/n}^1 \frac{1}{2} \frac{c_1^{\min}}{(1-x)^{1/2}} dx \tag{C.19}$$

$$= \int_{1-2/n}^1 \frac{1}{2} \frac{c_1^{\min}}{(1-x)^{1/2}} dx - \frac{1}{2n} \int_{1-2/n}^{1-1/n} \frac{c_1^{\min}}{(1-x)^{3/2}} dx = \frac{3\sqrt{2}-2}{2} c_1^{\min} \frac{1}{\sqrt{n}}. \tag{C.20}$$

Relations (C.15), (C.17), and (C.20) imply that

$$\Sigma_{1,n} \geq L_2^* \frac{1}{\sqrt{n}} - l_2 \frac{1}{n},$$

where $L_2^* := \frac{3\sqrt{2}-2}{2} c_1^{\min} > c_1$ and $l_2 := F_2(1-\delta) + C$.

D Requirements on $n_0(E)$

In this appendix, we collect the constants that we use throughout the paper to determine $n_0(E)$ as well as all the 12 requirements on the number $n_0 = n_0(E)$ entering Definition 3.1 of admissible E .

D.1 Constants not depending on a or E

1. $\tau_0 > 0$ is an arbitrarily fixed real number (see (4.9)).

2. Set (see (4.10))

$$A_0 := \sup_{n \geq 0, t \geq \tau_0} \sqrt{t} \left| y_n(t) - \sqrt{t} f \left(\frac{n}{\sqrt{t}} \right) \right|.$$

3. Set (see (4.10))

$$A_1 := \sup_{n \geq 0, t \geq \tau_0} t\sqrt{t} \left| \dot{y}_n(t) - \frac{1}{\sqrt{t}} h \left(\frac{n}{\sqrt{t}} \right) \right|.$$

4. Set (see (4.10))

$$\tilde{A}_1 := \sup_{n \geq 0, t \geq \tau_0} t\sqrt{t} \left| y_n(t) - \sqrt{t} f \left(\frac{n}{\sqrt{t}} \right) - \frac{1}{\sqrt{t}} \tilde{f} \left(\frac{n}{\sqrt{t}} \right) \right|.$$

5. Set (see (4.10))

$$A_2 := \sup_{n \geq 0, t \geq \tau_0} t^2\sqrt{t} \left| \ddot{y}_n(t) - \frac{1}{t\sqrt{t}} h'' \left(\frac{n}{\sqrt{t}} \right) \right|.$$

6. Set (see (4.11))

$$B_0 := \sup_{n \geq 0, t \geq \tau_0} t \left| \nabla y_n(t) - g \left(\frac{n}{\sqrt{t}} \right) - \frac{1}{2\sqrt{t}} h \left(\frac{n}{\sqrt{t}} \right) \right|.$$

7. Set (see (4.11))

$$B_1 := \sup_{n \geq 0, t \geq \tau_0} t\sqrt{t} \left| \nabla \dot{y}_n(t) - \frac{1}{t} h' \left(\frac{n}{\sqrt{t}} \right) \right|.$$

8. Set (see (4.12))

$$A_2^* := \sup_{n \geq 0, t \geq \tau_0} t\sqrt{t}|\ddot{y}_n(t)|.$$

9. Set (see (4.12))

$$B_2^* := \sup_{n \geq 0, t \geq \tau_0} t^2|\nabla\ddot{y}_n(t)|.$$

10. We use the notation (see (5.19))

$$R := \pi.$$

11. B_{h2} is given by (see (7.23))

$$B_{h2} := \sup_{x \in [0, \infty)} |h''(x)| = -h''(0).$$

12. B_{h4} is given by (see (7.23))

$$B_{h4} := \sup_{x \in [0, \infty)} |h''''(x)| = h''''(0).$$

D.2 Constants depending on a but not depending on E

1. a is a unique root of equation (4.18) (or equivalently (4.19), (4.20)).

2. Consider the function

$$h_a(x) := h\left(\frac{x}{\sqrt{a}}\right) + h\left(\frac{1}{x\sqrt{a}}\right).$$

Set (see (5.1))

$$D_a := \inf_{x \in (0, 1]} h_a(x).$$

3. Set (see (5.2))

$$p := \sup_{x \in (0, 1]} \frac{h'_a(x)x}{h_a(x)}.$$

4. N is a fixed natural number satisfying (see (5.9), (5.10))

$$\begin{aligned} & N \left(\left(1 + \frac{1}{N} \right)^{\frac{1+p}{2}} - 1 \right) - \left(2^{\frac{1-p}{2}} - 1 \right) \\ & - 2 \left(2^{\frac{1-p}{2}} - 1 + N \left(\left(1 + \frac{1}{N} \right)^{\frac{1+p}{2}} - 1 \right) \right) \left(2^{\frac{1-p}{2}} - 1 \right) \frac{1}{N} > 0. \end{aligned}$$

5. Set (see (5.3))

$$D_{p1} := 2^{\frac{1-p}{2}} - 1,$$

$$D_{p2} := N \left(\left(1 + \frac{1}{N} \right)^{\frac{1+p}{2}} - 1 \right).$$

6. Constants needed to define K and K' from Lemma 5.3

(a) Set (see (5.35))

$$L_1(F, N) := \sup_{n \geq N} n^{3/2} \left| \int_{-1}^1 F(x) dx - \sum_{|k| \leq n-1} \frac{1}{n} F\left(\frac{k}{n}\right) \right|,$$

where the supremum exists due to Proposition 4.3. Set

$$K'_1 := h_1 L_1(F, N).$$

(b) Set (see (5.36))

$$l_1 := h_1 A_0 \frac{1}{a^{1/2}} R.$$

(c) Set (see (5.37))

$$K' := K'_1 + l_1 \frac{1}{\sqrt{N}}.$$

(d) Set

$$K_1 := \sup_{n \geq N} n^{1/2} |C_{n+1}^{(1)} - C_n^{(1)}|,$$

where

$$C_l^{(1)} := h_1 \left(l^2 I_F - \sum_{|k| \leq l-1} \sqrt{a(l^2 - k^2)} f\left(\frac{l-k}{\sqrt{a(l^2 - k^2)}}\right) \right), \quad l = n, n+1.$$

Note that the supremum exists due to (5.38).

(e) Consider the function

$$\tilde{F}(x) := \frac{1}{\sqrt{a(1-x^2)}} \tilde{f}\left(\frac{1-x}{\sqrt{a(1-x^2)}}\right)$$

and constants $L_{\tilde{f}}, L_{\tilde{f}}^*, l_{\tilde{f}} \geq 0$ such that (see (5.40))

$$L_{\tilde{f}}^* \frac{1}{\sqrt{n}} - l_{\tilde{f}} \frac{1}{n} \leq - \int_{-1}^1 \tilde{F}(x) dx + \sum_{-n \leq k \leq n-1} \frac{1}{n} \tilde{F}\left(\frac{k}{n}\right) \leq L_{\tilde{f}} \frac{1}{\sqrt{n}}, \quad n \geq N.$$

Set (see (5.42))

$$K_2 := h_1 \left(L_{\tilde{f}} - L_{\tilde{f}}^* + l_{\tilde{f}} \frac{1}{\sqrt{N}} \right).$$

(f) Set (see (5.43))

$$K_3 := 2h_1 \tilde{A}_1 \frac{1}{a^{3/2}} T_{3/2}.$$

(g) Set (see (5.44))

$$K := K_1 + K_2 + K_3 \frac{1}{N}.$$

Remark D.1. In principle, due to Proposition 5.3, we could define $K' := \sup_{n \geq N} n^{-1/2} |C_n|$ and $K := \sup_{n \geq N} n^{1/2} |C_{n+1} - C_n|$. However, calculation of the values C_n is computationally consuming as it involves Bessel functions. We used the strategy described above, since it is based on error estimates of Riemann sums only.

7. Set (see (5.9))

$$\varkappa := D_{p2} - D_{p1} - 2(D_{p1} + D_{p2})D_{p1} \frac{1}{N} (> 0).$$

8. Set (see (5.13))

$$E_0 := \frac{K + \varkappa K'}{(h_1 - 2c)\varkappa} (> 0).$$

9. For $\alpha > 0$, set (see (5.17))

$$S_\alpha := \sup_{n \geq N} \left(\sqrt{n} \sum_{|k| \leq n-1} \frac{(n-k)^\alpha}{(n^2 - k^2)^{\alpha+1/2}} \right).$$

We use only the values of S_1 , S_2 , and S_3 to determine $n_0(E)$.

10. For $\alpha > 1$, set (see (5.18))

$$T_\alpha := \sup_{n \geq N} \left(\sum_{|k| \leq n-1} \frac{n^\alpha}{(n^2 - k^2)^\alpha} \right).$$

We use only the values of $T_{3/2}$, T_2 , and $T_{5/2}$ to determine $n_0(E)$.

11. Let constants $L_2^*(H, N) > \frac{1}{\sqrt{2a}} h(0)$ and $l_2(H, N) \geq 0$ be such that (see Proposition 4.4)

$$L_2^*(H, N) \frac{1}{n^{1/2}} - l_2(H, N) \frac{1}{n} \leq I_H - \sum_{k=-n}^{n-1} \frac{1}{n} H\left(\frac{k}{n}\right), \quad n \geq N,$$

where $H(x)$ is given by (4.15) and $I_H = I_H(a)$ is given by (4.17). Set (see (5.51))

$$\begin{aligned} C_H &:= L_2^*(H, N), \\ l_H &:= l_2(H, N). \end{aligned}$$

12. Consider the function $\bar{H}(x) := \frac{1}{1-x^2}H(x)$. Set (see (5.52))

$$C_{H2} := \sup_{n \geq N} n^{-1/2} \left| \sum_{|k| \leq n-1} \frac{1}{n} \bar{H}\left(\frac{k}{n}\right) \right|.$$

13. Set (see (6.3))

$$K_{h1} := \sup_{n \geq N} n^{-1} \left| \sum_{|k| \leq n-1} \frac{1}{a(1-(k/n)^2)} h' \left(\frac{1}{\sqrt{a}} \sqrt{\frac{1-k/n}{1+k/n}} \right) \right|.$$

14. Set (see Lemma 6.6 and (6.9))

$$K_g := \sup_{n \geq N} n^{3/2} \left| \sum_{|k| \leq n-1} \frac{1}{n} G\left(\frac{k}{n}\right) - I_G - \frac{1}{4n} \right|,$$

where $G(x)$ is given by (4.14), and $I_G = I_G(a)$ is given by (4.17).

15. Set (see Lemma 6.7 and (6.10))

$$K_h := \sup_{n \geq N} n^{1/2} \left| \sum_{|k| \leq n-1} \frac{1}{n} H\left(\frac{k}{n}\right) - I_H \right|,$$

where $H(x)$ is given by (4.15), and $I_H = I_H(a)$ is given by (4.17).

16. θ_0 satisfies (see (7.2))

$$\theta_0 K_h < \frac{1}{4}.$$

17. η satisfies (see (7.15))

$$\eta \in \left(0, \frac{1}{4\sqrt{\pi}}\right).$$

18. $x_0 \in [0, 1)$ and $\varepsilon_0 > 0$ satisfy (see (7.16), (7.17))

$$h'' \left(\frac{1-x}{\sqrt{a(1-x^2)} + \varepsilon_1} \right) + h'' \left(\frac{1+x}{\sqrt{a(1-x^2)} + \varepsilon_1} \right) \leq -\eta$$

for all $x \in [x_0, (n-1)/n]$ and ε_1 with

$$-\min \left(\varepsilon_0, \frac{a}{2n} \left(2 - \frac{1}{n} \right) \right) \leq \varepsilon_1 \leq \varepsilon_0, \quad n \geq \max \left(N, \frac{1}{1-x_0} \right)$$

19. b satisfies (see (7.20))

$$b > 2a.$$

20. Set (see (7.21))

$$R_1 := \sup_{n \geq N} n^2 \sum_{|k| < x_0 n} \frac{1}{(n^2 - k^2)^{3/2}}$$

D.3 Constants depending on E

1. Set (see (5.14))

$$a_n^{\min} := a - \frac{2En^{1/2}}{2n-1}.$$

2. Set (see (5.14))

$$a_n^{\max} := a + \frac{2En^{1/2}}{2n-1}.$$

3. Set (see (5.16))

$$\delta_n := \frac{2A_1}{aD_a} \frac{1}{n} + \frac{2A_2^* E a^{1/2}}{D_a (a_n^{\min})^{3/2}} \frac{1}{\sqrt{n}}.$$

4. Set (see (7.22))

$$R_2 := \inf_{n \geq \max(N, 1/(1-x_0))} \sum_{k \in [x_0 n, n-1]} \frac{n^{3/2}}{(a_n^{\max}(n^2 - k^2) + bn)^{3/2}}.$$

D.4 Requirements on $n_0(E)$

We assume that the following requirements hold for $n \geq n_0(E)$:

1. $n \geq N$.

2. $a_n^{\min} \geq \frac{\tau_0}{2n-1}$.

3. $\delta_n + \delta_{n+1} \leq 2D_{p1}$.

4. $C_H - \frac{1}{\sqrt{2a}} h(0) \geq l_H \frac{1}{\sqrt{n}} + \frac{\varkappa}{\sqrt{2a}} h(0) \frac{1}{n} + C_{H2} \delta_n + \frac{1}{\sqrt{2a}} \left(1 + \frac{\varkappa}{n}\right) \left(h(0) \frac{n+1}{2n+1} \delta_{n+1} + (1 + \delta_{n+1}) h\left(\sqrt{\frac{2n+1}{a}}\right) \right)$.

5. $2EK_{h1} n^{-1/2} + \left(\frac{2EB_1 T_{3/2}}{a^{3/2}} + \frac{2E^2 B_2^* T_2}{(a_n^{\min})^2} \right) n^{-1} + \sum_{|k| \leq n-1} \frac{B_0}{a(n^2 - k^2)} + \left(K_g + \frac{K_h}{2} \right) n^{-1/2} \leq \frac{3}{8}$.

6. $\left(K_{h1} + \frac{(2E+\theta_0)A_2^* T_{3/2}}{(a_n^{\min})^{3/2}} \right) n^{-1/2} + \left(\frac{B_1 T_{3/2}}{a^{3/2}} + \frac{(2E+\theta_0)B_2^* T_2}{(a_n^{\min})^2} + \frac{A_1 T_{3/2}}{a^{3/2}} \right) n^{-1} \leq \frac{K_h}{2}$.

7. $a(n-1)^2 - E(n-1)^{1/2} + b(n-1) \geq an^2 + En^{1/2}$.

8. $\frac{2E-\theta_0}{n^{3/2}} \leq \min\left(\varepsilon_0, \frac{a}{2n}\left(2 - \frac{1}{n}\right)\right)$ for $n \geq \max\left(n_0(E), \frac{1}{1-x_0}\right)$.
9. $2En^{-3/2} + bn^{-1} \leq \varepsilon_0$ for $n \geq \max\left(n_0(E), \frac{1}{1-x_0}\right)$.
10. $\tau_0 \leq \theta_0 n^{1/2}$.
11. $\frac{B_{h4}}{2\theta_0} n^{-1/2} + \sup_{x \geq \frac{2n+1}{\sqrt{bn}}} h''(x) + \frac{2A_2}{\theta_0^{1/2}} n^{-1/2} \leq \frac{1}{4\sqrt{\pi}}$.
12. $\left(\frac{B_{h2}R_1}{(a_n^{\min})^{3/2}} + \frac{B_2^*T_2}{(a_n^{\min})^2}\right) n^{-1/2} + \frac{A_2T_{5/2}}{(a_n^{\min})^{5/2}} n^{-1} \leq \eta R_2$.

Acknowledgement

The authors are grateful to Daria Neverova for her help in preparing the figures. Pavel Gurevich acknowledges the support of DFG through the Heisenberg fellowship and the Collaborative Research Center SFB 910 (Germany). Sergey Tikhomirov acknowledges the support of Humboldt postdoctoral fellowship, Chebyshev Laboratory under RF Government grant 11.G34.31.0026, JSC ‘‘Gazprom neft’’, Saint-Petersburg State University research grant 6.38.223.2014, RFBR 15-01-03797a. Both authors acknowledge the support of the German-Russian Interdisciplinary Science Center (G-RISC).

References

- [1] M. Abramowitz, I. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. National Bureau of Standards Applied Mathematics Series, 55 (1965).
- [2] T. Aiki, J. Kopfova, *A Mathematical Model for Bacterial Growth Described by a Hysteresis Operator*. Recent advances in nonlinear analysis – Proceedings of the International Conference on Nonlinear Analysis. Hsinchu, Taiwan, 2006.
- [3] H. W. Alt, *On the thermostat problem*. Control Cyb., **14** (1985), 171–193.
- [4] D. E. Amos, *Computation of modified Bessel functions and their ratios*. Math. Comp., **47**, No. 125 (1974), 239–251.
- [5] D. Apushkinskaya, N. Uraltseva, *On regularity properties of solutions to the hysteresis-type problems*. <http://arxiv.org/abs/1410.6719>, to appear in Interfaces and Free Boundaries, **17** (2015), 93–115, DOI: 10.4171/IFB/335.
- [6] D. Apushkinskaya, N. Uraltseva, *Free boundaries in problems with hysteresis*. <http://arxiv.org/abs/1411.5376>, to appear in Philos. Trans. A, **373** (2015).

- [7] D. Apushinskaya, N. Uraltseva, H. Shahgholian, *Lipschitz property of the free boundary in the parabolic obstacle problem*. St. Petersburg Math. J., **15** (2004), 375–391.
- [8] M. Brokate, J. Sprekels, *Hysteresis and Phase Transitions*, Springer, 1996.
- [9] M. Curran, *Local well-posedness of a reaction-diffusion equation with hysteresis*. Masters Thesis, Free University of Berlin, 2014.
- [10] P. Gurevich, *Asymptotics of parabolic Green’s functions on lattices*. Preprint. <http://arxiv.org/abs/1504.02673>
- [11] P. Gurevich, S. Tikhomirov, *Uniqueness of transverse solutions for reaction-diffusion equations with spatially distributed hysteresis*. Nonlinear Anal., **75** (2012), 6610–6619.
- [12] P. Gurevich, R. Shamin, S. Tikhomirov, *Reaction-diffusion equations with spatially distributed hysteresis*. SIAM J. Math. Anal., **45**, no. 3 (2013), 1328–1355.
- [13] P. Gurevich, S. Tikhomirov, *Systems of reaction-diffusion equations with spatially distributed hysteresis*. Math. Bohem., **139**, no. 2 (2014), 239–257.
- [14] F.C. Hoppensteadt, W. Jäger, *Pattern formation by bacteria*. Lecture Notes in Biomath. 38 (W. Jäger, H. Rost, P. Tautu, eds.). Springer, Berlin, 1980, pp. 68–81.
- [15] F.C. Hoppensteadt, W. Jäger, C. Pöppe, *A hysteresis model for bacterial growth patterns*. Modelling of patterns in space and time. Lecture Notes in Biomath. 55 (W. Jäger, J.D. Murray, eds.). Springer, Berlin, 1984, pp. 123–134.
- [16] A. M. Il’in, B. A. Markov, *A nonlinear diffusion equation and Liesegang rings*. Dokl. Math., **84**, no. 2, (2011), 730–733.
- [17] J. Kopfová, *Hysteresis in biological models*. Proceedings of the conference “International Workshop on Multi-rate processes and hysteresis”. Journal of Physics, Conference Series, **55** (2007), 130–134.
- [18] M. A. Krasnosel’skii, A. V. Pokrovskii. *Systems with Hysteresis*. Springer-Verlag. Berlin–Heidelberg–New York (1989). Translated from Russian: *Sistemy s Gistereziom*. Nauka. Moscow (1983).
- [19] P. Krejčí. *Hysteresis, convexity and dissipation in hyperbolic equations*. Gakuto International Series. Mathematical Sciences and Applications, 8. Gakkotosho Co., Ltd., Tokyo, 1996.
- [20] I. D. Mayergoyz, *Mathematical Models of Hysteresis*, Springer, 1991.
- [21] A. Mielke, *Evolution of rate-independent systems*, Evolutionary Equations. Vol. II, 461–559, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2005.

- [22] A. Petrosyan, H. Shahgholian, N. Uraltseva, Regularity of Free Boundaries in Obstacle-Type Problems. Graduate Studies in Mathematics, **36**, 2012.
- [23] H. Shahgholian, N. Uraltseva, G. Weiss, *A parabolic two-phase obstacle-like equation*. Adv. Math., **221** (2009), 861–881.
- [24] A. Visintin, *Evolution problems with hysteresis in the source term*. SIAM J. Math. Anal, **17** (1986), 1113–1138.
- [25] A. Visintin. *Differential Models of Hysteresis*. Springer-Verlag. Berlin — Heidelberg (1994).
- [26] A. Visintin. *Ten issues about hysteresis*. Acta Appl. Math., **132** (2014), 635–647.