

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

On absolute nonshadowability of transitive maps

by

Sergey Tikhomirov

Preprint no.: 36

2016



On absolute nonshadowability of transitive maps

Sergey Tikhomirov*

Abstract

We study shadowing property for random infinite pseudotrajectories of a continuous map f of a compact metric space. For the cases of transitive maps and transitive attractors we prove a dichotomy: either f satisfies shadowing property or random pseudotrajectory is shadowable with probability 0.

Keywords: shadowing, transitivity, attractor, Markov chain

1 Introduction

The theory of shadowing of approximate trajectories (pseudotrajectories) of dynamical systems is now a well-developed part of the global theory of dynamical systems (see the monographs [10, 11] and [13] for a survey of modern results). The shadowing problem is related to the following question: under which conditions, for any pseudotrajectory of f does there exist a close trajectory?

It is known that a diffeomorphism f has the shadowing property in a neighborhood of a hyperbolic set [3, 4]. Moreover if f is structurally stable (see definition for example in [7, 12]) then it has the shadowing property on the whole manifold. At the same time, it is easy to give an example of a diffeomorphism that is not structurally stable but has the shadowing

*Max Planck Institute for Mathematics in the Science Inselstrasse 22, 04103 Leipzig, Germany. Chebyshev Laboratory, Saint-Petersburg State University 14th lane 29B, Vasilievsky Island, St. Petersburg, 199178, Russia. email:sergey.tikhomirov@gmail.com

property (see [17], for instance). Thus, structural stability is not equivalent to shadowing.

At the same time it was proved that in several contexts shadowing and structural stability are equivalent. Sakai prove that the C^1 -interior of the set of diffeomorphisms having the shadowing property coincides with the set of structurally stable diffeomorphisms [16] (see [14] for a similar result for the orbital shadowing property). Abdenur and Diaz conjectured that a C^1 -generic diffeomorphism with the shadowing property is structurally stable; they have proved this conjecture for the so-called tame diffeomorphisms [2]. Jointly with S. Pilyugin the author proved that so-called Lipschitz shadowing is equivalent to structural stability [15]. Thus, set of not structurally stable diffeomorphisms satisfying shadowing property is not very reach.

It is a natural problem to find a shadowing property which is satisfied for a broader class of diffeomorphisms. One of the possible approaches is to consider random pseudotrajectories: endow the space of pseudotrajectories with a probability measure and find sufficient conditions for probability of a pseudotrajectory to be shadowable to be close to 1 or at least positive.

Such studies were initiated in [19]. In this work Yuan, Yorke constructed an open set of diffeomorphisms for which probability of a pseudotrajectory to be shadowable is 0. In a recent work [18] the author considered a special example of linear skew product and found probability of a finite pseudotrajectory to be shadowable.

Despite the naturalness of randomness approach currently consideration of finite pseudotrajectories is more developed [5,6,17,18]. One of the reasons is lack of positive results about shadowing of random pseudotrajectories.

In the present paper we prove that such a positive result is not possible: under transitivity assumption either all pseudotrajectories are shadowable or probability of a pseudotrajectory to be shadowable is 0. Precise statements of the results for transitive maps and transitive attractors are formulated in Theorems 1, 2 respectively.

2 Transitive maps

Let (M, dist) be a compact metric space endowed with a finite Borel measure μ , such that for any open set U the inequality $\mu(U) > 0$ holds. For $a > 0$, $x \in M$ denote by $B(a, x)$ the open ball of radius a centered at x . Let $f : M \rightarrow M$ be a homeomorphism.

Let $I = [0, N]$ or $I = [0, +\infty)$. For $d > 0$ we say that sequence $\{y_n\}_{n \in I}$ is a d -pseudotrajectory if

$$\text{dist}(y_{n+1}, f(y_n)) \leq d, \quad n, n+1 \in I. \quad (1)$$

We say that a d -pseudotrajectory $\{y_n\}_{n \in I}$ can be ε -shadowed for $\varepsilon > 0$ if there exists $x_0 \in M$ such that

$$\text{dist}(y_n, f^n(x_0)) \leq \varepsilon, \quad n \in I. \quad (2)$$

We say that f has *shadowing property* if for any $\varepsilon > 0$ there exists $d > 0$ such that any d -pseudotrajectory $\{y_n\}_{n \geq 0}$ can be ε -shadowed.

Remark 1. *In the definitions of pseudotrajectories and shadowing in equations (1), (2) usually are used strict inequalities. The definition of the shadowing property with not strict inequalities is equivalent to the classical one and allows us to simplify the notation in the proofs of main results.*

For $y_0 \in M$, $d > 0$, $N \in \mathbb{N} \cup \{+\infty\}$ denote by $\Omega_N(y_0, d)$ the set of all d -pseudotrajectories $\{y_n\}_{n=0}^{+\infty}$ starting at y_0 . Let us consider point y_{n+1} being chosen at random in $B(d, f(y_n))$ uniformly with respect to the measure μ : for a measurable set $A \subset M$ the equality

$$P(y_{n+1} \in A | y_n) = \frac{\mu(A \cap B(d, f(y_n)))}{\mu(B(d, f(y_n)))}$$

holds. Then $\Omega_N(y_0, d)$ forms a Markov chain. This naturally endows $\Omega_N(y_0, d)$ with a probability measure $P_N^{y_0, d}$. For simplicity for $N = +\infty$ we will omit it: $\Omega(y_0, d)$, $P^{y_0, d}$.

Remark 2. *This concept was introduced in [19] for infinite pseudotrajectories, see also [18] for a similar concept for finite pseudotrajectories.*

For any $y_0 \in M$, $d > 0$, $\varepsilon > 0$, $N \in \mathbb{N} \cup \{+\infty\}$ consider set $Sh_N(y_0, d, \varepsilon) \subset \Omega_N(y_0, d)$ of pseudotrajectories $\{y_n\}$ which can be ε -shadowed. For $N = +\infty$ we denote the set as $Sh(y_0, d, \varepsilon)$. Note that each of the sets $Sh_N(y_0, d, \varepsilon)$ is closed for $N \in \mathbb{N}$ and is measurable with respect to $P_N^{y_0, d}$. Hence $Sh(y_0, d, \varepsilon)$ is a countable intersection of measurable events and is measurable itself.

Denote

$$p(y_0, d, \varepsilon) = P^{y_0, d}(Sh(y_0, d, \varepsilon))$$

the probability that a d -pseudotrajectory starting at y_0 can be ε -shadowed.

We say that a map f is *absolutely not shadowable* if there exists $\varepsilon > 0$ such that for any $y_0 \in M$, $d > 0$, $p(y_0, d, \varepsilon) = 0$. Speaking informally almost any trajectory of f cannot be ε -shadowed. In [19] Yuan and Yorke provided a class of examples of absolutely not shadowable diffeomorphism. Based on the same technique Abdenur and Diaz proved that absolute nonshadowability is C^1 generic among not structurally stable maps.

We say that f is *transitive* if there exists $r \in M$ such that

$$\overline{O^+(r, f)} = M. \quad (3)$$

In [2] Abdenur and Diaz proved that C^1 -robustly transitive and not hyperbolic diffeomorphisms are absolutely not shadowable. Their proof is based on construction of periodic orbits with different indices.

In the present paper we remove the differentiability assumption, and what is more important, do not assume any properties of the perturbation of f .

Theorem 1. *If f is a transitive map then one of the following holds*

- (i) *f has the shadowing property;*
- (ii) *f is absolutely not shadowable.*

In the proof we will use the following folklore result.

Lemma 1. *If for any $\varepsilon > 0$ there exists $d > 0$ such that any finite d -pseudotrajectory $\{y_n\}_{n=0}^N$, $N > 0$ can be ε -shadowed then f satisfies the shadowing property.*

Proof of Theorem 1. Assume that f does not satisfy shadowing property.

Lemma 1 implies that there exists $\varepsilon > 0$ such that for any $d > 0$ there exists N and a $d/2$ -pseudotrajectory $\{p_n\}_{n=0}^N$ which cannot be ε -shadowed.

Take this $\varepsilon > 0$ and fix arbitrarily $d > 0$ and corresponding N and $\{p_n\}_{n=0}^N$ a not ε -shadowable $d/2$ -pseudotrajectory. Below we will show that

$$p(y_0, d, \varepsilon/2) = 0. \quad (4)$$

Let us note that any sequence $\{z_n\}_{n=0}^N$ satisfying

$$\text{dist}(p_n, z_n) < \varepsilon/2$$

is not $\varepsilon/2$ -shadowable.

Since f is continuous there exists $\delta > 0$ such that for any $x \in M$, $y \in B(d/2, f(x))$, $z \in B(\delta, x)$ the inclusion

$$B(\delta, y) \subset B(d, f(z)) \quad (5)$$

holds.

Denote

$$\eta = \frac{\inf_{x \in M} \mu(B(\delta, x))}{\sup_{x \in M} \mu(B(d, x))} \neq 0, \quad (6)$$

$$\mathcal{B} = \{\{q_n\}_{n=0}^N : q_n \in B(\delta, p_n)\}.$$

Let us show that for any $z_0 \in B(\delta, p_0)$ the inequality

$$P_N(\Omega_N(z_0, d) \cap \mathcal{B}) \geq \eta^N \quad (7)$$

holds. Indeed, let z_0, z_1, \dots, z_N be a random d -pseudotrajectory. Then for any $z_k \in B(\delta, p_k)$ the inequality

$$P_N(z_{k+1} \in B(\delta, p_{k+1}) | z_k) \geq \frac{\mu(B(\delta, p_{k+1}))}{\mu(B(d, f(z_k)))} \geq \eta$$

holds. Multiplying those inequalities for $k = 0, \dots, N-1$ and using Markov property we conclude that

$$P_N(z_{k+1} \in B(\delta, p_{k+1}), k = 0, \dots, N-1) \geq \eta^N,$$

which proves (7).

Let us consider finite covering $\{U_i\}$ of M by open balls of radius $\delta_1 = \delta/4$. Let $r \in M$ satisfies (3). Trajectory of point r visits each of $\{U_i\}$ infinitely many times. Let K_1 be such that $\{f^n(r)\}_{n=0}^{K_1}$ visits each of $\{U_i\}$ and K_2 be such that $\{f^n(f^{K_1+1}(r))\}_{n=0}^{K_2}$ visits each of $\{U_i\}$. Set $K = K_1 + K_2 + 1$. For any $z_0 \in M$ there exists $0 \leq n_1 \leq n_2 \leq K$ such that

$$\text{dist}(z_0, f^{n_1}(r)) < 2\delta_1, \quad \text{dist}(p_0, f^{n_2}(r)) < 2\delta_1. \quad (8)$$

Consider sequence

$$\{q_n\}_{n=0}^{n_2-n_1+N} = \{f^{n_1}(r), \dots, f^{n_2-1}(r), p_0, p_1, \dots, p_N\}.$$

Due to inequalities (8) sequence $\{q_n\}$ is a $d/2$ -pseudotrajectory. Since it contains $\{p_n\}_{n=0}^N$ it cannot be $\varepsilon/2$ -shadowed. Similarly to (7)

$$P^{z_0, d}(z_n \in B(\delta, q_n), n \in [0, n_2 - n_1 + N]) \geq \eta^{n_2 - n_1 + N} \geq \eta^{K+N}.$$

Denote $L = K + N + 1$. Hence for any z_0

$$P^{z_0, d}(\{z_n\}_{n=0}^{L-1} \text{ is not } \varepsilon/2\text{-shadowable} | z_0) \geq \eta^{L-1} \geq \eta^L.$$

Similarly for any $k \geq 0$ and $y_{kL} \in M$

$$P^{y_0, d}(\{y_n\}_{n=kL}^{(k+1)L-1} \text{ is not } \varepsilon/2\text{-shadowable} | y_{kL}) \geq \eta^L.$$

Combining those inequalities we conclude that for any $k \geq 0$

$$P^{y_0, d}(\{y_n\}_{n=0}^{(k+1)L-1} \text{ is not } \varepsilon/2\text{-shadowable}) \geq 1 - (1 - \eta^L)^k.$$

The right-hand side of the latter expression tends to 1 as $k \rightarrow \infty$ hence

$$P^{y_0, d}(\{y_n\} \text{ is not } \varepsilon/2\text{-shadowable}) = 1.$$

Theorem is proved. □

3 Transitive attractors

We say that an invariant compact set Λ is *an attractor* if there exists an open neighborhood U of Λ such that $f(U) \subset U$ and $\bigcap_{n \geq 0} f^n(U) = \Lambda$, see for instance [8]. See book [9] for systematic studies of properties of attractors.

We will use the following two properties of an attractor:

1. $\text{dist}(f^n(x), \Lambda) \rightarrow 0$ as $n \rightarrow \infty$ for any $x \in U$;
2. for any neighborhood $V \subset U$ of Λ there exists a neighborhood $W \subset V$ such that

$$\overline{f(W)} \subset W. \tag{9}$$

Denote by $D(\Lambda)$ the domain of attraction of Λ :

$$D(\Lambda) := \{x \in M : \text{dist}(f^n(x), \Lambda) \rightarrow 0\}.$$

Note that $D(\Lambda) = \bigcup_{n \geq 0} f^{-n}(U)$.

We say that f has the shadowing property on Λ if for any $\varepsilon > 0$ there exists $d > 0$ such that for any d -pseudotrajectory $\{y_n\}_{n \geq 0} \subset \Lambda$ there exists $x_0 \in M$ (not necessarily belonging to Λ) such that inequalities (2) hold.

We say that set Λ is transitive if there exists $r \in \Lambda$ such that

$$\Lambda = \overline{O^+(r, f)}.$$

The assumption for a map to be transitive is quite restrictive. At the same time it is quite common for attractors. In some works transitivity is included in the definition of attractor [1]. In this work it was proved that transitive attractors persists under C^1 -generic perturbations.

Theorem 2. *Let Λ be a transitive attractor not satisfying shadowing property then there exists $\varepsilon_0 > 0$ such that for any $y_0 \in D(\Lambda)$ there exists $d_0 > 0$ such that for any $d < d_0$ the probability of a d -pseudotrajectory starting at y_0 to be ε_0 -shadowed is 0:*

$$p(y_0, d, \varepsilon_0) = 0.$$

Remark 3. *Note that in Theorem 1 the choice of d was uniform with respect to y_0 and hence Theorem 1 does not follow from Theorem 2.*

Remark 4.

Proof. Similarly to Lemma 1 there exists $\varepsilon > 0$ such that for any $d > 0$ there exists $N > 0$ and a d -pseudotrajectory $\{p_n\}_{n=0}^N$ which cannot be ε -shadowed.

Fix $\varepsilon_0 = \varepsilon/4$, $V = B(\varepsilon_0, \Lambda) \cap U$ and W , satisfying (9).

Fix $y_0 \in D(\Lambda)$. There exists n_0 such that $f^{n_0}(y_0) \in W$. Take $d_0 < \varepsilon/4$ such that

1. for any $\{y_n\}_{n=0}^{n_0} \in \Omega(y_0, d_0)$ the inclusion

$$y_{n_0} \in W \tag{10}$$

holds;

2. $B(d_0, \overline{f(W)}) \subset W$.

Then for any $d < d_0$ and $\{y_n\} \in \Omega(y_0, d)$ the inclusions

$$y_n \in W, \quad \text{for } n \geq n_0 \tag{11}$$

holds.

Fix arbitrarily $d < d_0$ and find $N > 0$ and a d -pseudotrajectory $\{p_n\}_{n=0}^N \subset \Lambda$ which cannot be ε -shadowed.

Take $\delta < d/4$ such that inclusion (5) holds. Due to compactness of M and hence compactness of \overline{W} there exists $S > 0$ such that

$$f^n(z) \in B(\delta/4, \Lambda), \quad n \geq S, z \in W.$$

Arguing similarly to the proof of Theorem 1 there exists $K > 0$ such that for any $z_0 \in W$ there exists $0 < n_1 < n_2 < K$ such that

$$\text{dist}(f^S(z_0), f^{n_1}(r)) < \delta/2, \quad \text{dist}(f^{n_2}(r), p_0) < \delta/2.$$

Hence the sequence

$$\{q_n\}_{n=0}^{S+n_2-n_1+N} = \{z_0, f(z_0), \dots, f^{S-1}(z_0), f^{n_1}(r), \dots, f^{n_2-1}(r), p_0, \dots, p_N\}.$$

Define η by (6) and $L = S + K + N + 1$. Arguing similarly to the proof of Theorem 1 we conclude that

$$P^{z_0, d}(z_n \in B(\delta, q_n), n \in [0, S + n_2 - n_1 + N]) \geq \eta^{n_2-n_1+N} \geq \eta^{K+S+N}.$$

Again arguing similarly to Theorem 1 and using inclusion (11) we conclude that for any $z_0 \in W$ the equality

$$P^{z_0, d}(\{z_n\} \text{ is } \varepsilon\text{-shadowable}) = 0$$

holds. Combining the latter with inclusion (10) we conclude that

$$P^{y_0, d}(\{y_n\} \text{ is } \varepsilon\text{-shadowable}) = 0,$$

which completes the proof of Theorem 2. □

Acknowledgement. Research of the author is supported by the Russian Science Foundation grant 14-21-00035, Saint-Petersburg State University research grant 6.38.223.2014, and RFBR 15-01-03797a.

References

- [1] Abdenur F. Attractors of generic diffeomorphisms are persistent. *Non-linearity* 16 (2003), no. 1, 301311.
- [2] Abdenur F., Diaz L. J. Pseudo-orbit shadowing in the C^1 topology. – *Discrete Contin. Dyn. Syst.*, 7 (2003), 223-245.

- [3] Anosov D. V. On a class of invariant sets of smooth dynamical systems. *Proc. 5th Int. Conf. on Nonlin. Oscill.* 2 (1970), 39-45.
- [4] Bowen R. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*. Lecture Notes Math., vol. 470, Springer, Berlin, 1975.
- [5] Hammel S. M., Yorke J. A., Grebogi C. Do numerical orbits of chaotic dynamical processes represent true orbits. *J. of Complexity* **3** (1987), 136-145.
- [6] Hammel S. M., Yorke J. A., Grebogi C. Numerical orbits of chaotic processes represent true orbits. *Bulletin of the American Mathematical Society* 19 (1988), 465-469.
- [7] Hasselblatt B., Katok A. *Introduction to the modern theory of dynamical systems*. Encyclopedia of Mathematics and its Applications, 54. Cambridge University Press, Cambridge, 1995.
- [8] Hasselblatt B., Katok A. *A first course in dynamics. With a panorama of recent developments*. Cambridge University Press, New York, 2003.
- [9] Leonov G. A. *Strange Attractors and Classical Stability Theory*. St. Petersburg University Press, 2009.
- [10] Palmer K. *Shadowing in Dynamical Systems. Theory and Applications*. Kluwer, Dordrecht, 2000.
- [11] Pilyugin S. Yu. *Shadowing in Dynamical Systems*. Lecture Notes Math., vol. 1706, Springer, Berlin, 1999.
- [12] Pilyugin S. Yu. *Spaces of dynamical systems*. De Gruyter Studies in Mathematical Physics, 3. De Gruyter, Berlin, 2012.
- [13] Pilyugin S. Yu. Theory of pseudo-orbit shadowing in dynamical systems. *Diff. Eqs.* 47 (2011), 1929–1938.
- [14] Pilyugin S. Yu., Rodionova A. A., Sakai K. Orbital and weak shadowing properties. – *Discrete Contin. Dyn. Syst.*, 9 (2003), 287-308.
- [15] Pilyugin S. Yu., Tikhomirov S. B. Lipschitz Shadowing implies structural stability. *Nonlinearity* 23 (2010), 2509-2515

- [16] Sakai K., Pseudo orbit tracing property and strong transversality of diffeomorphisms of closed manifolds. *Osaka J. Math.*, **31** (1994), 373-386.
- [17] Tikhomirov S., Holder Shadowing on finite intervals. *Ergodic Theory Dynam. Systems*, 35 (2015), no. 6, 2000-2016.
- [18] Tikhomirov S., Shadowing in linear skew products. *J. Math. Sci. (N.Y.)*, 209 (2015), no. 6, 979-987.
- [19] Yuan G., Yorke J., An open set of maps for which every point is absolutely nonshadowable. *Proc. Amer. Math. Soc.* 128 (2000), 909-918.