

**Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig**

**Energy identity for a class of
approximate Dirac-harmonic maps from
surfaces with boundary**

(revised version: June 2017)

by

Jürgen Jost, Lei Liu, and Miaomiao Zhu

Preprint no.: 11

2017



ENERGY IDENTITY FOR A CLASS OF APPROXIMATE DIRAC-HARMONIC MAPS FROM SURFACES WITH BOUNDARY

JÜRGEN JOST, LEI LIU, AND MIAOMIAO ZHU

ABSTRACT. For a sequence of coupled fields $\{(\phi_n, \psi_n)\}$ from a compact Riemann surface M with smooth boundary to a general compact Riemannian manifold with uniformly bounded energy and satisfying the Dirac-harmonic system up to some uniformly controlled error terms, we show that the energy identity holds during a blow-up process near the boundary. As an application to the heat flow of Dirac-harmonic maps from surfaces with boundary, when such a flow blows up at infinite time, we obtain an energy identity.

1. INTRODUCTION

This paper is a contribution to the study of coupled field equations on Riemann surfaces, merging the theory of harmonic maps from surfaces with a mathematical version of the nonlinear supersymmetric of quantum field theory. The corresponding action functional couples a term involving what is called the energy of a map from a surface to some Riemannian manifold with a Dirac action for a nonlinear spinor field. The solutions of the resulting Euler-Lagrange equations are called Dirac-harmonic maps [2]. While they share many properties with harmonic maps, their analysis is much more subtle, because the Dirac action is not bounded from below. Therefore, standard variational methods do not apply to show the existence of solutions under general conditions. As an alternative, a new type of mixed parabolic-elliptic has been introduced [3] and further investigated [11] in order to develop new tools for the existence problem. The existence problem is still not fully solved. In order to make progress, results about the behavior at singularities that are known and classical for harmonic maps need to be extended to the Dirac-harmonic case. This is where the contribution of the present paper lies. We study the blow-up process and show a so-called energy identity, that is, all the energy that is removed from the map gets transferred to the bubbles that represent the singularity. In fact, we study this at the boundary, because boundary value problems currently offer the situation where the existence theory is best developed and most promising. In order that our results be applicable to the parabolic case, we have to consider approximate solutions, that is, fields that satisfy the Euler-Lagrange equations up to some controlled error term. This naturally makes the bubbling analysis more difficult.

We now fix the technical setting to describe our results in more precise terms. Let (M, h) be a compact Riemann surface with smooth boundary ∂M , equipped with a Riemannian

Date: June 21, 2017.

2010 Mathematics Subject Classification. 53C43 58E20.

Key words and phrases. Dirac-harmonic maps, approximate Dirac-harmonic maps, Dirac-harmonic map flow, energy identity, boundary blow-up.

The third author was supported in part by National Science Foundation of China (No. 11601325).

metric h and with a fixed spin structure, ΣM be the spinor bundle over M and $\langle \cdot, \cdot \rangle_{\Sigma M}$ be the metric on ΣM induced by the Riemannian metric h . Choosing a local orthonormal basis $e_\alpha, \alpha = 1, 2$ on TM , the usual Dirac operator is defined as $\not{D} := e_\alpha \cdot \nabla_{e_\alpha}$, where ∇ is the spin connection on ΣM , \cdot is the Clifford multiplication, which satisfies the skew-adjointness property

$$\langle X \cdot \psi_1, \psi_2 \rangle_{\Sigma M} = -\langle \psi_1, X \cdot \psi_2 \rangle_{\Sigma M}$$

for any $X \in \Gamma(TM)$, $\psi_i \in \Gamma(\Sigma M)$, $i = 1, 2$.

Let ϕ be a smooth map from M to another compact Riemannian manifold (N, g) with dimension $n \geq 2$. Let ϕ^*TN be the pull-back bundle of TN by ϕ and then we get the twisted bundle $\Sigma M \otimes \phi^*TN$. Naturally, there is a metric $\langle \cdot, \cdot \rangle_{\Sigma M \otimes \phi^*TN}$ on $\Sigma M \otimes \phi^*TN$ which is induced from the metrics on ΣM and ϕ^*TN . Also we have a natural connection $\tilde{\nabla}$ on $\Sigma M \otimes \phi^*TN$ which is induced from the connections on ΣM and ϕ^*TN . Let ψ be a section of the bundle $\Sigma M \otimes \phi^*TN$. In local coordinates, it can be written as

$$\psi = \psi^i \otimes \partial_{y^i}(\phi),$$

where each ψ^i is a usual spinor on M and ∂_{y^i} is the nature local basis on N . Then $\tilde{\nabla}$ becomes

$$(1.1) \quad \tilde{\nabla} \psi = \nabla \psi^i \otimes \partial_{y^i}(\phi) + (\Gamma_{jk}^i \nabla \phi^j) \psi^k \otimes \partial_{y^i}(\phi),$$

where Γ_{jk}^i are the Christoffel symbols of the Levi-Civita connection of (N, g) . The Dirac operator along the map ϕ is defined by $\not{D}\psi := e_\alpha \cdot \tilde{\nabla}_{e_\alpha} \psi$.

An important factor that will enable us to utilize tools from complex analysis is that the usual Dirac operator \not{D} on a surface can be seen as the Cauchy-Riemann operator. Consider \mathbb{R}^2 with the Euclidean metric $dx^2 + dy^2$. Let $e_1 = \frac{\partial}{\partial x}$ and $e_2 = \frac{\partial}{\partial y}$ be the standard orthonormal frame. A spinor field is simply a map $\psi : \mathbb{R}^2 \rightarrow \Delta_2 = \mathbb{C}^2$, and the action of e_1 and e_2 on spinors can be identified with multiplication with matrices

$$e_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

If $\psi := \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} : \mathbb{R}^2 \rightarrow \mathbb{C}^2$ is a spinor field, then the Dirac operator is

$$(1.2) \quad \not{D}\psi = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} \frac{\partial \psi_1}{\partial x} \\ \frac{\partial \psi_2}{\partial x} \end{pmatrix} + \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \begin{pmatrix} \frac{\partial \psi_1}{\partial y} \\ \frac{\partial \psi_2}{\partial y} \end{pmatrix} = 2 \begin{pmatrix} \frac{\partial \psi_2}{\partial \bar{z}} \\ -\frac{\partial \psi_1}{\partial z} \end{pmatrix},$$

where

$$\frac{\partial}{\partial z} = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).$$

For more details on spin geometry and Dirac operators, one can refer to [15].

We consider the following functional

$$\begin{aligned} L(\phi, \psi) &= \int_M (|d\phi|^2 + \langle \psi, \not{D}\psi \rangle_{\Sigma M \otimes \phi^*TN}) dvol \\ &= \int_M \left(g_{ij}(\phi) h^{\alpha\beta} \frac{\partial \phi^i}{\partial x^\alpha} \frac{\partial \phi^j}{\partial x^\beta} + g_{ij}(\phi) \langle \psi^i, \not{D}\psi^j \rangle_{\Sigma M} \right) dvol. \end{aligned}$$

The functional $L(\phi, \psi)$ is conformally invariant (see [2]). That is, for any conformal diffeomorphism $f : M \rightarrow M$, setting

$$\tilde{\phi} = \phi \circ f \quad \text{and} \quad \tilde{\psi} = \lambda^{-1/2} \psi \circ f,$$

here λ is the conformal factor of the conformal map f , i.e. $f^*h = \lambda^2h$. Then $L(\tilde{\phi}, \tilde{\psi}) = L(\phi, \psi)$. Critical points (ϕ, ψ) of L are called Dirac-harmonic maps from M to N .

The Euler-Lagrange equations of the functional L are

$$(1.3) \quad (\Delta\phi^i + \Gamma_{jk}^i h^{\alpha\beta} \phi_\alpha^j \phi_\beta^k) \frac{\partial}{\partial y^i}(\phi(x)) = R(\phi, \psi),$$

$$(1.4) \quad \not{D}\psi = 0,$$

where $R(\phi, \psi)$ is defined by

$$R(\phi, \psi) = \frac{1}{2} R_{lij}^m(\phi(x)) \langle \psi^i, \nabla\phi^l \cdot \psi^j \rangle \frac{\partial}{\partial y^m}(\phi(x)).$$

Here R_{lij}^m stands for the Riemann curvature tensor of the target manifold (N, g) . One can refer to [1, 2].

By Nash's embedding theorem, we embed N isometrically into some Euclidean space \mathbb{R}^K . Then, the critical points (ϕ, ψ) satisfy the Euler-Lagrange equations

$$(1.5) \quad -\Delta\phi = A(\phi)(d\phi, d\phi) - \text{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)),$$

$$(1.6) \quad \not{D}\psi = \mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi),$$

where \not{D} is the usual Dirac operator, A is the second fundamental form of N in \mathbb{R}^K , and

$$\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi) := (\nabla\phi^i \cdot \psi^j) \otimes A(\partial_{y^i}, \partial_{y^j}),$$

$$\text{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)) := P(A(\partial_{y^l}, \partial_{y^j}); \partial_{y^i}) \text{Re}(\langle \psi^i, d\phi^l \cdot \psi^j \rangle).$$

Here $P(\xi; \cdot)$ denotes the shape operator satisfying $\langle P(\xi; X), Y \rangle = \langle A(X, Y), \xi \rangle$ for any $X, Y \in \Gamma(TN)$ and $\text{Re}(z)$ denotes the real part of $z \in \mathbb{C}$. We refer to [1, 2, 32, 4, 26, 10] for more details.

Before we state our main results, let us recall a definition of approximate Dirac-harmonic map in [14]. Denote

$$W^{2,2}(M, N) := \{ \phi \in W^{2,2}(M, \mathbb{R}^K) \text{ with } \phi(x) \in N \text{ for a.e. } x \in M \},$$

$$W^{1,4/3}(M, \Sigma M \otimes \phi^*TN) := \{ \psi \in W^{1,4/3}(M, \Sigma M \otimes \mathbb{R}^K) \text{ with } \psi(x) \in \Sigma M \otimes \phi^*TN \\ \text{for a.e. } x \in M \}.$$

A pair of fields $(\phi, \psi) \in W^{2,2}(M, N) \times W^{1,4/3}(M, \Sigma M \otimes \phi^*TN)$ is called an approximate Dirac-harmonic map from M to N with boundary data $(\varphi(x), \chi(x))$, if there exists a pair of fields $(\tau(\phi, \psi), h(\phi, \psi)) \in L^1(M)$ such that

$$(1.7) \quad \tau(\phi, \psi) = \Delta\phi + A(d\phi, d\phi) - \text{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)),$$

$$(1.8) \quad h(\phi, \psi) = \not{D}\psi - \mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi),$$

with the boundary data

$$(1.9) \quad \begin{cases} \phi(x) = \varphi(x), & \text{on } \partial M; \\ \mathcal{B}\psi(x) = \mathcal{B}\chi(x), & \text{on } \partial M, \end{cases}$$

where $\mathcal{B} = \mathcal{B}^\pm$ is the chiral boundary operator defined by

$$(1.10) \quad \mathcal{B}^\pm : L^2(\partial M, \Sigma M \otimes \phi^*TN|_{\partial M}) \rightarrow L^2(\partial M, \Sigma M \otimes \phi^*TN|_{\partial M})$$

$$(1.11) \quad \psi \mapsto \frac{1}{2} (Id \pm \vec{n} \cdot G) \cdot \psi,$$

where \vec{n} is the outward unit normal vector field on ∂M , $G = ie_1 \cdot e_2$ is the chiral operator defined using a local orthonormal frame $\{e_\alpha\}_{\alpha=1}^2$ on TM and satisfying:

$$(1.12) \quad G^2 = Id, \quad G^* = G, \quad \nabla G = 0, \quad G \cdot X = -X \cdot G,$$

for any $X \in \Gamma(TM)$. See e.g. [4, 3] for the notion of chiral boundary condition.

Therefore, (ϕ, ψ) is a Dirac-harmonic map if and only if $\tau(\phi, \psi) = h(\phi, \psi) = 0$.

Dirac-harmonic maps were introduced in [1, 2]. They are motivated by a model from quantum field theory, the supersymmetric sigma model [5, 9]. This subject generalizes the theory of harmonic maps and harmonic spinors. Similarly to the case of two dimensional harmonic maps, the conformal invariance of the energy functional L leads to non-compactness of Dirac-harmonic maps in dimension 2 and hence one needs to study their blow-up theory, as in [1, 30, 31, 21]. For the blow-up theory of a more general model, whose critical points are called Dirac-harmonic maps with curvature terms, see [10]. For approximate harmonic maps in dimension two, one can refer to e.g. [25, 8, 22, 23, 6, 24, 19, 20, 16, 28, 27] for the interior blow-up case and [12, 13, 7] for the boundary blow-up cases under various boundary constraints.

In order to study the blow-up behavior of the Dirac-harmonic map flow from surfaces with boundary considered in [3, 11], we introduced the notion of approximate Dirac-harmonic maps in [14] and proved the energy identity and no neck result in the interior blow-up case for a sequence of such maps. In general, this sequence might blow up at a boundary point. In this paper, we shall consider the case that the sequence blows up at the boundary and hence complete the blow-up picture of the Dirac-harmonic map flow.

Denote the energy of ϕ on $\Omega \subset M$ by

$$E(\phi; \Omega) = \int_{\Omega} |\nabla \phi|^2 dM,$$

the energy of ψ on $\Omega \subset M$ by

$$E(\psi; \Omega) = \int_{\Omega} |\psi|^4 dM,$$

and the energy of the pair (ϕ, ψ) on $\Omega \subset M$ by

$$E(\phi, \psi; \Omega) = \int_{\Omega} (|\nabla \phi|^2 + |\psi|^4) dM.$$

We shall often omit the domain M from the notation and simply write $E(\phi) = E(\phi; M)$, $E(\psi) = E(\psi; M)$ and $E(\phi, \psi) = E(\phi, \psi; M)$.

Based on the interior blow-up results for approximate Dirac-harmonic maps studied in [14], we state our first main result in this paper concerning the boundary blow-up case:

Theorem 1.1. *Consider a sequence of approximate Dirac-harmonic maps $(\phi_n, \psi_n) \in C^2(M, N) \times C^1(M, \Sigma M \otimes \phi^*TN)$ from a compact Riemann surface M with smooth boundary ∂M to a compact Riemannian manifold N satisfying*

$$E(\phi_n, \psi_n) + \|\tau(\phi_n, \psi_n)\|_{L^2} + \|h(\phi_n, \psi_n)\|_{L^4} \leq \Lambda,$$

and with boundary data

$$\phi_n|_{\partial M} = \varphi, \quad \mathcal{B}\psi_n|_{\partial M} = \mathcal{B}\chi,$$

where $\varphi \in C^{2+\alpha}(\partial M, N)$, $\chi \in C^{1+\alpha}(\partial M, \Sigma M \otimes \phi^*TN)$ for some $0 < \alpha < 1$. We assume $(\phi_n, \psi_n) \rightharpoonup (\phi, \psi)$ weakly in $W^{1,2}(M, N) \times L^4(M, \Sigma M \otimes \phi^*TN)$. Define the blow-up set

$$(1.13) \quad \mathcal{S} := \bigcap_{r>0} \left\{ x \in M \mid \liminf_{n \rightarrow \infty} \int_{D(x,r)} (|d\phi_n|^2 + |\psi_n|^4) \geq \frac{\bar{\epsilon}^2}{2} \right\},$$

where $\bar{\epsilon} > 0$ is some constant depending on N . Then \mathcal{S} is a (possibly empty) finite set $\{p_1, \dots, p_q, \dots, p_I\}$, where $1 \leq q \leq I$, $\{p_1, \dots, p_q\} \in M \setminus \partial M$, $\{p_{q+1}, \dots, p_I\} \in \partial M$. Moreover, a subsequence, still denoted by $\{(\phi_k, \psi_k)\}$, converges weakly in $W_{loc}^{2,2}(M \setminus \mathcal{S}) \times W_{loc}^{1,2}(M \setminus \mathcal{S})$ to (ϕ, ψ) and for each $i = 1, \dots, I$, there is a finite set of Dirac-harmonic spheres $(\sigma_i^l, \xi_i^l) : S^2 \rightarrow N$, $l = 1, \dots, L_i$, such that

$$(1.14) \quad \lim_{k \rightarrow \infty} E(\phi_k) = E(\phi) + \sum_{i=1}^I \sum_{l=1}^{L_i} E(\sigma_i^l),$$

$$(1.15) \quad \lim_{k \rightarrow \infty} E(\psi_k) = E(\psi) + \sum_{i=1}^I \sum_{l=1}^{L_i} E(\xi_i^l),$$

and the image $\phi(M \setminus \partial M) \cup_{i=1}^q \bigcup_{l=1}^{L_i} (\sigma_i^l(S^2))$ is a connected set.

Remark 1.2. In Theorem 1.1, for those Dirac-harmonic spheres splitting off at the interior blow-up points, i.e. $(\sigma_i^l, \xi_i^l) : S^2 \rightarrow N$, $i = 1, \dots, q$; $l = 1, \dots, L_i$, we know that the image of the map parts σ_i^l , $i = 1, \dots, q$; $l = 1, \dots, L_i$, are connected to the map part ϕ of the base field (ϕ, ψ) in the target manifold; this is proved in [14]. However, for those Dirac-harmonic spheres splitting off at the boundary blow-up points, i.e. $(\sigma_i^l, \xi_i^l) : S^2 \rightarrow N$, $i = q+1, \dots, I$; $l = 1, \dots, L_i$, it is not clear whether the images of the map parts σ_i^l , $i = q+1, \dots, I$; $l = 1, \dots, L_i$ have the same property.

To prove the energy quantization result near the boundary in Theorem 1.1, we shall follow the general blow-up scheme developed for harmonic map type problems, however, the proofs in this case are subtle and there are new difficulties arising when carrying out the neck analysis. Firstly, the method of the three circle type theorem used in the interior case in [14] can not be applied to the boundary case and we need to apply certain integration argument to show the no neck energy property. Secondly, we need to establish a new Pohozaev type identity for approximate Dirac-harmonic maps from surfaces with boundary (see Lemma 2.2) which requires some algebraic property for the spinors, see (2.10). Moreover, we succeed in driving a new Pohozaev type estimate (see Corollary 2.5) by showing that some terms in

the Pohozaev type identity are actually having exponential decay, which are crucial in the proof of the above theorem. Finally, we would like to remark that the bubbling analysis at the boundary is more complicated than in the interior case and here we follow the scheme developed for approximate harmonic maps in [12, 13].

With the help of Theorem 1.1, we can now study the asymptotic behavior at infinite time for the Dirac-harmonic map flow in dimension 2.

The notion of Dirac-harmonic map flow was introduced in [3]. In this flow, one seeks a pair of fields $(\phi, \psi) : M \times [0, \infty) \rightarrow N \times (\Sigma M \otimes \phi^*TN)$ that solves

$$(1.16) \quad \begin{cases} \partial_t \phi = \tau(\phi) - Re(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)), & \text{in } M \times (0, \infty); \\ \not\partial \psi = \mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi), & \text{in } M \times (0, \infty). \end{cases}$$

with the following boundary-initial data:

$$(1.17) \quad \begin{cases} \phi(x, t) = \varphi(x), & \text{on } \partial M \times [0, \infty); \\ \phi(x, 0) = \phi_0(x), & \text{in } M; \\ \mathcal{B}\psi(x, t) = \mathcal{B}\chi(x), & \text{on } \partial M \times [0, \infty); \\ \phi_0(x) = \varphi(x), & \text{on } \partial M, \end{cases}$$

where $\tau(\phi) := \Delta\phi + A(\phi)(\nabla\phi, \nabla\phi)$ is the tension field of ϕ , M is a compact spin Riemannian manifold with smooth boundary ∂M and of dimension $\dim M \geq 2$ and $\phi_0 \in W^{1,2}(M, N)$, $\varphi \in C^{2+\alpha}(\partial M, N)$, $\chi \in C^{1+\alpha}(\partial M, \Sigma M \otimes \varphi^*TN)$ are given data. The short-time existence for the above flow (1.16) (1.17) was proved in [3].

When M is a surface, it was shown in [11] that there exists a unique global weak solution to (1.16) with initial-boundary data (1.17) under some smallness assumption for $\|\phi_0\|_{H^1} + \|\mathcal{B}\chi\|_{L^2}$, which has at most finitely many singular times and enjoys the following property:

$$(1.18) \quad E(\phi(t), \psi(t); M) + \int_{M^t} |\partial_t \phi|^2 dx dt \leq C(M, E(\phi_0), \|\mathcal{B}\chi\|_{L^2(\partial M)}), \quad \forall 0 \leq t < \infty.$$

It follows from (1.18) that there exists a sequence $t_n \uparrow \infty$ such that

$$(\phi_n, \psi_n) := (\phi(\cdot, t_n), \psi(\cdot, t_n)) \in C^{2+\alpha}(M, N) \times C^{1+\alpha}(M, \Sigma M \otimes \phi^*TN)$$

is a sequence of approximate Dirac-harmonic maps with boundary-data (φ, χ) and satisfying

$$h(\phi_n, \psi_n) = 0$$

and

$$\tau(\phi_n, \psi_n) := \partial_t \phi(\cdot, t_n) \quad \text{with} \quad \|\tau(\phi_n, \psi_n)\|_{L^2} \rightarrow 0.$$

When such a flow blows up at infinite time and at interior points, it was proved in [14] that an energy identity and no neck property hold during the blow-up process. In this paper, as an immediate corollary of Theorem 1.1 and as a complement of the blow-up picture at infinite time of such a flow given in [14], we obtain

Theorem 1.3. *Let M be a compact spin Riemann surface with smooth boundary ∂M . Let $\phi_0 \in H^1(M, N)$, $\varphi \in C^{2+\alpha}(\partial M, N)$, $\chi \in C^{1+\alpha}(\partial M, \Sigma M \otimes \varphi^*TN)$. Let $(\phi, \psi) : M \times [0, \infty) \rightarrow N \times (\Sigma M \otimes \phi^*TN)$ be a global weak solution of (1.16) and (1.17), which has*

finitely many singular times and satisfies (1.18). Then there exist $t_n \uparrow \infty$, a Dirac-harmonic map $(\phi_\infty, \psi_\infty) \in C^{2+\alpha}(M, N) \times C^{1+\alpha}(M, \Sigma M \otimes \phi_\infty^* TN)$ with boundary data $\phi_\infty|_{\partial M} = \varphi$ and $\mathcal{B}\psi_\infty|_{\partial M} = \mathcal{B}\chi$, nonnegative integer I and a possibly empty set with at most finitely many points $\{p_1, \dots, p_q, \dots, p_I\} \subset M$, where $1 \leq q \leq I$, $\{p_1, \dots, p_q\} \in M \setminus \partial M$, $\{p_{q+1}, \dots, p_I\} \in \partial M$ such that

$$(1) (\phi_n, \psi_n) := (\phi(\cdot, t_n), \psi(\cdot, t_n)) \rightharpoonup (\phi_\infty, \psi_\infty) \text{ in } W^{1,2}(M, N) \times L^4(M, \Sigma M \times \phi_\infty^* TN);$$

$$(2) (\phi_n, \psi_n) \rightarrow (\phi_\infty, \psi_\infty) \text{ in } W_{loc}^{1,2}(M \setminus \{p_1, \dots, p_I\}) \times L_{loc}^4(M \setminus \{p_1, \dots, p_I\});$$

(3) For $1 \leq i \leq I$, there exist a positive integer L_i and L_i nontrivial Dirac-harmonic spheres $(\sigma_i^l, \xi_i^l) : S^2 \rightarrow N$, $i = 1, \dots, I$; $l = 1, \dots, L_i$ such that

$$(1.19) \quad \lim_{n \rightarrow \infty} E(\phi_n) = E(\phi_\infty) + \sum_{i=1}^I \sum_{l=1}^{L_i} E(\sigma_i^l),$$

$$(1.20) \quad \lim_{n \rightarrow \infty} E(\psi_n) = E(\psi_\infty) + \sum_{i=1}^I \sum_{l=1}^{L_i} E(\xi_i^l).$$

and the image $\phi_\infty(M \setminus \partial M) \cup_{i=1}^q \cup_{l=1}^{L_i} (\sigma_i^l(S^2))$ is a connected set.

This paper is organized as follows. In Section 2, we extend some basic lemmas to the boundary case, such as small energy regularity, Pohozaev's identity and removable singularity. Then, we recall some known results which will be used in this paper. In Section 3, we prove our main Theorem 1.1.

Notations: We denote $\mathbb{R}_+^2 = \{(x, y) \in \mathbb{R}^2 | y \geq 0\}$, $D_r(x) = \{y \in \mathbb{R}^2 | |y - x| \leq r\}$, $D_r^+(x) = D_r(x) \cap \mathbb{R}_+^2$, $\partial^+ D_r^+(x) = \partial D_r(x) \cap \mathbb{R}_+^2$, $\partial^0 D_r^+(x) = D_r(x) \cap \partial \mathbb{R}_+^2$.

For simplicity, we also denote $D_r(0)$, $D_r^+(0)$, $D_1(0)$, $D_1^+(0)$ as D_r , D_r^+ , D , D^+ respectively.

2. SOME BASIC LEMMAS

In this section, we will prove some basic lemmas and recall some known results which will be used in this paper.

By the standard elliptic theory, there exists a unique solution $u \in C^{2+\alpha}(M, \mathbb{R}^K)$ of

$$\begin{cases} \Delta u = 0, & \text{in } M, \\ u = \varphi, & \text{on } \partial M, \end{cases}$$

satisfying

$$\|u\|_{C^{2+\alpha}(M)} \leq C(\alpha, M) \|\varphi\|_{C^{2+\alpha}(\partial M)}.$$

For simplicity of notation, in the sequel, we will also denote this solution as φ .

Firstly, we prove a small energy regularity theorem for the boundary case. For similar results for approximate harmonic maps, one can refer to the main estimate 3.2 in [25] and Lemma 2.1 in [6] for the interior case and one can also refer to Lemma 4.1 in [12], Lemma 2.4 in [13] for various boundary cases.

Theorem 2.1. *There is a small constant $\epsilon_0 > 0$ depending only on p, q and N , such that if $(\phi, \psi) \in W^{2,p}(D^+, N) \times W^{1,q}(D^+, \Sigma D^+ \otimes \phi^* TN)$ is an approximate Dirac-harmonic map from the upper unit disc $D^+ \subset \mathbb{R}^2$ to a compact Riemannian manifold (N, g) with $\tau(\phi, \psi) \in L^p$ and $h(\phi, \psi) \in L^q$ for some $1 < p \leq 2$ and some $\frac{4}{3} < q \leq 2$, and with boundary data (1.9), satisfying*

$$(2.1) \quad E(\phi, \psi; D^+) = \int_{D^+} (|d\phi|^2 + |\psi|^4) dx < (\epsilon_0)^2,$$

then

$$\begin{aligned} \|\phi - \bar{\varphi}\|_{W^{2,p}(D^+_{\frac{1}{2}})} &\leq C(\|d\phi\|_{L^p(D^+)} + \|\psi\|_{L^{2p}(D^+)} + \|\tau\|_{L^p(D^+)} + \|\nabla\varphi\|_{W^{1,p}(D^+)}), \\ \|\psi\|_{W^{1,q}(D^+_{\frac{1}{2}})} &\leq C(\|\psi\|_{L^q(D^+)} + \|h\|_{L^q(D^+)} + \|\mathcal{B}\chi\|_{W^{1-1/q,q}(\partial^0 D^+)}), \end{aligned}$$

where $\bar{\varphi} := \int_{\partial^0 D^+_{1/2}} \varphi$ and $C > 0$ is a constant depending only on $p, q, N, \|\varphi\|_{C^2}, \|\chi\|_{C^1}$.

Moreover, by the Sobolev embedding $W^{2,p}(\mathbb{R}^2) \subset C^0(\mathbb{R}^2)$, we have

$$(2.2) \quad \|\phi\|_{Osc(D^+_{1/2})} = \sup_{x,y \in D^+_{1/2}} |\phi(x) - \phi(y)| \leq C(\|\nabla\phi\|_{L^2(D^+)} + \|\tau(u)\|_{L^p(D^+)} + \|\nabla\varphi\|_{W^{1,p}(D^+)}).$$

Proof. Without loss of generality, we assume $\int_{\partial^0 D^+_{\frac{1}{2}}} \varphi = 0$.

Choosing a cut-off function $\eta \in C_0^\infty(D^+)$ satisfying $0 \leq \eta \leq 1, \eta|_{D^+_{3/4}} \equiv 1, |\nabla\eta| + |\nabla^2\eta| \leq C$, by elliptic theory of first order elliptic operator, for any $1 < q < 2$, we have

$$\begin{aligned} \|\eta\psi\|_{W^{1,q}(D^+)} &\leq C(\|\mathcal{D}(\eta\psi)\|_{L^q(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/q,q}(\partial^0 D^+)}) \\ &\leq C(\|\nabla\eta \cdot \psi + \eta\mathcal{D}\psi\|_{L^q(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/q,q}(\partial^0 D^+)}) \\ &\leq C(\|\psi\|_{L^q(D^+)} + \|d\phi\|\|\eta\psi\|_{L^q(D^+)} + \|h\|_{L^q(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/q,q}(\partial^0 D^+)}) \\ &\leq C\|d\phi\|_{L^2(D^+)}\|\eta\psi\|_{L^{\frac{2q}{2-q}}(D^+)} + C(\|\psi\|_{L^q(D^+)} + \|h\|_{L^q(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/q,q}(\partial^0 D^+)}) \\ &\leq C\epsilon_0\|\eta\psi\|_{L^{\frac{2q}{2-q}}(D^+)} + C(\|\psi\|_{L^q(D^+)} + \|h\|_{L^q(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/q,q}(\partial^0 D^+)}). \end{aligned}$$

Taking $\epsilon_0 > 0$ sufficiently small, by Sobolev embedding, we get

$$(2.3) \quad \|\eta\psi\|_{L^{\frac{2q}{2-q}}(D^+)} \leq \|\eta\psi\|_{W^{1,q}(D^+)} \leq C(\|\psi\|_{L^q(D^+)} + \|h\|_{L^q(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/q,q}(\partial^0 D^+)}).$$

Computing directly, we obtain

$$\begin{aligned} |\Delta(\eta\phi)| &= |\eta\Delta\phi + 2\nabla\eta\nabla\phi + \phi\Delta\eta| \\ &\leq C(|\phi| + |d\phi| + |d\phi|\|\eta d\phi\| + |\psi|^2|\eta d\phi| + |\tau|) \\ &\leq C(|d\phi| + |\psi|^2)|d(\eta\phi)| + C(|\phi| + |d\phi| + |\psi|^2 + |\tau|). \end{aligned}$$

By standard elliptic estimates and Poincaré's inequality, for any $1 < p < 2$, we have

$$\begin{aligned}
\|\eta\phi\|_{W^{2,p}(D^+)} &\leq C(\| |d\phi| + |\psi|^2 \|_{L^p(D^+)} \|d(\eta\phi)\|_{L^p(D^+)} + C(\|d\phi\|_{L^p(D^+)} + \|\psi\|^2_{L^p(D^+)}) \\
&\quad + \|\tau\|_{L^p(D^+)} + \|\varphi\|_{W^{2,p}(D^+)}) \\
&\leq C\|d(\eta\phi)\|_{L^{\frac{2p}{2-p}}(D^+)} (\|d\phi\|_{L^2(D^+)} + \|\psi\|^2_{L^4(D^+)}) + C(\|d\phi\|_{L^p(D^+)} \\
&\quad + \|\psi\|^2_{L^{2p}(D^+)} + \|\tau\|_{L^p(D^+)} + \|\varphi\|_{W^{2,p}(D^+)}) \\
&\leq C\epsilon_0\|d(\eta\phi)\|_{L^{\frac{2p}{2-p}}(D^+)} + C(\|d\phi\|_{L^p(D^+)} + \|\psi\|^2_{L^{2p}(D^+)} + \|\tau\|_{L^p(D^+)} \\
&\quad + \|\nabla\varphi\|_{W^{1,p}(D^+)}).
\end{aligned}$$

Taking $\epsilon_0 > 0$ sufficiently small, we have

$$\begin{aligned}
\|\nabla(\eta\phi)\|_{L^{\frac{2p}{2-p}}(D^+)} &\leq C\|\eta\phi\|_{W^{2,p}(D^+)} \\
(2.4) \quad &\leq C(\|d\phi\|_{L^p(D^+)} + \|\psi\|^2_{L^{2p}(D^+)} + \|\tau\|_{L^p(D^+)} + \|\nabla\varphi\|_{W^{1,p}(D^+)}).
\end{aligned}$$

So, we have proved the theorem in the case $1 < p < 2$, $4/3 < q < 2$.

For the case $p = 2$, $4/3 < q < 2$, taking $p = \frac{q}{2(q-1)} \in (1, 2)$ and $p = \frac{4}{3}$ in (2.4), by Sobolev embedding, we have

$$\begin{aligned}
&\|\nabla\phi\|_{L^4(D_{3/4}^+)} + \|\nabla\phi\|_{L^{\frac{2q}{3q-4}}(D_{3/4}^+)} \\
(2.5) \quad &\leq C(\|d\phi\|_{L^2(D^+)} + \|\psi\|^2_{L^4(D^+)} + \|\tau\|_{L^2(D^+)} + \|\nabla\varphi\|_{W^{1,2}(D^+)}).
\end{aligned}$$

By (2.3) and the $W^{2,2}$ -estimate for the Laplace operator, we obtain

$$\begin{aligned}
\|\phi\|_{W^{2,2}(D_{1/2}^+)} &\leq C(\|\Delta\phi\|_{L^2(D_{3/4}^+)} + \|\nabla\phi\|_{L^2(D^+)} + \|\nabla\varphi\|_{W^{1,2}(D^+)}) \\
&\leq C(\|\nabla\phi\|_{L^4(D_{3/4}^+)}^2 + \|\nabla\phi\|_{L^{\frac{2q}{3q-4}}(D_{3/4}^+)} \|\psi\|^2_{L^{\frac{q}{2-q}}(D_{3/4}^+)} + \|\nabla\phi\|_{L^2(D^+)}) \\
&\quad + \|\nabla\varphi\|_{W^{1,2}(D^+)}) \\
&\leq C(\|d\phi\|_{L^2(D^+)} + \|\psi\|_{L^4(D^+)} + \|\tau\|_{L^2(D^+)} + \|\nabla\varphi\|_{W^{1,2}(D^+)}).
\end{aligned}$$

For the case $q = 2$, $1 < p < 2$, taking $q = \frac{2p}{3p-2} \in (1, 2)$ in (2.3), we get

$$(2.6) \quad \|\psi\|_{L^{\frac{p}{p-1}}(D_{3/4}^+)} \leq C(\|\psi\|_{L^2(D^+)} + \|h\|_{L^2(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/2,2}(\partial^0 D^+)}).$$

By (2.4) and $W^{1,2}$ -estimates for the Dirac operator, we arrive at

$$\begin{aligned}
\|\psi\|_{W^{1,2}(D_{1/2}^+)} &\leq C(\|\not\partial\psi\|_{L^2(D_{3/4}^+)} + \|\psi\|_{L^4(D_{3/4}^+)} + \|\mathcal{B}\chi\|_{W^{1-1/2,2}(\partial^0 D^+)}) \\
&\leq C(\|\nabla\phi\|_{L^{\frac{2p}{2-p}}(D_{3/4}^+)} \|\psi\|_{L^{\frac{p}{p-1}}(D_{3/4}^+)} + \|\psi\|_{L^4(D_{3/4}^+)} + \|\mathcal{B}\chi\|_{W^{1-1/2,2}(\partial^0 D^+)}) \\
&\leq C(\|\psi\|_{L^2(D^+)} + \|h\|_{L^2(D^+)} + \|\mathcal{B}\psi\|_{W^{1-1/2,2}(\partial^0 D^+)}).
\end{aligned}$$

For the case $p = q = 2$, taking $q = \frac{8}{5}$ in (2.3) and $p = \frac{4}{3}$ in (2.4), we will obtain a $L^8(D_{3/4}^+)$ -bound for ψ and a $L^4(D_{3/4}^+)$ bound for $\nabla\phi$. Then one can apply the $W^{2,2}$ -boundary estimate for the Laplace operator and the $W^{1,2}$ -boundary estimate for the Dirac operator to get the conclusion of the theorem. \square

Next we shall derive a Pohozaev type identity for approximate Dirac-harmonic maps with boundary data, extending the interior case given in Lemma 2.3 in [14]. For corresponding results for two dimensional approximate harmonic maps, one can refer to Lemma 2.4 [19] for the interior case and refer to Lemma 4.3 in [12] and Lemma 2.5 in [13] for various boundary cases.

Lemma 2.2. (Pohozaev type identity) *Let $\Omega \subset \mathbb{R}^2$ be a bounded smooth domain. If $D^+ \subset \Omega \subset \mathbb{R}_+^2$ and $(\phi, \psi) \in C^2(\Omega, N) \times C^1(\Omega, \Sigma\Omega \otimes \phi^*TN)$ is an approximate Dirac-harmonic map with boundary data (1.9) on $\partial^0\Omega$, then for any $0 < t < \frac{1}{2}$, we have*

$$\begin{aligned}
(2.7) \quad t \int_{\partial^+ D_t^+} (|\phi_r|^2 - \frac{1}{2}|\nabla\phi|^2) &= \frac{1}{2} \int_{\partial^+ D_t^+} \langle \psi, r^{-1}\partial_\theta \cdot \psi_\theta \rangle - \frac{1}{2} \int_{D_t^+} \langle \psi, \not{D}\psi \rangle dx - \operatorname{Re} \int_{D_t^+} \langle \not{D}\psi, r\psi_r \rangle dx \\
&+ \int_{D_t^+} r(\phi - \varphi)_r \tau dx + \frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle \\
&+ \int_{\partial^+ D_t^+} r\phi_r \varphi_r - \int_{D_t^+} \nabla\phi(\nabla\varphi + r\nabla\varphi_r) dx \\
&+ \int_{D_t^+} \langle r\varphi_r, A(\phi)(d\phi, d\phi) - \operatorname{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)) \rangle dx,
\end{aligned}$$

where (r, θ) are polar coordinates in D centered at 0, $\phi_r = \frac{\partial\phi}{\partial r}$, $\psi_r = \tilde{\nabla}_{\frac{\partial}{\partial r}}\psi$ and $\psi_\theta = \tilde{\nabla}_{\frac{\partial}{\partial\theta}}\psi$.

Before we prove this lemma, let us recall two basic properties for Dirac operators and spinors with chiral boundary constraint,

Lemma 2.3 (Proposition 3.1 in [4]). *For any $\psi, \omega \in W^{1,3/4}(M, \Sigma M \otimes \phi^*TN)$ satisfying*

$$\mathcal{B}\psi|_{\partial M} = \mathcal{B}\omega|_{\partial M} = 0,$$

we have

$$(2.8) \quad \langle \vec{n} \cdot \psi, \omega \rangle = 0 \text{ on } \partial M,$$

where \vec{n} is the unit normal vector field on ∂M .

Lemma 2.4 (Proposition 3.2 in [4]). *For any $\psi, \omega \in W^{1,3/4}(M, \Sigma M \otimes \phi^*TN)$, we have*

$$(2.9) \quad \int_M \langle \psi, \not{D}\omega \rangle dx = \int_M \langle \not{D}\psi, \omega \rangle dx - \int_{\partial M} \langle \vec{n} \cdot \psi, \omega \rangle$$

where $\langle \psi, \omega \rangle := h_{ij}\langle \psi^i, \omega^j \rangle$.

Proof of Lemma 2.2: Multiplying the equation (1.7) by $r(\phi - \varphi)_r$ and integrating over D_t^+ , noting the fact that $r\partial_r\phi = x^\beta\partial_\beta\phi$ and recalling Proposition 2.2 in [14] that

$$(2.10) \quad \langle \psi, \tilde{\nabla}_{\frac{\partial}{\partial x^\beta}}(\not{D}\psi) \rangle = 2\langle \operatorname{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)), \nabla_{\frac{\partial}{\partial x^\beta}}\phi \rangle + \langle \psi, \not{D}\tilde{\nabla}_{\frac{\partial}{\partial x^\beta}}\psi \rangle,$$

we get

$$\begin{aligned}
\int_{D_t^+} r(\phi - \varphi)_r \tau dx &= \int_{D_t^+} r(\phi - \varphi)_r \Delta \phi dx - \int_{D_t^+} \langle r\phi_r, \operatorname{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)) \rangle dx \\
&\quad - \int_{D_t^+} \langle r\varphi_r, A(\phi)(d\phi, d\phi) - \operatorname{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)) \rangle dx \\
&= \int_{D_t^+} r(\phi - \varphi)_r \Delta \phi dx + \frac{1}{2} \int_{D_t^+} \langle x^\beta \psi, \mathcal{D}\psi_\beta \rangle dx - \frac{1}{2} \int_{D_t^+} \langle x^\beta \psi, \tilde{\nabla}_{\partial_\beta} \mathcal{D}\psi \rangle dx \\
&\quad - \int_{D_t^+} \langle r\varphi_r, A(\phi)(d\phi, d\phi) - \operatorname{Re}(P(\mathcal{A}(d\phi(e_\alpha), e_\alpha \cdot \psi); \psi)) \rangle dx \\
&:= \text{II} + \text{III} + \text{IIII} + \text{IV}.
\end{aligned}$$

On one hand, by integrating by parts, we have

$$\begin{aligned}
\text{II} &= \int_{\partial^+ D_t^+} r|\phi_r|^2 - \int_{\partial^+ D_t^+} r\phi_r \varphi_r - \int_{D_t^+} \nabla \phi \nabla (r(\phi - \varphi)_r) dx \\
&= \int_{\partial^+ D_t^+} r|\phi_r|^2 - \int_{\partial^+ D_t^+} r\phi_r \varphi_r - \int_{D_t^+} \nabla \phi \nabla (\phi - \varphi) dx - \frac{1}{2} \int_{D_t^+} r \partial_r |\nabla \phi|^2 dx \\
&\quad + \int_{D_t^+} r \nabla \phi \nabla \varphi_r dx \\
&= t \int_{\partial^+ D_t^+} (|\phi_r|^2 - \frac{1}{2} |\nabla \phi|^2) - \int_{\partial^+ D_t^+} r\phi_r \varphi_r + \int_{D_t^+} \nabla \phi (\nabla \varphi + r \nabla \varphi_r) dx,
\end{aligned}$$

where the last equality follows from the fact that

$$\begin{aligned}
-\frac{1}{2} \int_{D_t^+} r \partial_r |\nabla \phi|^2 dx &= -\frac{1}{2} \int_{\partial^+ D_t^+} \int_0^t r^2 \partial_r |\nabla \phi|^2 dr d\theta \\
&= -\frac{1}{2} \int_{\partial^+ D_t^+} t |\nabla \phi|^2 + \int_{D_t^+} |\nabla \phi|^2 dx.
\end{aligned}$$

On the other hand, by Lemma 2.4, we get

$$\begin{aligned}
2\text{III} &= \int_{D_t^+} \langle x^\beta \psi, \mathcal{D}\psi_\beta \rangle dx \\
&= \int_{D_t^+} \langle \mathcal{D}(x^\beta \psi), \psi_\beta \rangle dx - \int_{\partial^+ D_t^+} \langle \frac{\partial}{\partial r} \cdot x^\beta \psi, \psi_\beta \rangle + \int_{\partial^0 D_t^+} \langle \frac{\partial}{\partial x^2} \cdot x^\beta \psi, \psi_\beta \rangle \\
(2.11) \quad &= - \int_{D_t^+} \langle \psi, \mathcal{D}\psi \rangle dx + \int_{D_t^+} \langle \mathcal{D}\psi, r\psi_r \rangle dx + \int_{\partial^+ D_t^+} \langle \psi, r\partial_r \cdot \psi_r \rangle - \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle.
\end{aligned}$$

Integrating by parts, it follows that

$$\begin{aligned}
2\text{IIII} &= - \int_{D_t^+} \langle x^\beta \psi, \tilde{\nabla}_{\partial_\beta} \mathcal{D}\psi \rangle dx \\
&= - \int_{\partial^+ D_t^+} \langle r\psi, \mathcal{D}\psi \rangle dx + \int_{D_t^+} \langle \tilde{\nabla}_{\partial_\beta} (x^\beta \psi), \mathcal{D}\psi \rangle dx \\
(2.12) \quad &= 2 \int_{D_t^+} \langle \psi, \mathcal{D}\psi \rangle dx + \int_{D_t^+} \langle r\psi_r, \mathcal{D}\psi \rangle dx - \int_{\partial^+ D_t^+} \langle r\psi, \mathcal{D}\psi \rangle.
\end{aligned}$$

Thus, we have

$$\begin{aligned}
\text{III} + \text{IIII} &= \frac{1}{2} \int_{D_t^+} \langle \psi, \mathcal{D}\psi \rangle dx + Re \int_{D_t^+} \langle \mathcal{D}\psi, r\psi_r \rangle dx - \frac{1}{2} \int_{\partial^+ D_t^+} \langle \psi, r^{-1} \partial_\theta \cdot \psi_\theta \rangle \\
&\quad - \frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle.
\end{aligned}$$

Combining these estimates, we get (2.7). This finishes the proof of the lemma. \square

As a consequence of Lemma 2.2, we derive the following Pohozaev type estimate, which plays a key role in the proof of Theorem 1.1.

Corollary 2.5. (Pohozaev type estimate) *Under the assumption of Lemma 2.2, if*

$$E(\phi, \psi; D^+) + \|\tau(\phi, \psi)\|_{L^2} + \|h(\phi, \psi)\|_{L^4} \leq \Lambda,$$

then for any $0 < t < \frac{1}{4}$ and $0 < \varepsilon < \frac{1}{4}$, we have

$$\begin{aligned}
\int_{D_{2t}^+ \setminus D_t^+} (|\phi_r|^2 - \frac{1}{2} |\nabla \phi|^2) dx &\leq \varepsilon \int_{D_{2t}^+ \setminus D_t^+} \left| r^{-1} \frac{\partial \phi}{\partial \theta} \right|^2 dx + \frac{C}{\varepsilon} \int_{D_{2t}^+ \setminus D_t^+} |\psi|^4 dx \\
(2.13) \quad &+ C \int_{D_{2t}^+ \setminus D_t^+} \left| r^{-1} \frac{\partial \psi}{\partial \theta} \right|^{\frac{4}{3}} dx + C\sqrt{t},
\end{aligned}$$

where C is a positive constant depending only on Λ , N , $\|\varphi\|_{C^2}$, $\|\chi\|_{C^1}$.

Proof. Firstly, by equation (1.8) and elliptic theory, we have

$$\|\psi\|_{W^{1, \frac{4}{3}}(D_{\frac{1}{2}}^+)} \leq C(\|\nabla \phi\|_{L^2(D^+)} \|\psi\|_{L^4(D^+)} + \|h\|_{L^{\frac{4}{3}}(D^+)} + \|\mathcal{B}\chi\|_{W^{1/4, 4/3}(\partial^0 D^+)}) \leq C.$$

Thanks to Lemma 2.2, for any $0 < t < \frac{1}{2}$, we have

$$(2.14) \quad t \int_{\partial^+ D_t^+} (|\phi_r|^2 - \frac{1}{2} |\nabla \phi|^2) := \mathbb{I}_1 + \dots + \mathbb{I}_8.$$

Using Young's inequality and the fact that

$$(2.15) \quad \psi_r = \tilde{\nabla}_{\frac{\partial}{\partial r}} \psi = \frac{\partial \psi}{\partial r} + \psi^i \otimes A(d\phi(\frac{\partial}{\partial r}), \frac{\partial}{\partial y^i}),$$

where $\frac{\partial \psi}{\partial r} = (\frac{\partial \psi^1}{\partial r}, \dots, \frac{\partial \psi^K}{\partial r})$, we obtain

$$\begin{aligned}
\mathbb{I}_2 + \mathbb{I}_3 + \mathbb{I}_4 &\leq Ct(\|\psi\|_{L^4(D_t^+)} + \|\psi_r\|_{L^{\frac{4}{3}}(D_t^+)})\|h\|_{L^4(D_t^+)} \\
&\quad + Ct(\|\nabla\phi\|_{L^2(D_t^+)} + \|\nabla\varphi\|_{L^2(D_t^+)})\|\tau\|_{L^2(D_t^+)} \\
&\leq Ct(\|\psi\|_{L^4(D_t^+)} + \|\nabla\psi\|_{L^{\frac{4}{3}}(D_t^+)} + \|\nabla\phi\|_{L^2(D_t^+)})\|\psi\|_{L^4(D_t^+)} + Ct \\
(2.16) \quad &\leq Ct.
\end{aligned}$$

As for \mathbb{I}_5 , we have

$$\begin{aligned}
\mathbb{I}_5 &= \frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle \\
&= \frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi - \chi, \frac{\partial}{\partial x^2} \cdot r(\psi - \chi)_r \rangle + \frac{1}{2} \int_{\partial^0 D_t^+} \langle \chi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle \\
&\quad + \frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\chi_r \rangle - \frac{1}{2} \int_{\partial^0 D_t^+} \langle \chi, \frac{\partial}{\partial x^2} \cdot r\chi_r \rangle \\
(2.17) \quad &= \frac{1}{2} \int_{\partial^0 D_t^+} \langle \chi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle + \frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\chi_r \rangle - \frac{1}{2} \int_{\partial^0 D_t^+} \langle \chi, \frac{\partial}{\partial x^2} \cdot r\chi_r \rangle,
\end{aligned}$$

where the last equality follows from Lemma 2.3 which tells us that

$$\frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi - \chi, \frac{\partial}{\partial x^2} \cdot r(\psi - \chi)_r \rangle = 0.$$

Computing directly, we get

$$\begin{aligned}
\frac{1}{2} \int_{\partial^0 D_t^+} \langle \chi, \frac{\partial}{\partial x^2} \cdot r\psi_r \rangle &= -\frac{1}{2} \int_{-t}^t \langle x^1 \frac{\partial}{\partial x^2} \cdot \chi, \tilde{\nabla}_{\frac{\partial}{\partial x^1}} \psi \rangle dx^1 \\
&= -\frac{1}{2} \int_{-t}^t \frac{\partial}{\partial x^1} \langle x^1 \frac{\partial}{\partial x^2} \cdot \chi, \psi \rangle dx^1 + \frac{1}{2} \int_{-t}^t \langle \tilde{\nabla}_{\frac{\partial}{\partial x^1}} (x^1 \frac{\partial}{\partial x^2} \cdot \chi), \psi \rangle dx^1 \\
&\leq Ct(|\psi|(t, 0) + |\psi|(-t, 0)) + C\sqrt{t}\|\psi\|_{L^2(\partial^0 D_{\frac{1}{2}}^+)}.
\end{aligned}$$

By Hölder's inequality and trace theory, we have

$$\begin{aligned}
\frac{1}{2} \int_{\partial^0 D_t^+} \langle \psi, \frac{\partial}{\partial x^2} \cdot r\chi_r \rangle - \frac{1}{2} \int_{\partial^0 D_t^+} \langle \chi, \frac{\partial}{\partial x^2} \cdot r\chi_r \rangle &\leq C(\sqrt{t}\|\psi\|_{L^2(\partial^0 D_{\frac{1}{2}}^+)} + t) \\
&\leq C(\sqrt{t}\|\psi\|_{W^{1, \frac{4}{3}}(D_{\frac{1}{2}}^+)} + t),
\end{aligned}$$

where C is a constant depending only on $\|\chi\|_{C^1}$.

Then (2.17) implies

$$(2.18) \quad \mathbb{I}_5 \leq Ct(|\psi|(t, 0) + |\psi|(-t, 0)) + C\sqrt{t}\|\psi\|_{L^2(\partial^0 D_{\frac{1}{2}}^+)} + Ct.$$

For \mathbb{I}_7 and \mathbb{I}_8 , it is easy to see that

$$(2.19) \quad \mathbb{I}_7 + \mathbb{I}_8 \leq Ct.$$

Multiplying (2.14) by $\frac{1}{t}$ and integrating from t to $2t$, we get

$$\begin{aligned}
& \int_{D_{2t}^+ \setminus D_t^+} (|\phi_r|^2 - \frac{1}{2}|\nabla\phi|^2) dx \\
& \leq \frac{1}{2} \int_t^{2t} \frac{1}{r} \int_{\partial^+ D_r^+} \langle \psi, r^{-1} \partial_\theta \cdot \psi_\theta \rangle d\theta dr + \int_t^{2t} \frac{1}{r} \int_{\partial^+ D_r^+} r \phi_r \varphi_r d\theta dr \\
& \quad + C \int_t^{2t} (|\psi|(r, 0) + |\psi|(-r, 0) + 1 + \frac{1}{\sqrt{r}}) dr \\
& \leq C \|\psi\|_{L^4(D_{2t}^+ \setminus D_t^+)} \|r^{-1} \psi_\theta\|_{L^{\frac{4}{3}}(D_{2t}^+ \setminus D_t^+)} + Ct(1 + \|d\phi\|_{L^2(D_{2t}^+ \setminus D_t^+)}) + C\sqrt{t}(1 + \|\psi\|_{L^2(\partial^0 D_{\frac{1}{2}}^+)}) \\
& \leq \varepsilon \int_{D_{2t}^+ \setminus D_t^+} \left| r^{-1} \frac{\partial\phi}{\partial\theta} \right|^2 dx + \frac{C}{\varepsilon} \int_{D_{2t}^+ \setminus D_t^+} |\psi|^4 dx + C \int_{D_{2t}^+ \setminus D_t^+} \left| r^{-1} \frac{\partial\psi}{\partial\theta} \right|^{\frac{4}{3}} dx + C\sqrt{t},
\end{aligned}$$

where the last inequality follows from Young's inequality, the trace theory

$$\|\psi\|_{L^2(\partial^0 D_{\frac{1}{2}}^+)} \leq C \|\psi\|_{W^{1, \frac{4}{3}}(D_{\frac{1}{2}}^+)}$$

and the fact

$$\psi_\theta = \tilde{\nabla}_{\frac{\partial}{\partial\theta}} \psi = \frac{\partial\psi}{\partial\theta} + \psi^i \otimes A(d\phi(\frac{\partial}{\partial\theta}), \frac{\partial}{\partial y^i}).$$

This finishes the proof. \square

In the end of this section, we recall some known results for (approximate) Dirac-harmonic maps which are used in this paper.

Theorem 2.6 (Theorem 2.1., [14]). *There is a small constant $\epsilon'_0 > 0$ depending on p, q and N , such that if $(\phi, \psi) \in W^{2,p}(D, N) \times W^{1,q}(D, \Sigma D \otimes \phi^* TN)$ is an approximate Dirac-harmonic map from the unit disc D in \mathbb{R}^2 to a compact Riemannian manifold (N, g) with $\tau(\phi, \psi) \in L^p$ and $h(\phi, \psi) \in L^q$ for some $\frac{4}{3} \leq p \leq 2$ and some $\frac{8}{5} \leq q \leq 2$, and satisfies*

$$(2.20) \quad E(\phi, \psi; D) = \int_D (|d\phi|^2 + |\psi|^4) dx < (\epsilon'_0)^2,$$

then

$$\begin{aligned}
\|\phi - \bar{\phi}\|_{W^{2,p}(D_{\frac{1}{2}})} & \leq C(\|d\phi\|_{L^2(D)} + \|\tau\|_{L^p(D)}), \\
\|\psi\|_{W^{1,q}(D_{\frac{1}{2}})} & \leq C(\|\psi\|_{L^4(D)} + \|h\|_{L^q(D)}),
\end{aligned}$$

where $\bar{\phi} := \frac{1}{|D_{1/2}|} \int_{D_{1/2}} \phi dx$ and $C > 0$ is a constant depending only on p, q, N .

Moreover, by the Sobolev embedding $W^{2,p}(\mathbb{R}^2) \subset C^0(\mathbb{R}^2)$, we have

$$(2.21) \quad \|\phi\|_{\text{Osc}(D_{1/2})} = \sup_{x,y \in D_{1/2}} |\phi(x) - \phi(y)| \leq C(\Lambda, N)(\|\nabla\phi\|_{L^2(D)} + \|\tau(u)\|_{L^p(D)}).$$

Proposition 2.7 (Theorem 3.1 in [1]). *There exists an $\epsilon_1 > 0$ depending on N such that if (ϕ, ψ) is a smooth Dirac-harmonic map from the standard sphere S^2 to a compact Riemannian manifold N satisfying*

$$\int_{S^2} (|d\phi|^2 + |\psi|^4) dx < \epsilon_1,$$

then ϕ is a constant map and $\psi \equiv 0$.

Theorem 2.8 (Theorem 1.4 in [11]). *Let $(\phi, \psi) : \mathbb{R}_+^2 \rightarrow N$ be a smooth Dirac-harmonic map with boundary data $\phi|_{\partial\mathbb{R}_+^2} = \text{const.}$ and $\mathcal{B}\psi|_{\partial\mathbb{R}_+^2} = 0$ and satisfying*

$$\int_{\mathbb{R}_+^2} |\nabla\phi|^2 dx + \int_{\mathbb{R}_+^2} |\psi|^4 dx < \infty.$$

Then ϕ is a constant map and $\psi \equiv 0$.

3. ENERGY IDENTITY

In this section, we will prove our main Theorem 1.1. Since the interior blow-up behavior was already studied in [14], we only need to consider the boundary blow-up behavior.

Firstly, we consider the following simpler case of a boundary blow-up point.

Theorem 3.1. *Let $\phi_n \in C^2(D_1^+(0), N)$, $\psi_n \in C^1(D_1^+(0), \Sigma D_1^+(0) \otimes \phi_n^* TN)$ be a sequence of approximate Dirac-harmonic maps satisfying*

$$(a) \quad \|\phi_n\|_{W^{1,2}(D^+)} + \|\psi_n\|_{L^4(D^+)} + \|\tau_n\|_{L^2(D^+)} + \|h_n\|_{L^4(D^+)} \leq \Lambda,$$

$$(b) \quad (\phi_n, \psi_n) \rightharpoonup (\phi, \psi) \text{ weakly in } W_{loc}^{2,2}(D^+ \setminus \{0\}) \times W_{loc}^{1,2}(D^+ \setminus \{0\}) \text{ as } n \rightarrow \infty.$$

Then there exist a subsequence of (ϕ_n, ψ_n) (still denoted by (ϕ_n, ψ_n)) and a nonnegative integer L such that, for any $i = 1, \dots, L$, there exist points x_n^i , positive numbers λ_n^i and a nonconstant Dirac-harmonic sphere $(\sigma^i, \xi^i) : S^2 \rightarrow N$ such that:

$$(1) \quad x_n^i \rightarrow 0, \lambda_n^i \rightarrow 0, \text{ as } n \rightarrow \infty;$$

$$(2) \quad \frac{\text{dist}(x_n^i, \partial^0 D^+)}{\lambda_n^i} \rightarrow \infty, \text{ as } n \rightarrow \infty;$$

$$(3) \quad \lim_{n \rightarrow \infty} \left(\frac{\lambda_n^i}{\lambda_n^j} + \frac{\lambda_n^j}{\lambda_n^i} + \frac{|x_n^i - x_n^j|}{\lambda_n^i + \lambda_n^j} \right) = \infty \text{ for any } i \neq j;$$

$$(4) \quad (\sigma^i, \xi^i) \text{ is the weak limit of } (\phi_n(x_n^i + \lambda_n^i x), \psi_n(x_n^i + \lambda_n^i x)) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2) \times L_{loc}^4(\mathbb{R}^2);$$

(5) **Energy identity:** we have

$$(3.1) \quad \lim_{n \rightarrow \infty} E(\phi_n) = E(\phi) + \sum_{i=1}^L E(\sigma^i),$$

$$(3.2) \quad \lim_{n \rightarrow \infty} E(\psi_n) = E(\psi) + \sum_{i=1}^L E(\xi^i).$$

Proof. By assumption, without loss of generality, we may assume that 0 is the only blow-up point of the sequence $\{(\phi_n, \psi_n)\}$ in D^+ , i.e.

$$(3.3) \quad \liminf_{n \rightarrow \infty} E(\phi_n, \psi_n; D_r^+) \geq \frac{\bar{\epsilon}^2}{2} \text{ for all } r > 0$$

where $\bar{\epsilon} = \min\{\epsilon_0, \epsilon'_0\}$ and ϵ_0, ϵ'_0 are the constants in Theorem 2.1 and Theorem 2.6 by taking suitable p and q . By the standard argument of blow-up analysis (see e.g. [6, 1]), we can assume that, for any n , there exist sequences $x_n \rightarrow 0$ and $r_n \rightarrow 0$ such that

$$(3.4) \quad E(\phi_n, \psi_n; D_{r_n}^+(x_n)) = \sup_{\substack{x \in D^+, r \leq r_n \\ D_r^+(x) \subset D^+}} E(\phi_n, \psi_n; D_r^+(x)) = \frac{\bar{\epsilon}^2}{4}.$$

Firstly, we make a **Claim 1:** $\limsup_{n \rightarrow \infty} \frac{\text{dist}(x_n, \partial^0 D^+)}{r_n} = \infty$.

If not, after taking a subsequence, we may assume $\lim_{n \rightarrow \infty} \frac{\text{dist}(x_n, \partial^0 D^+)}{r_n} = a \geq 0$. Set

$$u_n(x, t) := \phi_n(x_n + r_n x), \quad v_n(x, t) := \sqrt{r_n} \psi_n(x_n + r_n x),$$

and

$$B_n := \{x \in \mathbb{R}^2 \mid x_n + r_n x \in D^+\}.$$

Then

$$B_n \rightarrow \mathbb{R}^2 := \{(x^1, x^2) \mid x^2 \geq -a\},$$

as $n \rightarrow \infty$. It is easy to see (u_n, v_n) lives in B_n and satisfies

$$(3.5) \quad \begin{cases} r_n^2 \tau(\phi_n) = \Delta u_n + A(du_n, du_n) - \text{Re}(P(\mathcal{A}(du_n(e_\alpha), e_\alpha \cdot v_n); v_n)), & \text{in } B_n; \\ r_n^{\frac{4}{3}} h_n = \not\partial v_n - \mathcal{A}(du_n(e_\alpha), e_\alpha \cdot v_n), & \text{in } B_n, \end{cases}$$

with the boundary data

$$(3.6) \quad \begin{cases} u_n(x) = \varphi(x_1 + r_n x), & \text{if } x_n + r_n x \in \partial M; \\ \mathcal{B}v_n(x) = \sqrt{r_n} \mathcal{B}\chi(x_n + r_n x), & \text{if } x_n + r_n x \in \partial M. \end{cases}$$

By (3.4), Theorem 2.6 and Theorem 2.1, we have

$$(3.7) \quad \|u_n\|_{W^{2,2}(D_{4R}(0) \cap B_n)} + \|v_n\|_{W^{1,2}(D_{4R}(0) \cap B_n)} \leq C(R, N)$$

for any $D_R(0) \subset \mathbb{R}^2$ which implies

$$\|u_n(x - (0, \frac{d_n}{r_n}))\|_{W^{2,2}(D_{3R}^+(0))} + \|v_n(x - (0, \frac{d_n}{r_n}))\|_{W^{1,2}(D_{3R}^+(0))} \leq C(R, N)$$

when n, R are large, where $d_n := \text{dist}(x_n, \partial^0 D^+)$.

Then there exist a subsequence of (u_n, v_n) (also denoted by (u_n, v_n)) and a Dirac-harmonic map $(\tilde{u}, \tilde{v}) \in W^{2,2}(\mathbb{R}_+^2) \times W^{1,2}(\mathbb{R}_+^2)$ with the boundary data $(\tilde{u}, \mathcal{B}\tilde{v})|_{\partial\mathbb{R}_+^2} = (\varphi(x_0), 0)$, such that for any $R > 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n(x - (0, \frac{d_n}{r_n})) - \tilde{u}(x)\|_{W^{1,2}(D_{3R}^+(0))} &= 0 \\ \lim_{n \rightarrow \infty} \|v_n(x - (0, \frac{d_n}{r_n})) - \tilde{v}(x)\|_{L^4(D_{3R}^+(0))} &= 0. \end{aligned}$$

Set $\tilde{u}^1(x) := \tilde{u}(x + (0, a))$ and $\tilde{v}^1(x) := \tilde{v}(x + (0, a))$, then we get, for any $R > 0$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n(x) - \tilde{u}^1(x)\|_{W^{1,2}(D_{2R}(0) \cap B_n \cap \mathbb{R}_a^2)} &= 0 \\ \lim_{n \rightarrow \infty} \|v_n(x) - \tilde{v}^1(x)\|_{L^4(D_{2R}(0) \cap B_n \cap \mathbb{R}_a^2)} &= 0. \end{aligned}$$

Combining this with (3.7) and noting that the measure of $D_{2R}(0) \cap B_n \setminus \mathbb{R}_a^2$ goes to zero, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n(x)\|_{W^{1,2}(D_R(0) \cap B_n)} &= \|\tilde{u}^1(x)\|_{W^{1,2}(D_R(0) \cap \mathbb{R}_a^2)} \\ \lim_{n \rightarrow \infty} \|v_n(x)\|_{L^4(D_R(0) \cap B_n)} &= \|\tilde{v}^1(x)\|_{L^4(D_R(0) \cap \mathbb{R}_a^2)}. \end{aligned}$$

Therefore, by (3.4), we can obtain $E(\tilde{u}^1, \tilde{v}^1; D_1(0) \cap \mathbb{R}_a^2) = \frac{\bar{\epsilon}^2}{4}$. However, by Theorem 2.8, we know \tilde{u}^1 is a constant map and $\tilde{v}^1 \equiv 0$. This is a contradiction. We proved **Claim 1**.

Under the assumption $\limsup_{n \rightarrow \infty} \frac{\text{dist}(x_n, \partial^0 D^+)}{r_n} = \infty$, we can see that (u_n, v_n) lives in B_n which tends to \mathbb{R}^2 as $n \rightarrow \infty$. Moreover, for any $x \in \mathbb{R}^2$, when n is sufficiently large, by (3.4), we have

$$(3.8) \quad E(u_n, v_n; D_1(x)) \leq \frac{\bar{\epsilon}^2}{4}.$$

According to Theorem 2.6, there exist a subsequence of (u_n, v_n) (we still denote it by (u_n, v_n)) and a Dirac-harmonic map $(u^1(x), v^1(x)) \in W^{2,2}(\mathbb{R}^2, N) \times W^{1,2}(\mathbb{R}^2, \Sigma\mathbb{R}^2 \otimes (u^1)^*TN)$ such that

$$(3.9) \quad u_n(x) \rightarrow u^1(x) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2), \quad v_n(x) \rightarrow v^1(x) \text{ in } L_{loc}^4(\mathbb{R}^2),$$

as $n \rightarrow \infty$. Besides, we know $E(u^1, v^1; D_1(0)) = \frac{\bar{\epsilon}^2}{4}$. By the standard theory of Dirac-harmonic maps [1], $(u^1(x), v^1(x))$ can be extended to a nontrivial Dirac-harmonic sphere which is usually called the first bubble.

By the standard induction argument in [6], we only need to prove the theorem in the case where there is only one bubble. For the more bubbles case, i.e. the bubble tree, we just need to distinguish ‘‘neck domains’’ which is almost the same as in the blow-up theory of approximate harmonic maps. See [17, 18] for details. Then we can estimate the energy concentration on each ‘‘neck domain’’ by using the proof of the one bubble case.

Under this assumption, we have the following:

Claim 2: for any $\epsilon > 0$, there exist $\delta > 0$ and $R > 0$ such that

$$(3.10) \quad E(\phi_n, \psi_n; D_{8t}^+(x_n) \setminus D_t^+(x_n)) \leq \epsilon^2 \text{ for any } t \in \left(\frac{1}{2}r_n R, 2\delta\right)$$

when n is large enough.

In fact, if (3.10) is not true, then we can find $t_n \rightarrow 0$, such that $\lim_{n \rightarrow \infty} \frac{t_n}{r_n} = \infty$ and $\epsilon' > 0$ such that

$$(3.11) \quad E(\phi_n, \psi_n; D_{8t_n}^+(x_n) \setminus D_{t_n}^+(x_n)) \geq \epsilon' > 0.$$

Passing to a subsequence, we may assume $\lim_{n \rightarrow \infty} \frac{d_n}{t_n} = b \in [0, \infty]$. For simplicity of notation, we also denote

$$u_n(x) := \phi_n(x_n + t_n x), v_n(x) := \sqrt{t_n} \psi_n(x_n + t_n x).$$

Denoting $B'_n := \{x \in \mathbb{R}^2 | x_n + t_n x \in D^+\}$, then it is easy to see that $(u_n(x), v_n(x))$ lives in B'_n and 0 is also an energy concentration point for (u_n, v_n) . We have to consider the following two cases:

(a) $b < \infty$.

Then B'_n tends to \mathbb{R}_b^2 as $n \rightarrow \infty$. Here, we also need to consider two cases.

(a-1) (u_n, v_n) has no other energy concentration points except 0.

By Theorem 2.6, Theorem 2.1 and the proof of **Claim 1**, there exists a Dirac-harmonic map $(u, v) : \mathbb{R}_b^2 \rightarrow N$ with boundary data $u|_{\partial \mathbb{R}_b^2} = \varphi(0)$, $\mathcal{B}v|_{\partial \mathbb{R}_b^2} = 0$ satisfying, passing to a subsequence, for any $\lambda, R > 0$, there hold

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n(x) - u(x)\|_{W^{1,2}(D_{2R}(0) \cap B'_n \cap \mathbb{R}_b^2 \setminus D_\lambda(0))} &= 0 \\ \lim_{n \rightarrow \infty} \|v_n(x) - v(x)\|_{L^4(D_{2R}(0) \cap B'_n \cap \mathbb{R}_b^2 \setminus D_\lambda(0))} &= 0, \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} \|u_n(x)\|_{W^{1,2}(D_R(0) \cap B'_n \setminus D_\lambda(0))} &= \|u(x)\|_{W^{1,2}(D_R(0) \cap \mathbb{R}_b^2 \setminus D_\lambda(0))} \\ \lim_{n \rightarrow \infty} \|v_n(x)\|_{L^4(D_R(0) \cap B'_n \setminus D_\lambda(0))} &= \|v(x)\|_{L^4(D_R(0) \cap \mathbb{R}_b^2 \setminus D_\lambda(0))}. \end{aligned}$$

According to (3.11), we have

$$E(u, v; (D_8(0) \setminus D_1(0)) \cap \mathbb{R}_b^2) = \lim_{n \rightarrow \infty} E(u_n, v_n; (D_8(0) \setminus D_1(0)) \cap B'_n) \geq \epsilon'.$$

However, Theorem 2.8 tells us that u is a constant map and $v \equiv 0$. This is a contradiction.

(a-2) (u_n, v_n) has another energy concentration point $p \neq 0$.

Without loss of generality, we may assume p is the only energy concentration point in $D_{r_0}(p)$ for some $r_0 > 0$. By the standard argument of blow-up analysis, there exist sequences $x'_n \rightarrow p$ and $r'_n \rightarrow 0$ such that

$$(3.12) \quad E(u_n, v_n; D_{r'_n}(x'_n) \cap B'_n) = \sup_{\substack{x \in D_{r_0}(p), r \leq r'_n \\ D_r(x) \subset D_{r_0}(p)}} E(u_n, v_n; D_r(x) \cap B'_n) = \frac{\bar{\epsilon}^2}{4}.$$

By (3.4), we have $r'_n t_n \geq r_n$ and taking a subsequence, we may assume

$$\lim_{n \rightarrow \infty} \frac{d_n}{r'_n t_n} = d \in [0, \infty].$$

Furthermore, we know d must be ∞ (the proof is the same as for **Claim 1**). Then similar to the process of constructing the first bubble, there exists a nontrivial Dirac-harmonic map $(u^2(x), v^2(x)) \in W^{2,2}(\mathbb{R}^2, N) \times W^{1,2}(\mathbb{R}^2, \Sigma\mathbb{R}^2 \otimes (u^2)^*TN)$ such that

$$u_n(x'_n + r'_n x) \rightarrow u^2(x) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2), \quad \sqrt{r'_n} v_n(x'_n + r'_n x) \rightarrow v^2(x) \text{ in } L_{loc}^4(\mathbb{R}^2),$$

as $n \rightarrow \infty$. This is

$$\phi_n(x_n + t_n x'_n + t_n r'_n x) \rightarrow v^2(x) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2) \text{ and } \sqrt{t_n r'_n} \psi_n(x_n + t_n x'_n + t_n r'_n x) \rightarrow v^2(x) \text{ in } L_{loc}^4(\mathbb{R}^2).$$

Thus, (u^2, v^2) is also a bubble for the sequence (ϕ_n, ψ_n) . This is a contradiction to the one bubble assumption.

(b) $b = \infty$.

In this case, B'_n will tend to \mathbb{R}^2 as $n \rightarrow \infty$. Again, we need to consider the following two cases.

(b-1) (u_n, v_n) has no other energy concentration points except 0.

According to (3.11), Theorem 2.6, Theorem 2.1 and the process of constructing the first bubble, we know that there exists a nontrivial Dirac-harmonic map $(u^2, v^2) : \mathbb{R}^2 \rightarrow N$ such that, passing to a subsequence,

$$u_n(x) \rightarrow u^2(x) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2 \setminus \{0\}) \text{ and } v_n(x) \rightarrow v^2(x) \text{ in } L_{loc}^4(\mathbb{R}^2 \setminus \{0\}),$$

as $n \rightarrow \infty$. Then, we get the second bubble $(u^2(x), v^2(x))$ which contradicts the ‘‘one bubble’’ assumption.

(b-2) (u_n, v_n) has another energy concentration point $p \neq 0$.

Similar to **Case (a-2)**, there exist sequences $x'_n \rightarrow p$ and $r'_n \rightarrow 0$ satisfying (3.12) and

$$\lim_{n \rightarrow \infty} \frac{d_n}{r'_n t_n} = \infty.$$

Moreover, by the process of constructing the first bubble, there exists a nontrivial Dirac-harmonic map $(u^2, v^2) : \mathbb{R}^2 \rightarrow N$ such that, as $n \rightarrow \infty$,

$$u_n(x'_n + r'_n x) \rightarrow v^2(x) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2) \text{ and } \sqrt{r'_n} v_n(x'_n + r'_n x) \rightarrow v^2(x) \text{ in } L_{loc}^4(\mathbb{R}^2)$$

that is

$$\phi_n(x_n + t_n x'_n + t_n r'_n x) \rightarrow v^2(x) \text{ in } W_{loc}^{1,2}(\mathbb{R}^2) \text{ and } \sqrt{t_n r'_n} \psi_n(x_n + t_n x'_n + t_n r'_n x) \rightarrow v^2(x) \text{ in } L_{loc}^4(\mathbb{R}^2).$$

So, we get the second bubble $(u^2(x), v^2(x))$. This also contradicts the “one bubble” assumption. Thus, we proved **Claim 2**.

Under the “one bubble” assumption, by (3.9), it is easy to see that energy identity (3.1) and (3.2) are equivalent to

$$(3.13) \quad \lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E(\phi_n; D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)) = 0$$

and

$$(3.14) \quad \lim_{R \rightarrow \infty} \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} E(\psi_n; D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)) = 0.$$

Without loss of generality, we may assume $\delta = 2^{m_n} r_n R$ for some positive integer m_n which tends to ∞ as $n \rightarrow \infty$. We denote $P_i := D_{2^i r_n R}^+(x_n) \setminus D_{2^{i-1} r_n R}^+(x_n)$.

Firstly we use a finite decomposition argument that is similar to those in [29, 30] to separate $\Sigma := D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)$ into finite parts

$$\Sigma = \cup_{j=1}^{s_n} Q_j, \quad Q_j := \cup_{i=k_{j-1}+1}^{k_j} P_i, \quad 0 = k_0 < k_1 < \dots < k_{s_n} = m_n$$

such that $s_n \leq S$ and

$$(3.15) \quad E(\phi_n, \psi_n; Q_j) \leq \frac{1}{C_1(N)}, \quad j = 1, \dots, s_n,$$

where $C_1(N) > 0$ is a constant depending only on N to be determined later and S is a uniform integer for all n large enough.

From (3.10), for any $\epsilon < \frac{1}{2C_1(N)}$, we have

$$E(\phi_n, \psi_n; P_i) < \epsilon < \frac{1}{2C_1(N)}, \quad i = 1, \dots, m_n$$

when n is large.

If

$$E(\phi_n, \psi_n; \Sigma) \leq \frac{1}{C_1(N)},$$

let $k_1 = m_n$ and then $Q_1 = \Sigma$. Otherwise, we can choose an integer $1 \leq k_1 < m_n$ such that

$$\frac{1}{2C_1(N)} < E(\phi_n, \psi_n; Q_1) \leq \frac{1}{C_1(N)} \quad \text{and} \quad E(\phi_n, \psi_n; Q_1 \cup P_{k_1+1}) > \frac{1}{C_1(N)}.$$

This is the first step of the division. Inductively, suppose that k_j is chosen such that $E(\phi_n, \psi_n; Q_j) \leq \frac{1}{C_1(N)}$. If

$$E(\phi_n, \psi_n; \cup_{i=k_j+1}^{m_n} P_i) \leq \frac{1}{C_1(N)},$$

let $k_{j+1} = m_n$, thus $Q_{j+1} = \cup_{i=k_{j+1}}^{m_n} P_i$. If not, then similar to the first step, we can find $k_j < k_{j+1} < m_n$ such that

$$\frac{1}{2C_1(N)} < E(\phi_n, \psi_n; Q_{j+1}) \leq \frac{1}{C_1(N)} \quad \text{and} \quad E(\phi_n, \psi_n; Q_{j+1} \cup P_{k_{j+1}+1}) > \frac{1}{C_1(N)}.$$

Since $E(\phi_n, \psi_n)$ is uniformly bounded by Λ , we will finish our division after at most $S = [2C_1(N)\Lambda] + 1$ steps. So we have finished the division.

Take a cut-off function $\eta \in C_0^\infty(D_{2^{k_j+1}r_nR}^+(x_n) \setminus D_{2^{k_j-1}r_nR}^+(x_n))$ such that $0 \leq \eta \leq 1$ and $\eta|_{D_{2^{k_j}r_nR}^+(x_n) \setminus D_{2^{k_j-1}r_nR}^+(x_n)} \equiv 1$ and

$$\begin{aligned} |\nabla\eta| &\leq \frac{C}{2^{k_j}r_nR} \quad \text{on} \quad D_{2^{k_j+1}r_nR}^+(x_n) \setminus D_{2^{k_j}r_nR}^+(x_n) \quad \text{and} \\ |\nabla\eta| &\leq \frac{C}{2^{k_j-1}r_nR} \quad \text{on} \quad D_{2^{k_j-1}r_nR}^+(x_n) \setminus D_{2^{k_j-2}r_nR}^+(x_n). \end{aligned}$$

By the standard elliptic estimates, we have

$$\begin{aligned} &\|\eta\psi_n\|_{W^{1,4/3}(D_1^+)} \\ &\leq C\|\eta\phi_n + \nabla\eta \cdot \psi_n\|_{L^{\frac{4}{3}}(D_1^+)} + C\|\eta\mathcal{B}\chi\|_{W^{1,4,4/3}(\partial D_1^+)} \\ &\leq \frac{1}{4}C(N)(\|d\phi_n\|_{L^2(\Sigma)}\|\eta\psi_n\|_{L^{\frac{4}{3}}(\Sigma)} + \|\eta|h_n|\|_{L^{\frac{4}{3}}(\Sigma)}) + C\|\nabla\eta\|_{L^{\frac{4}{3}}(\Sigma)}\|\psi_n\|_{L^{\frac{4}{3}}(\Sigma)} + C\|\eta\mathcal{B}\chi\|_{W^{1,4,4/3}(\partial^0 D_\delta^+)} \\ &\leq \frac{1}{4}C(N)\|d\phi_n\|_{L^2(D_{2^{k_j+1}r_nR}^+(x_n) \setminus D_{2^{k_j-1}r_nR}^+(x_n))}\|\eta\psi_n\|_{L^4(\Sigma)} + C\|h_n\|_{L^{\frac{4}{3}}(\Sigma)} \\ &\quad + C\|\nabla\eta\psi_n\|_{L^{\frac{4}{3}}(P_{k_j-1} \cup P_{k_j+1})} + C\|\eta\mathcal{B}\chi\|_{W^{1,4,4/3}(\partial^0 D_\delta^+)} \\ &\leq \frac{1}{4}C(N)\frac{2}{\sqrt{C_1(N)}}\|\eta\psi_n\|_{L^4(\Sigma)} + C\|\psi_n\|_{L^4(P_{k_j-1} \cup P_{k_j+1})} + C\|h_n\|_{L^{\frac{4}{3}}(\Sigma)} + C\|\eta\mathcal{B}\chi\|_{W^{1,4,4/3}(\partial^0 D_\delta^+)}, \end{aligned}$$

where the last inequality is from (3.10) and (3.15). Then, taking $C_1(N) = C^2(N) + 1$, by (3.10) and Sobolev embedding, we have

$$\begin{aligned} \|\psi_n\|_{L^4(Q_j)} + \|\nabla\psi_n\|_{L^{4/3}(Q_j)} &\leq C\|\psi_n\|_{L^4(P_{k_j-1} \cup P_{k_j+1})} + C\|h_n\|_{L^{\frac{4}{3}}(\Sigma)} + C\|\eta\mathcal{B}\chi\|_{W^{1,4,4/3}(\partial^0 D_\delta^+)} \\ &\leq C(\Lambda, \|\chi\|_{C^1})(\sqrt{\epsilon} + \delta). \end{aligned}$$

So,

$$\|\psi_n\|_{L^4(\Sigma)} + \|\nabla\psi_n\|_{L^{4/3}(\Sigma)} \leq \sum_{j=1}^{s_n} (\|\psi_n\|_{L^4(Q_j)} + \|\nabla\psi_n\|_{L^{4/3}(Q_j)}) \leq CS(\sqrt{\epsilon} + \delta).$$

This is (3.14).

Suppose $x'_n \in \partial^0 D^+$ is the projection of x_n , i.e. $d_n = \text{dist}(x_n, \partial^0 D^+) = |x_n - x'_n|$. Similar to the boundary blow-up cases for approximate harmonic maps studied in [12, 13], we

decompose the neck domain $D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n)$ as follows

$$\begin{aligned} D_\delta^+(x_n) \setminus D_{r_n R}^+(x_n) &= D_\delta^+(x_n) \setminus D_{\frac{\delta}{2}}^+(x'_n) \cup D_{\frac{\delta}{2}}^+(x'_n) \setminus D_{2d_n}^+(x'_n) \\ &\quad \cup D_{2d_n}^+(x'_n) \setminus D_{d_n}^+(x_n) \cup D_{d_n}^+(x_n) \setminus D_{r_n R}^+(x_n) \\ &:= \Omega_1 \cup \Omega_2 \cup \Omega_3 \cup \Omega_4, \end{aligned}$$

when n is large.

Since $\lim_{n \rightarrow \infty} d_n = 0$ and $\lim_{n \rightarrow \infty} \frac{d_n}{r_n} = \infty$, when n is large enough, it is easy to see that

$$\Omega_1 \subset D_\delta^+(x_n) \setminus D_{\frac{\delta}{4}}^+(x_n), \quad \text{and} \quad \Omega_3 \subset D_{4d_n}^+(x_n) \setminus D_{d_n}^+(x_n).$$

Moreover, for any $d_n \leq t \leq \delta$, there holds

$$D_{2t}^+(x'_n) \setminus D_t^+(x'_n) \subset D_{4t}^+(x_n) \setminus D_{t/2}^+(x_n).$$

By assumption (3.10), we have

$$(3.16) \quad E(\phi_n, \psi_n; \Omega_1) + E(\phi_n, \psi_n; \Omega_3) \leq \epsilon^2$$

and

$$(3.17) \quad E(\phi_n, \psi_n; D_{2t}^+(x'_n) \setminus D_t^+(x'_n)) \leq \epsilon^2 \text{ for any } t \in (d_n, \delta).$$

By (3.10), Theorem 2.6, Theorem 2.1 and the standard scaling argument, we get

$$\begin{aligned} &Osc_{D_{2t}^+(x'_n) \setminus D_t^+(x'_n)} \phi_n \\ &\leq C(\|\nabla \phi_n\|_{L^2(D_{4t}^+(x'_n) \setminus D_{t/2}^+(x'_n))} + \|\psi_n\|_{L^4(D_{4t}^+(x'_n) \setminus D_{t/2}^+(x'_n))} + \|\nabla \varphi\|_{L^2(D_{4t}^+(x'_n) \setminus D_{t/2}^+(x'_n))} \\ &\quad + t\|\nabla^2 \varphi\|_{L^2(D_{4t}^+(x'_n) \setminus D_{t/2}^+(x'_n))} + t\|\tau(u_n)\|_{L^2(D_{4t}^+(x'_n) \setminus D_{t/2}^+(x'_n))}) \\ (3.18) \quad &\leq C(\sqrt{\epsilon} + \delta), \end{aligned}$$

for any $t \in (2r_n R, \frac{1}{2}\delta)$, where $C = C(\Lambda, N, \|\varphi\|_{C^2}, \|\chi\|_{C^1})$ is a positive constant.

Noting that $\Omega_4 = D_{d_n}^+(x_n) \setminus D_{r_n R}^+(x_n) = D_{d_n}(x_n) \setminus D_{r_n R}(x_n)$, by the energy identity of approximate Dirac-harmonic maps with interior blow-up points (see Theorem 1.2 in [14]), there holds

$$(3.19) \quad \lim_{R \rightarrow \infty} \lim_{n \rightarrow 0} E(u_n; D_{d_n}(x_n) \setminus D_{r_n R}(x_n)) = 0.$$

Therefore, we just need to estimate the energy concentration in Ω_2 . Here, we use a similar method as in [12, 13].

Define $\widehat{\Omega}_2 := D_{\frac{\delta}{2}}^+(x'_n) \setminus D_{2d_n}(x'_n)$, $\Phi_n(x) := \phi_n(x) - \varphi(x)$, $x \in \Omega_2$ and

$$(3.20) \quad \widehat{\Phi}_n(x) := \begin{cases} \Phi_n(x), & x \in \Omega_2, \\ -\Phi_n(x'), & x \in \widehat{\Omega}_2 \setminus \Omega_2, \end{cases}$$

where $x = (x^1, x^2)$ and $x' = (x^1, -x^2)$. It is easy to see that $\widehat{\Phi}_n(x) \in W^{2,\infty}(\widehat{\Omega}_2)$ and satisfies the following equation

$$(3.21) \quad \Delta \widehat{\Phi}_n(x) = \begin{cases} \Delta \phi_n(x) - \Delta \varphi(x), & x \in \Omega_2, \\ -\Delta \phi_n(x') + \Delta \varphi(x'), & x \in \widehat{\Omega}_2 \setminus \Omega_2, \end{cases}$$

where $\Delta\phi_n(x) = -A(d\phi_n, d\phi_n)(x) + \text{Re}(P(\mathcal{A}(d\phi_n(e_\alpha), e_\alpha \cdot \psi_n); \psi_n))(x) + \tau(u_n)(x)$.

Without loss of generality, we may also assume $\frac{1}{2}\delta = 2^{m'_n}(2d_n)$, where m'_n is a positive integer which tends to ∞ as $n \rightarrow \infty$. Setting $P'_i := D_{2^{i+1}d_n}^+(x'_n) \setminus D_{2^i d_n}^+(x'_n)$ and $\widehat{P}'_i := D_{2^{i+1}d_n}(x'_n) \setminus D_{2^i d_n}(x'_n)$,

Set

$$\widehat{\Phi}_n^*(r) := \frac{1}{2\pi} \int_0^{2\pi} \widehat{\Phi}_n(r, \theta) d\theta,$$

where (r, θ) are the polar coordinates at x'_n . By (3.18) and (3.20), we have

$$\begin{aligned} \|\widehat{\Phi}_n(x) - \widehat{\Phi}_n^*(x)\|_{L^\infty(\widehat{\Omega}_2)} &\leq \sup_{1 \leq i \leq m'_n} \|\widehat{\Phi}_n(x) - \widehat{\Phi}_n^*(x)\|_{L^\infty(\widehat{P}'_i)} \leq \sup_{1 \leq i \leq m'_n} \|\widehat{\Phi}_n(x)\|_{\text{Osc}(\widehat{P}'_i)} \\ &\leq 2 \sup_{1 \leq i \leq m'_n} \|\Phi_n(x)\|_{\text{Osc}(P'_i)} \leq 2 \sup_{1 \leq i \leq m'_n} \|\phi_n(x)\|_{\text{Osc}(P'_i)} + C\delta \|\nabla\varphi\|_{L^\infty} \\ (3.22) \quad &\leq C(N, \Lambda, \|\varphi\|_{C^2}, \|\chi\|_{C^1})(\sqrt{\epsilon} + \delta). \end{aligned}$$

Integrating by parts, we get

$$\int_{\widehat{P}'_i} \nabla \widehat{\Phi}_n \nabla (\widehat{\Phi}_n - \widehat{\Phi}_n^*) dx = \int_{\partial \widehat{P}'_i} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} - \int_{\widehat{P}'_i} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \Delta \widehat{\Phi}_n dx.$$

On the one hand, we have

$$\begin{aligned} \int_{\widehat{P}'_i} \nabla \widehat{\Phi}_n \nabla (\widehat{\Phi}_n - \widehat{\Phi}_n^*) dx &= \int_{\widehat{P}'_i} |\nabla \widehat{\Phi}_n|^2 dx - \int_{\widehat{P}'_i} \frac{\partial \widehat{\Phi}_n}{\partial r} \frac{\partial \widehat{\Phi}_n^*}{\partial r} dx \\ &\geq \int_{\widehat{P}'_i} |\nabla \widehat{\Phi}_n|^2 dx - \left(\int_{\widehat{P}'_i} \left| \frac{\partial \widehat{\Phi}_n}{\partial r} \right|^2 dx \right)^{\frac{1}{2}} \left(\int_{\widehat{P}'_i} \left| \frac{\partial \widehat{\Phi}_n^*}{\partial r} \right|^2 dx \right)^{\frac{1}{2}} \\ &\geq \int_{\widehat{P}'_i} |\nabla \widehat{\Phi}_n|^2 dx - \int_{\widehat{P}'_i} \left| \frac{\partial \widehat{\Phi}_n}{\partial r} \right|^2 dx \\ &= \frac{1}{2} \int_{\widehat{P}'_i} |\nabla \widehat{\Phi}_n|^2 dx - \int_{\widehat{P}'_i} \left(\left| \frac{\partial \widehat{\Phi}_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla \widehat{\Phi}_n|^2 \right) dx \\ &= \int_{\widehat{P}'_i} |\nabla \Phi_n|^2 dx - 2 \int_{\widehat{P}'_i} \left(\left| \frac{\partial \Phi_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla \Phi_n|^2 \right) dx. \end{aligned}$$

By direct computation, we obtain

$$\begin{aligned} &\int_{P'_i} |\nabla \Phi_n|^2 dx - 2 \int_{P'_i} \left(\left| \frac{\partial \Phi_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla \Phi_n|^2 \right) dx \\ &= \int_{P'_i} |\nabla \phi_n|^2 dx - 2 \int_{P'_i} \left(\left| \frac{\partial \phi_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla \phi_n|^2 \right) dx + 4 \int_{P'_i} \left(\frac{\partial \phi_n}{\partial r} \frac{\partial \varphi}{\partial r} - \nabla \phi_n \nabla \varphi \right) dx \\ &\quad + 2 \int_{P'_i} \left(|\nabla \varphi|^2 - \left| \frac{\partial \varphi}{\partial r} \right|^2 \right) dx \\ &\geq \int_{P'_i} |\nabla \phi_n|^2 dx - 2 \int_{P'_i} \left(\left| \frac{\partial \phi_n}{\partial r} \right|^2 - \frac{1}{2} |\nabla \phi_n|^2 \right) dx - C2^i d_n. \end{aligned}$$

On the other hand, by (3.21) and (3.22), we have

$$\begin{aligned} \int_{\widehat{P}'_i} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \Delta \widehat{\Phi}_n dx &\leq C(\sqrt{\epsilon} + \delta) \int_{P'_i} |d\phi_n|^2 dx + C(\sqrt{\epsilon} + \delta) \int_{P'_i} |d\phi| |\psi_n|^2 dx \\ &\quad + C(\sqrt{\epsilon} + \delta) \int_{P'_i} (|\tau_n| + |\Delta\varphi|) dx \\ &\leq C(\sqrt{\epsilon} + \delta) \int_{P'_i} |d\phi_n|^2 dx + C(\sqrt{\epsilon} + \delta) \int_{P'_i} |\psi_n|^4 dx + C(\sqrt{\epsilon} + \delta) 2^i d_n. \end{aligned}$$

From the above, by Corollary 2.5 (taking $\epsilon = \frac{1}{2}$), we get

$$\begin{aligned} \int_{P'_i} |d\phi_n|^2 dx &\leq \int_{\partial \widehat{P}'_i} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} + \int_{P'_i} (|\frac{\partial \phi_n}{\partial r}|^2 - \frac{1}{2} |\nabla \phi_n|^2) dx + C(\sqrt{\epsilon} + \delta) \int_{P'_i} |d\phi_n|^2 dx \\ &\quad + C(\sqrt{\epsilon} + \delta) \int_{P'_i} |\psi_n|^4 dx + C 2^i d_n \\ &\leq \int_{\partial \widehat{P}'_i} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} + (\frac{1}{2} + C(\sqrt{\epsilon} + \delta)) \int_{P'_i} |d\phi_n|^2 dx + C \int_{P'_i} |\psi_n|^4 dx \\ &\quad + C \int_{P'_i} |\nabla \psi_n|^{\frac{4}{3}} dx + C 2^i d_n. \end{aligned}$$

Summing i from 1 to m'_n , we get

$$\begin{aligned} (\frac{1}{2} - C(\sqrt{\epsilon} + \delta)) \int_{\Omega_2} |\nabla \phi_n|^2 dx &\leq \int_{\partial D_{\delta/2}(x'_n)} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} - \int_{\partial D_{2d_n}(x'_n)} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} \\ (3.23) \quad &\quad + C \int_{\Omega_2} |\nabla \psi_n|^{4/3} dx + C \int_{\Omega_2} |\psi_n|^4 dx + C\delta. \end{aligned}$$

As for the boundary term, by trace theory, we have

$$\begin{aligned} \int_{\partial D_{\delta/2}(x'_n)} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} &\leq C(\sqrt{\epsilon} + \delta) \int_{\partial D_{\delta/2}(x'_n)} |\nabla \widehat{\Phi}_n| \\ &\leq C(\sqrt{\epsilon} + \delta) \int_{\partial^+ D_{\delta/2}(x'_n)} (|\nabla \phi_n| + |\nabla \varphi|) \\ &\leq C(\sqrt{\epsilon} + \delta) \left(\|\nabla \phi_n\|_{L^2(D_{\delta}^+ \setminus D_{\frac{1}{4}\delta}^+)} + \delta \|\nabla^2 \phi_n\|_{L^2(D_{\delta}^+ \setminus D_{\frac{1}{4}\delta}^+)} + 1 \right) \\ &\leq C(\sqrt{\epsilon} + \delta) \left(\|\nabla \phi_n\|_{L^2(D_{\frac{4}{3}\delta}^+ \setminus D_{\frac{1}{6}\delta}^+)} + \|\psi_n\|_{L^4(D_{\frac{4}{3}\delta}^+ \setminus D_{\frac{1}{6}\delta}^+)} \right. \\ &\quad \left. + \|\nabla \varphi\|_{L^2(D_{\frac{4}{3}\delta}^+ \setminus D_{\frac{1}{6}\delta}^+)} + \delta \|\nabla^2 \varphi\|_{L^2(D_{\frac{4}{3}\delta}^+ \setminus D_{\frac{1}{6}\delta}^+)} \right. \\ &\quad \left. + \delta \|\tau_n\|_{L^2(D_{\frac{4}{3}\delta}^+ \setminus D_{\frac{1}{6}\delta}^+)} + 1 \right) \\ &\leq C(\sqrt{\epsilon} + \delta), \end{aligned}$$

where the last second inequality can be derived from Theorem 2.1.

Also, there holds

$$\int_{\partial D_{2d_n}} (\widehat{\Phi}_n - \widehat{\Phi}_n^*) \frac{\partial \widehat{\Phi}_n}{\partial r} \leq C(\sqrt{\epsilon} + \delta).$$

Putting these in (3.23) and taking ϵ and δ sufficient small, we have

$$(3.24) \quad \int_{\Omega_2} |\nabla \phi_n|^2 dx \leq C \int_{\Omega_2} |\nabla \psi_n|^{4/3} dx + C \int_{\Omega_2} |\psi_n|^4 dx + C(\sqrt{\epsilon} + \delta).$$

Combining this with (3.16), (3.19) and (3.14), we will obtain (3.13) and we finished the proof of Theorem 3.1. \square

Proof of Theorem 1.1: It is easy to see that Theorem 1.1 is a consequence of the interior blow-up case, i.e. Theorem 1.2 in [14] and the model case of boundary blow-ups, i.e. Theorem 3.1. \square

REFERENCES

- [1] Q. Chen, J. Jost, J. Li and G. Wang, *Regularity theorems and energy identities for Dirac-harmonic maps*, *Mathematische Zeitschrift* **251** (2005), no. 1, 61–84.
- [2] Q. Chen, J. Jost, J. Li and G. Wang, *Dirac-harmonic maps*, *Mathematische Zeitschrift* **254** (2006), no. 2, 409–432.
- [3] Q. Chen, J. Jost, L. Sun and M. Zhu, *Estimates for solutions of Dirac equations and an application to a geometric elliptic-parabolic problem*, MPI MIS Preprint: 79/2014.
- [4] Q. Chen, J. Jost, G. Wang and M. Zhu, *The boundary value problem for Dirac-harmonic maps*, *J. Eur. Math. Soc. (JEMS)*. Volume 15, Issue 3, 2013, 997–1031.
- [5] P. Deligne, *Quantum fields and strings: a course for mathematicians*, vol. 2, Amer Mathematical Society, 1999.
- [6] W. Ding and G. Tian, *Energy identity for a class of approximate harmonic maps from surfaces*, *Comm. Anal. Geom.* **3** (1995), no. 3-4, 543–554.
- [7] T. Huang and C. Wang, *Boundary bubbling analysis of approximate harmonic maps under either weak or strong anchoring conditions in dimension two*, arXiv:1607.05357.
- [8] J. Jost, *Two-dimensional geometric variational problems*, New York, Wiley, 1991.
- [9] J. Jost, *Geometry and physics*, Springer, 2009
- [10] J. Jost, L. Liu and M. Zhu, *Geometric analysis of the action functional of the nonlinear supersymmetric sigma model*, MPI MIS Preprint: 77/2015.
- [11] J. Jost, L. Liu and M. Zhu, *A global weak solution of the Dirac-harmonic map flow*, MPI MIS Preprint: 1/2016. To appear in *Ann. Inst. H. Poincaré Anal. Non Linéaire* (2017). <https://doi.org/10.1016/j.anihpc.2017.01.002>
- [12] J. Jost, L. Liu and M. Zhu, *The qualitative behavior at the free boundary for approximate harmonic maps from surfaces*, MPI MIS Preprint: 26/2016.
- [13] J. Jost, L. Liu and M. Zhu, *Bubbling analysis near the Dirichlet boundary for approximate harmonic maps from surfaces*, MPI MIS Preprint: 38/2016. To appear in *Comm. Anal. Geom.* (2017).
- [14] J. Jost, L. Liu and M. Zhu, *Blow-up analysis for approximate Dirac-harmonic maps in dimension 2 with applications to Dirac-harmonic heat flow*, MPI MIS Preprint: 65/2016. To appear in *Calc. Var. Partial Differ. Equ.* (2017).
- [15] H. Lawson and M. Michelsohn, *Spin geometry*, vol. 38, Princeton University Press, 1989.

- [16] J. Li and X. Zhu, *Energy identity for the maps from a surface with tension field bounded in L^p* , Pacific Journal of Mathematics 260 (2012), no. 1, 181-195.
- [17] Y. Li and Y. Wang, *A weak energy identity and the length of necks for a sequence of Sacks-Uhlenbeck α -harmonic maps*, Adv. Math. 225 (2010), no. 3, 1134-1184.
- [18] J. Chen and Y. Li, *Homotopy classes of harmonic maps of the stratified 2-spheres and applications to geometric flows*, Adv. Math. 263 (2014), 357-388.
- [19] F. Lin and C. Wang, *Energy identity of harmonic map flow from surfaces at finite singular time*, Calculus of Variations and Partial Differential Equations, 6 (1998), 369-380.
- [20] F. Lin and C. Wang, *Harmonic and quasi-harmonic spheres. II*. Comm. Anal. Geom. 10 (2002), no. 2, 341-375.
- [21] L. Liu, *No neck for Dirac-harmonic maps*, Calculus of Variations and Partial Differential Equations 52 (2015), no. 1-2, 1-15.
- [22] T. Parker, *Bubble tree convergence for harmonic maps*, J. Diff. Geom 44 (1996), no. 3, 595-633.
- [23] J. Qing, *On singularities of the heat flow for harmonic maps from surfaces into spheres*. Comm. Anal. Geom. 3 (1995), 297-315.
- [24] J. Qing and G. Tian, *Bubbling of the heat flows for harmonic maps from surfaces*, Communications on pure and applied mathematics 50 (1997), no. 4, 295-310.
- [25] J. Sacks and K. Uhlenbeck, *The existence of minimal immersions of 2-spheres*, Ann. Math.113 (1981), 1-24.
- [26] B. Sharp and M. Zhu, *Regularity at the free boundary for Dirac-harmonic maps from surfaces*, Calc. Var. Partial Differ. Equ. 55 (2016), no. 2, 55:27.
- [27] C. Wang, *Remarks on approximate harmonic maps in dimension two*, Calc. Var. Partial Differential Equations 56 (2017), no. 2, 56:23, 24 pp.
- [28] W. Wang, D. Wei and Z. Zhang, *Energy identity for approximate harmonic maps from surface to general targets*, J. Funct. Anal. 272 (2017), no. 2, 776-803.
- [29] R. Ye, *Gromov's compactness theorem for pseudo-holomorphic curves*, Trans. Am. Math. Soc. 342(2), 671-694 (1994).
- [30] L. Zhao, *Energy identities for Dirac-harmonic maps*, Calc. Var. Partial Differential Equations 28 (2007), no. 1, 121-138.
- [31] M. Zhu, *Dirac-harmonic maps from degenerating spin surfaces I: the Neveu-Schwarz case*, Calc. Var. Partial Differ. Equ. 35 (2009), no. 2, 169-189.
- [32] M. Zhu, *Regularity of weakly Dirac-harmonic maps to hypersurfaces*, Ann. Global Anal. Geom. 35 (2009), no.4 405-412.

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, 04103 LEIPZIG, GERMANY

DEPARTMENT OF MATHEMATICS, LEIPZIG UNIVERSITY, 04081 LEIPZIG, GERMANY
E-mail address: jost@mis.mpg.de

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, 04103 LEIPZIG, GERMANY

E-mail address: leiliu@mis.mpg.de or llei1988@mail.ustc.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, 800 DONGCHUAN ROAD, SHANGHAI, 200240, CHINA

E-mail address: mizhu@sjtu.edu.cn