

**Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig**

Two-copy Quantum Teleportation

by

*Quan Quan, Ming-Jing Zhao, Shao-Ming Fei,
Heng Fan, and Gui-Lu Long*

Preprint no.: 3

2019



Two-copy Quantum Teleportation

Quan Quan^{1,2}, Ming-Jing Zhao³, Shao-Ming Fei^{4,5,*}, Heng Fan^{6,7,2}, Wen-Li Yang^{2,8,*} and Gui-Lu Long¹

¹*State Key Laboratory of Low-Dimensional Quantum Physics and*

Department of Physics, Tsinghua University, Beijing 100084, China

²*Institute of Modern Physics, Northwest University, Xi'an 710069, China*

³*School of Science, Beijing Information Science and Technology University, Beijing, 100192, China*

⁴*School of Mathematical Sciences, Capital Normal University, Beijing 100048, China*

⁵*Max-Planck-Institute for Mathematics in the Science, Leipzig 04103, Germany*

⁶*Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*

⁷*Collaborative Innovation Center of Quantum Matter, Beijing 100190, China*

⁸*Shaanxi Key Laboratory for Theoretical Physics Frontiers, Xi'an 710069, China*

We investigate two-copy scenario of quantum teleportation based on Bell measurements. The detailed protocol is presented and the general expression of the corresponding optimal teleportation fidelity is derived, which is given by the two-copy fully entangled fraction that is invariant under local unitary transformations. We prove that under a specific case of the protocol, which is significant for improving the optimal fidelity, the set of states with their two-copy fully entangled fractions bounded by a threshold value that required for useful two-copy teleportation is convex and compact. Hence the witness operators exist to separate states that are useful for two-copy teleportation from the rest ones. Moreover, we show that the fidelity of two-copy optimal teleportation surpasses that of the original one copy teleportation.

PACS numbers: 03.67.-a, 03.65.Ud, 03.65.Ta

INTRODUCTION

Quantum teleportation plays an important role in quantum information processing [1]. It gives ways to transmit an unknown quantum state from a sender traditionally named “Alice” to a receiver “Bob” who are spatially separated, using classical communication and quantum resources [2–5]. In [6, 7], the authors consider the original one copy teleportation: Alice and Bob previously share a pair of particles in an arbitrary mixed entangled state χ . In order to teleport an unknown state to Bob, Alice first performs joint Bell measurements on her particles and tells the results to Bob by classical communication. Bob tries his best to choose particular unitary transformations to get the optimal transmission fidelity. The optimal transmission fidelity of such teleportation is given by the fully entangled fraction (FEF) [8] of the quantum resource. It shows that when the resource χ is a maximally entangled pure state, the corresponding optimal fidelity is equal to 1. However, Alice and Bob usually share a mixed entangled state due to decoherence, and the optimal fidelity is less than 1.

One way to improve the fidelity of teleportation is to distill entanglement [10], which refers to the procedure of converting mixed entangled states into singlets by using many copies of the entangled resources. The distillation of pure states is often referred to as entanglement concentration [11]. For mixed states, since the distillation protocol presented in [10], fruitful results have been obtained [12–15]. However, the problem of distillation is that the complicated protocol may have to be repeated for infinitely many times to bring out a singlet. Moreover, in each round the desired results are usually ob-

tained probabilistically, usually with an extremely low possibility to get an expected measurement outcome.

Inspired by this, to improve the teleportation fidelity, we propose two-copy quantum teleportation scenario directly, instead of bringing out a singlet as the resource of traditional one copy teleportation scenario. Specifically, we introduce a quantum teleportation protocol based on Bell measurements. The corresponding optimal teleportation fidelity is derived analytically. The fact that the fidelity of the optimal two-copy teleportation can surpass that in one copy scenario (the traditional quantum teleportation) is shown by analytical means together with numerical methods. In particular, we discuss a specific case which is significant for improving the optimal fidelity. It shows that the set of quantum states with their two-copy fully entangled fraction bounded by a threshold value required for useful two-copy teleportation is convex and compact, which demonstrates the existence of teleportation witness in two-copy quantum teleportation.

TWO-COPY TELEPORTATION PROTOCOL BASED ON BELL MEASUREMENT

The two-copy teleportation protocol is as follows. Initially Alice and Bob share two pairs of entangled resources, see Fig. 1. Particles 1 and 2 (resp. 3 and 4) are in an entangled state χ . Particles 1 and 3 are in Alice’s side, while particles 2 and 4 are in Bob’s side. Alice wants to transmit an unknown state ρ_{in} of particle 0 to Bob. Firstly, Alice (resp. Bob) performs a joint local unitary operation W (resp. V) on particles 1 (resp. 2) and particle 3 (resp. 4). Then she makes joint Bell




FIG. 1. Scheme of two-copy teleportation protocol based on Bell measurements. Alice and Bob share two copies of entangled resource χ , with particles 1 and 3 in Alice's side, and particles 2 and 4 in Bob's side. Alice wants to transmit the unknown state ρ_{in} of particle 0 to Bob with optimal fidelity. The two-channel teleportation protocol based on Bell measurements is as follows: firstly, Alice (resp. Bob) performs a joint local unitary operation W (resp. V) on particles 1 (resp. 2) and particle 3 (resp. 4) to correlate these two particles. Then Alice makes joint Bell measurements on particles 0 and 1 and informs Bob the measurement results by classical communication. According to the measurement results, Bob chooses corresponding unitary transformations $\{T\}$ on his particles 2 and 4 to restore the input state ρ_{in} on particle 2.

measurements on particles 0 and 1. She informs Bob the measurement results by classical means. According to these measurement results, Bob chooses corresponding unitary transformations $\{T\}$ on particle 2 to achieve the optimal teleportation fidelity.

Let H denotes an n -dimensional Hilbert space, with $\{|j\rangle, j = 0, \dots, n-1, n < \infty\}$ as orthogonal normalized basis. A set of unitary matrices $\{U_{st}\}$ in H can be written as: $U_{st} = h^t g^s$, where h and g are $n \times n$ matrices such that $h|k\rangle = |(k+1)/\text{mod } n\rangle$ and $g|k\rangle = w^k|k\rangle$, with $w = \exp\{-2i\pi/n\}$. $\{U_{st}\}$ has the following relations [16]: $\text{tr}(U_{st}^\dagger U_{s't'}) = n\delta_{tt'}\delta_{ss'}$, $U_{st}U_{st}^\dagger = I_{n \times n}$, where I is the identity matrix. The generalized Bell states [6] are given by $|\Phi_{st}\rangle = (I_{n \times n} \otimes U_{st})|\Phi\rangle$, where $|\Phi\rangle = |\Phi_{00}\rangle = \frac{1}{\sqrt{n}} \sum_{j=0}^{n-1} |jj\rangle$ is the maximally entangled pure state. The n^2 generalized Bell states $\{|\Phi_{st}\rangle\} = \frac{1}{\sqrt{n}} \sum_{j,k} (U_{st})_{jk}^* |jk\rangle$ form a complete orthogonal normalized basis of the $H \otimes H$ space. Throughout this paper we adopt the standard notations: for any matrix $A \in \text{End}(H)$, A_α is an embedded operator in the tensor space $H \otimes H \otimes \dots \otimes H$, which acts as A on the α -th space and as identity on the other spaces; and for any matrix $U \in \text{End}(H \otimes H)$, $U_{\alpha\beta}$ is an embedded operator in $H \otimes H \otimes \dots \otimes H$, which acts as identity on the spaces except for the α -th and β -th ones. After some tedious calculation, we get the output state under the scenario of two-copy teleportation:

Theorem 1 For any unknown input state ρ_{in} , the two-copy teleportation protocol under Bell measurements

maps the state ρ_{in} to state $\Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in})$,

$$\begin{aligned} & \Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in}) \\ &= \frac{1}{n^3} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \sum_{s, t} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \\ & \text{tr}_4 [(T_{st})_2 V_{24} (U_{s_1 t_1})_2 (U_{s_2 t_2})_4 W_{24} (U_{st})_2^\dagger (\rho_{in})_2 \\ & (U_{st})_2 W_{24}^\dagger (U_{s'_1 t'_1})_2^\dagger (U_{s'_2 t'_2})_4^\dagger V_{24}^\dagger (T_{st})_2^\dagger], \end{aligned} \quad (1)$$

where W and V are the unitary transformations Alice and Bob apply to their two particles, respectively. $T_{st} \in \{T\}$ is the unitary operator that Bob performs on particle 2 to achieve the optimal teleportation fidelity.

Proof. First consider that the unknown initial input state ρ_{in} that Alice wants to teleport is a pure state, $|\phi\rangle = \sum_\nu \alpha_\nu |\nu\rangle$.

1). The two entangled resource states shared by Alice and Bob are pure: $\chi^{\otimes 2} = |\Psi\rangle_{1234} \langle \Psi|$, where $|\Psi\rangle_{1234} = \sum_{j,k=0}^{n-1} \sum_{l,m=0}^{n-1} a_{jk} |jk\rangle \otimes a_{lm} |lm\rangle$, $\sum_{j,k=0}^{n-1} |a_{jk}|^2 = 1$. Alice and Bob apply the unitary transformations W and V to their two resource particles respectively. Before the measurement, the initial state becomes $|\phi\rangle_0 W_{13} V_{24} |\Psi\rangle_{1234} = \sum_{j,k=0}^{n-1} \sum_{l,m=0}^{n-1} \sum_{j',k'=0}^{n-1} \sum_{l',m'=0}^{n-1} \sum_\nu a_{ij} a_{kl} W_{i'k'}^{ik} V_{j'l'}^{jl} \alpha_\nu |\nu i' j' k' l'\rangle_{01234}$.

After Alice's joint Bell measurements based on $|\Phi_{st}\rangle$ on particles 0 and 1, we get: $\langle \Phi_{st} |_{01} (|\phi\rangle_0 W_{13} V_{24} |\Psi\rangle_{1234}) = V_{24} A_2 A_4 W_{24} (U_{st}^\dagger)_2 |\phi\rangle_2 |\Phi\rangle_{34}$, where A is the $n \times n$ matrix with elements $(A)_{jk} = a_{jk}$. Receiving Alice's measurement outcomes, correspondingly Bob applies unitary operators $\{T\}$ on particle 2. The resulting state becomes $(T_{st})_2 V_{24} A_2 A_4 W_{24} (U_{st}^\dagger)_2 |\phi\rangle_2 |\Phi\rangle_{34}$. Taking partial trace over the spaces with respect to particles 3 and 4, we have

$$\begin{aligned} & \Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in}) \\ &= \sum_{s,t} \frac{1}{n} \text{tr}_4 [(T_{st})_2 V_{24} A_2 A_4 W_{24} (U_{st}^\dagger)_2 |\phi\rangle_2 \langle \phi|_2 \\ & (U_{st})_2 W_{24}^\dagger A_2^\dagger A_4^\dagger V_{24}^\dagger (T_{st})_2^\dagger], \end{aligned} \quad (2)$$

which is equivalent to Eq. (1).

2). Now consider the case of arbitrary entangled mixed resources, $\chi^{\otimes 2} = \sum_{\alpha, \beta} P_\alpha P_\beta |\Psi_{\alpha\beta}\rangle \langle \Psi_{\alpha\beta}|$, where $|\Psi_{\alpha\beta}\rangle = \sum_{j,k=0}^{n-1} \sum_{l,m=0}^{n-1} a_{jk}^{(\alpha)} |jk\rangle \otimes a_{lm}^{(\beta)} |lm\rangle$, $0 \leq P_{\alpha(\beta)} \leq 1$ and $\sum_{\alpha(\beta)} P_{\alpha(\beta)} = 1$. Similar to the derivation of (2), we have

$$\begin{aligned} & \Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in}) \\ &= \frac{1}{n} \sum_{s,t} \sum_{\alpha, \beta} P_\alpha P_\beta \text{tr}_4 [(T_{st})_2 V_{24} A_2^{(\alpha)} A_4^{(\beta)} W_{24} (U_{st}^\dagger)_2 |\phi\rangle_2 \langle \phi|_2 \\ & (U_{st})_2 W_{24}^\dagger A_2^{(\alpha)\dagger} A_4^{(\beta)\dagger} V_{24}^\dagger (T_{st})_2^\dagger], \end{aligned}$$

where $(A)_{jk}^{(\alpha/\beta)} = a_{jk}^{(\alpha/\beta)}$. Since each matrix $A^{(\alpha)}$ can be decomposed in the basis of U_{st} : $A^{(\alpha)} = \sum_{s,t} a_{st}^{(\alpha)} U_{st}$, by

using the relation [6], $n \sum_{\alpha} p_{\alpha} a_{st}^{(\alpha)} a_{s't'}^{(\alpha)*} = \langle \Phi_{st} | \chi | \Phi_{s't'} \rangle$, we can straightforwardly show that Eq. (1) is also valid for any mixed input state ρ_{in} . \square

Remark The two-copy teleportation scenario is trace preserving, $\text{tr}[\Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in})] = 1$, see proof in Appendix A.

Utilizing the output state, we get the optimal teleportation fidelity, see Appendix B for detailed proof:

Theorem 2 The optimal teleportation fidelity $f_2(\chi)_{max}$ of the two-copy teleportation protocol is given by

$$f_2(\chi)_{max} = \frac{nF_2(\chi)}{(n+1)} + \frac{1}{n+1}, \quad (3)$$

where $\Omega_{13}^T = W_{13}(U_{st})_1(T_{st})_3$, $F_2(\chi)$ is two-copy fully entangled fraction,

$$F_2(\chi) = \max_{\Omega, V \in U(n^2)} \{ \langle \Phi |_{12} \text{tr}_{34} [\Omega_{13} V_{24} \chi_{12} \chi_{34} \Omega_{13}^{\dagger} V_{24}^{\dagger}] | \Phi \rangle_{12} \}.$$

From Theorem 2 we see that the two-copy optimal teleportation fidelity $f_2(\chi)$ solely depends on the two-copy fully entangled fraction $F_2(\chi)$. It can be shown that $F_2(\chi)$ given by (3) is an invariant under local unitary transformations: $\chi_{12} \chi_{34} \rightarrow (\mathfrak{U})_{13} (\mathfrak{V})_{24} \chi_{12} \chi_{34} (\mathfrak{U})_{13}^{\dagger} (\mathfrak{V})_{24}^{\dagger}$, where \mathfrak{U} and \mathfrak{V} are unitary operators on $H \otimes H$. Theorem 2 also tells us that a resource state χ is useful, namely, it gives better teleportation fidelity than classical channels, if $F_2(\chi) > \frac{1}{n}$.

The original one copy optimal teleportation fidelity is given by [6, 7]

$$f_1(\chi)_{max} = \frac{nF_1(\chi)}{n+1} + \frac{1}{n+1}, \quad (4)$$

where $F_1(\chi) = \max_{U \in U(n)} \{ \langle \Phi |_{12} U_2^{\dagger} \chi_{12} U_2 | \Phi \rangle_{12} \}$ is the original fully entangled fraction. To show that the two-copy teleportation protocol is always better, or at least as good as the original one copy case, let us simply choose the unitary matrix $W_{13} = V_{24} = I_{n^2 \times n^2}$ in Theorem 2, which does not necessarily reach the value of $f_2(\chi)_{max}$. Then the output state $\Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in})$ of the two-copy teleportation protocol reduces to the output state $\Lambda^{(x)}(\{T\})(\rho)$ of one-copy teleportation protocol [6, 7], which can be seen from the proof of Theorem 1. Therefore, we have the following relation between the two-copy optimal teleportation fidelity and the original one-copy optimal teleportation fidelity,

$$f_2(\chi)_{max} \geq f_1(\chi)_{max}. \quad (5)$$

From Eq. (3) and Eq. (4), one can see that both optimal teleportation fidelities for two-copy and one-copy teleportation protocols are linear functions of the corresponding fully entangled fractions. These fully entangled fractions characterize the usefulness of the entangled resource states in quantum teleportation. To compare $f_2(\chi)_{max}$ with $f_1(\chi)_{max}$, one only needs to compare




FIG. 2. Hollow (solid) triangles stand for 3 (4)-dimensional randomly generated states. Horizontal axis is the one-copy fully entangled fraction F_1 . Vertical axis denotes the differences $\Delta F = F'_2 - F_1$. It is seen that $\Delta F > 0$, and hence the lower bound of the optimal fidelity of the two-copy teleportation is better than the optimal fidelity of original one-copy teleportation for all these randomly generated states.

$F_2(\chi)$ with $F_1(\chi)$. Unfortunately, both $F_2(\chi)$ and $F_1(\chi)$ are formidably difficult to calculate analytically. Analytical formulae for $F_1(\chi)$ are only available for some special states [17, 18]. Generally one has only estimations of the upper and lower bounds of $F_1(\chi)$ [18, 19]. The computation of $F_2(\chi)$ is much more difficult than that of $F_1(\chi)$. However, if one takes $W = V$ to be identity, or takes Ω and V in (3) to be the tensor of two unitary operators $\Upsilon \otimes \Gamma$ with $\Upsilon, \Gamma \in U(n)$, then one gets $F_2(\chi) = F_1(\chi)$. Thus the extreme value range of F_2 is larger than that of F_1 . Therefore, for any arbitrary state χ , $f_2(\chi)_{max} \geq f_1(\chi)_{max}$, i.e., the two-copy optimal teleportation fidelity is always greater than or equal to that of the original one copy protocol.

In the following, we give numerical calculations of F_2 and F_1 by using the Conjugate Gradient Algorithm [20, 21]. Following the modified Polak-Ribiere-Polyak method introduced in [21], we can get the numerical result of F_1 . For F_2 , to simplify the computation, we take $V = I_{n \times n}$. Then we get a lower bound $F'_2(\chi)$ of $F_2(\chi)$:

$$F'_2(\chi) = \max_{\Omega \in U(n^2)} \left\{ \sum_j \langle \Phi |_{12} \langle j |_3 \Omega_{23} \chi_{12} \Omega_{23}^{\dagger} \rho_3^* | j \rangle_3 | \Phi \rangle_{12} \right\},$$

where $\rho_3 = \text{tr}_4(\chi_{34})$; ρ_3^* is the conjugate of ρ_3 , see Appendix D. Denote $\mathfrak{F}_2(\chi) = \sum_j \langle \Phi |_{12} \langle j |_3 \Omega_{23} \chi_{12} \Omega_{23}^{\dagger} \rho_3^* | j \rangle_3 | \Phi \rangle_{12}$. We get $F'_2(\chi) = \max_{\Omega_{23} \in U(n^2)} \mathfrak{F}_2$. Set $\Delta F = F'_2(\chi) - F_1(\chi)$. Fig. 2 shows that for these randomly generated states, one has $F'_2(\chi) > F_1(\chi)$, i.e., the lower bound of the optimal fidelity of two-copy teleportation is better than the optimal fidelity of the one-copy teleportation.

Let us investigate further the lower bound $F'_2(\chi)$ of the two-channel fully entangled fraction $F_2(\chi)$, obtained by setting $V = I_{n \times n}$ in the two-copy teleportation protocol. For the original one copy teleportation, it is shown that

the resource states χ satisfying $F_1(\chi) > 1/n$ are useful for teleportation [22]. Since the set of states satisfying $F_1(\chi) \leq \frac{1}{n}$ is convex and compact, there exist witness operators that detect entangled states which are useful for teleportation [22, 23]. Similarly, for the two-copy teleportation protocol with $V = I_{n \times n}$, if resource states satisfying $F'_2 > \frac{1}{n}$, they are useful for teleportation. Denote $\mathbb{S} = \{\chi : F'_2(\chi) \leq \frac{1}{n}\}$. We have:

Theorem 3 The set \mathbb{S} is convex and compact. For the two-copy teleportation protocol with $V = I_{n \times n}$, there exist witnesses to identify the usefulness of an unknown resource states experimentally.

Proof. For the proof of that set \mathbb{S} is convex and compact, see Appendix E. According to the Hahn-Banach theorem [24], any $\chi \notin \mathbb{S}$ can be separated from \mathbb{S} by a hyperplane. This feature enables for the existence of hermitian witness operators and thus experimental ways to detect the usefulness of an unknown resource state. \square

CONCLUSIONS AND DISCUSSIONS

To conclude, we have proposed a general two-copy quantum teleportation protocol based on Bell measurement systematically. The corresponding optimal teleportation fidelities have been analytically derived. Interestingly, the formulae of optimal teleportation fidelity in two-copy scenario and one-copy scenario are similar. Both of them are the one-variable linear function of the corresponding fully entangled fractions, which are invariant under local unitary transformations on the resource states. Analytical analysis together with numerical results illustrate that the optimal teleportation fidelity can be improved in two-copy teleportation protocol when compared with the one-copy scenario. Therefore, in order to improve the teleportation fidelity, two-copy teleportation protocol is significant both theoretically and experimentally. Furthermore, we have shown that in the context of two-copy teleportation protocol, if one considers a specific case that Bob conducts identity transformation on his resource states, the optimal fidelity can be still be improved. The quantum states with their two-copy fully entangled fraction bounded by a threshold value required for useful two-copy teleportation is convex and compact. It demonstrates the existence of teleportation witness, completing the theory of two-copy quantum teleportation. Besides, this two-copy quantum teleportation protocol can be generalized to many-copy cases and result in further improvement on the teleportation fidelity. Different protocols and methods are worthwhile to conceive and investigate in the future.

ACKNOWLEDGEMENTS

Q. Q. is grateful to Prof. Kai Chen for discussion and to Huangjun Zhu and Xiaolong Du for help on numerical algorithm. We gratefully thank for the supports by NSFC (Grant Nos. 11275131, 11375141, 11434013, 11425522) and the “973” program (2010CB922904). Q. Q. acknowledges financial supports from NSFC (Grant No. 11704217). M. J. Zhao acknowledges financial supports from NSFC (Grant No. 11401032).

-
- *
- [1] Nielsen, M. A. & Chuang, I. L. *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
 - [2] Bennett, C. H. et al. Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels. *Phys. Rev. Lett.* **70**, 1895 (1993).
 - [3] D’Ariano, G. M., Lo Presti, P. & Sacchi, M. F. Bell measurements and observables. *Phys. Lett. A* **272**, 32 (2000).
 - [4] Bowen, G. & Bose, S. Teleportation as a Depolarizing Quantum Channel, Relative Entropy, and Classical Capacity. *Phys. Rev. Lett.* **87**, 267901 (2001).
 - [5] Alberverio, S., Fei, S. M. & Yang, W. L., Teleportation with an Arbitrary Mixed Resource as a Trace-Preserving Quantum Channel. *Commun. Theor. Phys.* **38**, 301-304 (2002).
 - [6] Alberverio, S., Fei, S. M. & Yang, W. L., teleportation based on bell measurements. *Phys. Rev. A* **66**, 012301 (2002).
 - [7] Horodecki, M., Horodecki, P. & Horodecki, R., *Phys. Rev. A* **60** 1888 (1999).
 - [8] Alber, G. et al. *Quantum Information: An Introduction to Basic Concepts and Experiments* (Springer Tracts in Modern Physics, 2001).
 - [9] Abeyesinghe, A., Devetak, I., Hayden, P. & Winter, A. The mother of all protocols: restructuring quantum informations family tree. *Proc. R. Soc. A* **465(2108)**: 2537-2563 (2009).
 - [10] Bennett, C. H. et al., Purification of Noisy Entanglement and Faithful Teleportation via Noisy Channels. *Phys. Rev. Lett.* **76**, 722-725 (1996).
 - [11] Bennett, C. H., Bernstein, H. J., Popescu, S. & Schumacher, B. Concentrating partial entanglement by local operations. *Phys. Rev. A* **53**, 2046 (1996).
 - [12] Horodecki, M., Horodecki, P. & Horodecki, R. Inseparable Two Spin-12 Density Matrices Can Be Distilled to a Singlet Form. *Phys. Rev. Lett.* **78** 574 (1997).
 - [13] Horodecki, M., Horodecki, P. & Horodecki, R. Mixed-State Entanglement and Distillation: Is there a Bound Entanglement in Nature? *Phys. Rev. Lett.* **80** 5239 (1998).
 - [14] Peres, A. Separability Criterion for Density Matrices. *Phys. Rev. Lett.* **77** 1413 (1996).
 - [15] Horodecki, R., Horodecki, P., Horodecki, M. & Horodecki, K. Quantum entanglement. *Rev. Mod. Phys.* **81**, 865 (2009).
 - [16] Werner, R. F. All teleportation and dense coding schemes. *J. Phys. A: Math. Gen.* **34** 7081(2001).

- [17] Grondalski, J., Etlinger, D. M. & James, D. F. V. The fully entangled fraction as an inclusive measure of entanglement applications. *Phys. Lett. A* **300**, 573 (2002).
- [18] Zhao, M. J., Li, Z. G., Fei, S. M. & Wang, Z. X. A Note on Fully Entangled Fraction. *J. Phys. A* **43**, Math. Theor. 275203 (2010).
- [19] Li, M., Fei, S. M. & Wang, Z. X. Upper bound of the fully entangled fraction. *Phys. Rev. A* **78**, 032332 (2008).
- [20] Shrwchuk, J. R. *An Introduction to the Conjugate Gradient Method Without the Agonizing Pain*, (August 1994).
- [21] Zhang, Li., Zhou, W. J. & Li, D. H. A descent modified PolakCRibi'ereCPolyak conjugate gradient method and its global convergence. *Journal of Numerical analysis* **26** 629-640 (2006).
- [22] Ganguly, N., Adhikari, S., Majumdar, A. S. & Chatterjee, J. Entanglement Witness Operator for Quantum Teleportation. *Phys. Rev. Lett.* **107**, 270501 (2011).
- [23] Zhao, M. J., Fei, S. M. & Li-Jost, X. Complete entanglement witness for quantum teleportation. *Phys. Rev. A* **85**, 054301 (2012).
- [24] Holmes, R. B. *Geometric Functional Analysis and its Applications*, (Springer-Verlag 1975).

Appendix A. Proof of the Remark

The output state $\Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}$ of the two-copy teleportation is trace preserving:

$$\begin{aligned}
& \text{tr}[\Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in})] \\
&= \frac{1}{n^3} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \\
& \quad \text{tr}_2 \{ (T_{st})_2 (U_{s_1 t_1})_2 \text{tr}_4 [(U_{s_2 t_2})_4 W_{24} \\
& \quad W_{24}^\dagger (U_{s'_2 t'_2})_4^\dagger] (U_{s'_1 t'_1})_2 (T_{st})_2^\dagger \} \\
&= \frac{1}{n^2} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \\
& \quad \text{tr}_2 [(U_{s_1 t_1})_2 (U_{s'_1 t'_1})_2] \text{tr}_4 [(U_{s_2 t_2})_4 (U_{s'_2 t'_2})_4^\dagger] \\
&= 1,
\end{aligned}$$

where in the first equality we have used the relation $\sum_{s,t} U_{st}^\dagger A U_{st} = n \text{tr}(A) I_{n \times n}$ for any $n \times n$ matrix A .

Appendix B. Proof of Theorem 2

Let $U(n)$ be an irreducible n -dimensional representation of unitary group G . By using the Schur's lemma

$$\begin{aligned}
& \int_G dg (U^\dagger(g) \otimes U^\dagger(g)) \sigma (U(g) \otimes U(g)) = \alpha_1 I \otimes I + \alpha_2 P, \\
& \alpha_1 = \frac{n^2 \text{tr}(\sigma) - n \text{tr}(\sigma P)}{n^2(n^2 - 1)}, \quad \alpha_2 = \frac{n^2 \text{tr}(\sigma P) - n \text{tr}(\sigma)}{n^2(n^2 - 1)},
\end{aligned}$$

where σ is any operator acting on the tensor space, P is the flip operator, dg is the Haar measure on G normalized by $\int_G dg = 1$, we get the fidelity of the two-copy

teleportation protocol,

$$\begin{aligned}
& f_2(\chi) = \overline{\langle \phi_{in} | \Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in}) | \phi_{in} \rangle} \\
&= \frac{1}{n^3} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \\
& \quad \sum_{s, t, j, k} \langle 00 | \int_G [U(g)^\dagger \otimes U(g)^\dagger] [|j\rangle_4 (T_{st})_2 V_{24} (U_{s_1 t_1})_2 \\
& \quad (U_{s_2 t_2})_4 W_{24} (U_{st})_2^\dagger |k\rangle_4] \otimes [\langle k |_4 (U_{st})_2 W_{24}^\dagger (U_{s'_1 t'_1})_2 \\
& \quad (U_{s'_2 t'_2})_4 V_{24}^\dagger (T_{st})_2^\dagger |j\rangle_4] [U(g) \otimes U(g)] dg |00\rangle \\
&= \frac{1}{n^4(n+1)} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \\
& \quad \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \sum_{s, t, j, k, l, l'} \{ \text{tr}_2 [\langle j |_4 (T_{st})_2 V_{24} (U_{s_1 t_1})_2 \\
& \quad (U_{s_2 t_2})_4 |l\rangle_4 \langle l |_4 W_{24} (U_{st})_2^\dagger |k\rangle_4] \text{tr}_4 [\langle k |_4 (U_{st})_2 W_{24}^\dagger \\
& \quad (U_{s'_1 t'_1})_2 (U_{s'_2 t'_2})_4^\dagger |l'\rangle_4 \langle l' |_4 V_{24}^\dagger (T_{st})_2^\dagger |j\rangle_4] \\
& \quad + \frac{1}{n^2(n+1)} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \\
& \quad \text{tr}_2 [(U_{s_1 t_1})_2 (U_{s'_1 t'_1})_2] \text{tr}_4 [(U_{s_2 t_2})_4 (U_{s'_2 t'_2})_4^\dagger] \\
&= \frac{1}{n^4(n+1)} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \sum_{s, t, j, k} \langle \Phi |_{12} \langle \Phi |_{34} \\
& \quad \text{tr}_{24} [W_{24} (U_{st})_2^\dagger |k\rangle_4 \langle j |_4 (T_{st})_2 V_{24} (U_{s_1 t_1})_2 (U_{s_2 t_2})_4] \\
& \quad (U_{s'_1 t'_1})_2 (U_{s'_2 t'_2})_4^\dagger \chi_{12} \chi_{34} \text{tr}_{24} [V_{24}^\dagger (T_{st})_2^\dagger |j\rangle_4 \langle k |_4 (U_{st})_2 W_{24}^\dagger \\
& \quad (U_{s'_1 t'_1})_2 (U_{s'_2 t'_2})_4^\dagger] (U_{s'_1 t'_1})_2 (U_{s'_2 t'_2})_4 | \Phi \rangle_{12} | \Phi \rangle_{34} + \frac{1}{n+1} \\
&= \frac{1}{(n+1)} \sum_{s, t} \langle \Phi |_{12} \langle \Phi |_{34} W_{24} (U_{st})_2^\dagger (T_{st})_2 \text{tr}_4 [V_{24} \chi_{12} \chi_{34} V_{24}^\dagger] \\
& \quad (T_{st})_2^\dagger (U_{st})_2 W_{24}^\dagger | \Phi \rangle_{12} | \Phi \rangle_{34} + \frac{1}{n+1},
\end{aligned}$$

where $\overline{\langle \phi_{in} | \dots | \phi_{in} \rangle}$ represents the average over all input states $|\phi_{in}\rangle$.

Then the optimal teleportation fidelity is given by the maximal fidelity of $f_2(\chi)$,

$$\begin{aligned}
& f_2(\chi)_{max} = \frac{n^2}{(n+1)} \max_{\Omega, V \in U(n^2)} \{ \langle \Phi |_{12} \langle \Phi |_{34} \Omega_{24} \\
& \quad \text{tr}_4 [V_{24} \chi_{12} \chi_{34} V_{24}^\dagger] \Omega_{24}^\dagger | \Phi \rangle_{12} | \Phi \rangle_{34} \} + \frac{1}{n+1} \\
&= \frac{n^2}{(n+1)} \max_{\Omega, V \in U(n^2)} \{ \langle \Phi |_{12} \langle \Phi |_{34} \Omega_{13}^T \\
& \quad \text{tr}_4 [V_{24} \chi_{12} \chi_{34} V_{24}^\dagger] \Omega_{13}^* | \Phi \rangle_{12} | \Phi \rangle_{34} \} + \frac{1}{n+1},
\end{aligned}$$

where $\Omega_{24} = W_{24}(U_{st})_2(T_{st})_2$. Rewriting Ω^T as Ω , we get (3). \square

Appendix C. Proof of (5)

When we choose $W_{13} = V_{24} = I_{n^2 \times n^2}$, the output state $\Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in})$ of the two-copy teleportation protocol reduces to that of one-copy teleportation protocol $\Lambda^{(x)}(\{T\})(\rho)$ in [6]:

$$\begin{aligned} & \Lambda_{\chi^{\otimes 2}}^{\{T_{st}, W, V\}}(\rho_{in}) \\ &= \frac{1}{n^3} \sum_{s_1, t_1} \sum_{s'_1, t'_1} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \sum_{s, t} \langle \Phi_{s_1 t_1} | \chi | \Phi_{s'_1 t'_1} \rangle \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \\ & \quad \text{tr}_4[(T_{st})_2 (U_{s_1 t_1})_2 (U_{s_2 t_2})_4 (U_{st})_2^\dagger | \phi \rangle_2 \langle \phi |_2 (U_{st})_2 \\ & \quad (U_{s'_1 t'_1}^\dagger)_2 (U_{s'_2 t'_2}^\dagger)_4 (T_{st})_2^\dagger], \\ &= \Lambda^{(x)}(\{T\})(\rho_{in}) \frac{1}{n} \sum_{s_2, t_2} \sum_{s'_2, t'_2} \langle \Phi_{s_2 t_2} | \chi | \Phi_{s'_2 t'_2} \rangle \text{tr}[U_{s_2 t_2} U_{s'_2 t'_2}^\dagger] \\ &= \Lambda^{(x)}(\{T\})(\rho_{in}). \end{aligned}$$

Appendix D. Derivation of $F'_2(\chi)$

$$\begin{aligned} F'_2(\chi) &= n \max_{\Omega \in U(n^2)} \{ \langle \Phi |_{12} \langle \Phi |_{34} \Omega_{24} \chi_{12} \rho_3 \Omega_{24}^\dagger | \Phi \rangle_{12} | \Phi \rangle_{34} \} \\ &= \max_{\Omega \in U(n^2)} \{ \sum_{j, j'} \langle \Phi |_{12} \langle j |_4 \Omega_{24} \chi_{12} \Omega_{24}^\dagger | j' \rangle_4 | \Phi \rangle_{12} \langle j |_3 \rho_3 | j' \rangle_3 \} \\ &= \max_{\Omega \in U(n^2)} \{ \sum_j \langle \Phi |_{12} \langle j |_3 \Omega_{23} \chi_{12} \Omega_{23}^\dagger \rho_3^* | j \rangle_3 | \Phi \rangle_{12} \}, \end{aligned}$$

where $\rho_3 = \text{tr}_4(\chi_{34})$.

Appendix E. \mathbb{S} is convex and compact

Proof. 1). The set \mathbb{S} is convex: Let χ_a and $\chi_b \in \mathbb{S}$, namely, $F'_2(\chi_a) \leq \frac{1}{n}$, $F'_2(\chi_b) \leq \frac{1}{n}$. Consider $\chi_c = \xi \chi_a + (1 - \xi) \chi_b$, where $\xi \in [0, 1]$. By the definition of $F'_2(\chi) = \max_{\Omega_{23} \in U(n^2)} \{ \sum_j \langle \Phi_{00} |_{12} \langle j |_3 \Omega_{23} \chi_{12} \Omega_{23}^\dagger \rho_3^* | j \rangle_3 | \Phi_{00} \rangle_{12} \}$, we get that $F'_2(\chi_c) \leq \xi F'_2(\chi_a) + (1 - \xi) F'_2(\chi_b) \leq \frac{1}{n}$. Thus $\chi_c \in \mathbb{S}$, i.e. \mathbb{S} is convex.

2). The set \mathbb{S} is compact: For finite dimensional Hilbert spaces, to show that a set is compact, it is enough to show the set is closed and bounded. \mathbb{S} is bounded since the eigenvalues of χ lies in $[0, 1]$ [22]. To see that it is closed, assume that for any two density matrices χ_a and χ_b , the value of $F'_2(\chi_a + \chi_b)$ and $F'_2(\chi_a)$ are obtained at Ω_{a+b} and Ω_a respectively, where $\Omega \in U(n^2)$. Therefore

$$\begin{aligned} & F'_2(\chi_a + \chi_b) - F'_2(\chi_a) \\ & \leq \sum_j \langle \Phi_{00} |_{12} \langle j |_3 (\Omega_{a+b})_{23} (\chi_a)_{12} (\Omega_{a+b})_{23}^\dagger \text{tr}_4[(\chi_b)_{34}]^* | j \rangle_3 | \Phi_{00} \rangle_{12} \\ & \quad + \sum_j \langle \Phi_{00} |_{12} \langle j |_3 (\Omega_{a+b})_{23} (\chi_b)_{12} (\Omega_{a+b})_{23}^\dagger \text{tr}_4[(\chi_a)_{34}]^* | j \rangle_3 | \Phi_{00} \rangle_{12} \\ & \quad + \sum_j \langle \Phi_{00} |_{12} \langle j |_3 (\Omega_{a+b})_{23} (\chi_b)_{12} (\Omega_{a+b})_{23}^\dagger \text{tr}_4[(\chi_b)_{34}]^* | j \rangle_3 | \Phi_{00} \rangle_{12} \\ & \leq \sum_{j, j'} \|\Phi_{00}\|^2 \| |j\rangle \langle j'| \|^2 \|\Omega_{a+b}\|^2 (2\|\chi_a\| + \|\chi_b\|) \|\chi_b\| \end{aligned}$$

where $\|\chi_a\|$ is the maximal eigenvalue of χ_a satisfying $\|\chi_a\| \leq 1$. Since the set of all unitary operators is bounded, $\|\Omega_{a+b}\|^2 \leq v$, where v is a positive real number. Thus $F'_2(\chi_a + \chi_b) - F'_2(\chi_a) \leq n^2 v (2 + \|\chi_b\|) \|\chi_b\|$. \square